

Early Weight Loss Nomograms for Exclusively Breastfed Newborns

Valerie J. Flaherman, MD, MPH^{a,b}, Eric W. Schaefer, MS^c, Michael W. Kuzniewicz, MD, MPH^{a,d}, Sherian X. Li, MS^d, Eileen M. Walsh, MPH, RN^d, Ian M. Paul, MD, MS^{c,e}

abstract

BACKGROUND: The majority of newborns are exclusively breastfed during the birth hospitalization, and weight loss is nearly universal for these neonates. The amount of weight lost varies substantially among newborns with higher amounts of weight loss increasing risk for morbidity. No hour-by-hour newborn weight loss nomogram exists to assist in early identification of those on a trajectory for adverse outcomes.

METHODS: For 161 471 term, singleton neonates born at ≥ 36 weeks' gestation at Northern California Kaiser Permanente hospitals in 2009–2013, data were extracted from the birth hospitalization regarding delivery mode, race/ethnicity, feeding type, and weights from electronic records. Quantile regression was used to create nomograms stratified by delivery mode that estimated percentiles of weight loss as a function of time among exclusively breastfed neonates. Weights measured subsequent to any nonbreastmilk feeding were excluded.

RESULTS: Among this sample, 108 907 newborns had weights recorded while exclusively breastfeeding with 83 433 delivered vaginally and 25 474 delivered by cesarean. Differential weight loss by delivery mode was evident 6 hours after delivery and persisted over time. Almost 5% of vaginally delivered newborns and $>10\%$ of those delivered by cesarean had lost $\geq 10\%$ of their birth weight 48 hours after delivery. By 72 hours, $>25\%$ of newborns delivered by cesarean had lost $\geq 10\%$ of their birth weight.

CONCLUSIONS: These newborn weight loss nomograms demonstrate percentiles for weight loss by delivery mode for those who are exclusively breastfed. The nomograms can be used for early identification of neonates on a trajectory for greater weight loss and related morbidities.



WHAT'S KNOWN ON THIS SUBJECT: Exclusively breastfed newborns lose weight daily in the first few days after birth. The amount of weight lost varies substantially between newborns, with higher amounts of weight loss increasing risk for morbidity.

WHAT THIS STUDY ADDS: This study presents nomograms demonstrating percentiles for weight loss by delivery mode for those who are exclusively breastfed. The nomograms have potential to be used for early identification of neonates on a trajectory for greater weight loss and related morbidities.

Departments of ^aPediatrics, and ^bEpidemiology and Biostatistics, School of Medicine, University of California, San Francisco, California; Departments of ^cPublic Health Sciences, and ^dPediatrics, Penn State College of Medicine, Hershey, Pennsylvania; and ^eDivision of Research, Kaiser Permanente, Oakland, California

Dr Flaherman conceptualized and designed this study, oversaw all aspects of the data collection and analysis, and drafted the initial manuscript; Mr Schaefer contributed to the study design, analyzed the data, and critically revised the manuscript for important scientific content; Dr Kuzniewicz contributed to the study design, participated in data analysis, and critically revised the manuscript for important scientific content; Ms Li contributed to the study design, coordinated data collection, carried out preliminary analyses, and critically revised the manuscript for important scientific content; Ms Walsh participated in study design and data analysis and critically revised the manuscript for important scientific content; Dr Paul conceptualized and designed this study, obtained funding, and critically revised the manuscript for important scientific content; and all authors approved this manuscript as submitted.

www.pediatrics.org/cgi/doi/10.1542/peds.2014-1532

DOI: 10.1542/peds.2014-1532

Accepted for publication Oct 3, 2014

The World Health Organization, the Centers for Disease Control and Prevention, and the American Academy of Pediatrics all recommend breastfeeding exclusively during the birth hospitalization without any supplementary formula or water,¹⁻³ and in the United States, 60% of newborns are breastfed exclusively for at least the first 2 days.⁴ Among such exclusively breastfed neonates, enteral intake is low during the time of colostrum production,⁵⁻⁷ and initial postnatal weight loss is nearly universal.⁸⁻¹⁰ Although the brief period of low enteral intake and weight loss is well tolerated by most newborns, some develop complications of weight loss such as hyperbilirubinemia and dehydration, the 2 most common causes of neonatal morbidity.¹¹⁻¹⁷

To monitor for potential morbidity, weight and jaundice are evaluated daily during the birth hospitalization, with weight generally decreasing and bilirubin levels generally increasing. A nomogram published in 1999 depicting bilirubin levels by hour of age¹⁸ is widely used to evaluate trajectories of neonatal hyperbilirubinemia¹⁹ and has been incorporated into the American Academy of Pediatrics guideline on hyperbilirubinemia management.²⁰ However, similarly detailed hour-specific values and trajectories for newborn weight loss have not been previously reported. Therefore, we sought to develop a detailed graphical depiction of early trajectories of weight loss for exclusively breastfed newborns. Such a weight loss nomogram would be of great clinical utility because newborn weight is measured daily at varying hours of age, and these measurements become the basis for multiple clinical decisions including timing of discharge, need for lactation support or supplementation, and timing and type of newborn follow-up.^{21,22}

METHODS

Participants and Outcomes

The analysis included 161 471 newborns who were born at ≥ 36 weeks' gestation at 1 of 14 Kaiser Permanente Northern California hospitals between January 1, 2009, and December 31, 2013, who survived to discharge home and who did not receive Level II or Level III care. For this reason, newborns with infectious disease or congenital abnormalities requiring Level II or Level III care were not included in this cohort. For included newborns, data were extracted on all weights obtained during the birth hospitalization as well as on gestational age, method of delivery, length of stay, hospital of birth, maternal race/ethnicity, and timing (hour of age) and type (breast milk or formula) of all inpatient feedings. From this cohort, newborns were excluded if they had missing data on type of delivery, weight or feeding, birth weight < 2000 g or > 5000 g, multiple birth, discrepantly reported birth weights between data sources, or no weight

documented after 6 hours of age and before initiation of formula feeding.

Weight change was defined as the difference between birth weight and each weight recorded subsequently, calculated as a percentage of birth weight as is typically done daily in clinical practice. Excess weight loss was defined as the loss of $\geq 10\%$ of birth weight.^{9,23} Newborns with implausible weight loss or weight gain values ($> 10\%$ loss in the first 24 hours, $> 15\%$ loss at any time thereafter, gain $> 5\%$) were excluded. Because length of stay varies by method of delivery and because neonates are usually weighed once daily during the birth hospitalization but not usually within 6 hours of birth, weight loss percentiles were determined from 6 to 72 hours for vaginal births and from 6 to 96 hours for cesarean births, reflecting the difference in length of stay by delivery mode and the corresponding variation in availability of weight measurements between vaginally delivered newborns and those delivered by cesarean.²⁴




FIGURE 1
Derivation of the final analytic cohort.

TABLE 1 Demographic and Clinical Characteristics of Included Newborns, by Type of Delivery

	Vaginal (<i>n</i> = 83 433)	Cesarean (<i>n</i> = 25 474)
Birth wt (g)		
Mean (SD)	3416.9 (426.2)	3487.7 (462.6)
Median	3402	3470
Interquartile range	(3120–3700)	(3170–3790)
Range	(2000–5000)	(2010–5000)
Gestational age (wk), <i>n</i> (%)		
36	1429 (1.7)	458 (1.8)
37	5817 (7.0)	1707 (6.7)
38	15114 (18.1)	3830 (15.0)
39	27335 (32.8)	11515 (45.2)
40	24011 (28.8)	4743 (18.6)
41	9375 (11.2)	3041 (11.9)
42	352 (0.4)	175 (0.7)
43	0 (0.0)	5 (<0.1)
Gestational age (wk)		
Mean (SD)	39.2 (1.2)	39.1 (1.1)
Median	39	39
Interquartile range	(38–40)	(39–40)
Range	(36–42)	(36–43)
Maternal race/ethnicity, <i>n</i> (%)		
Hispanic	20234 (24.5)	5897 (23.1)
American Indian/Eskimo	335 (0.4)	104 (0.4)
Asian	19765 (23.7)	6079 (23.9)
Black, non-Hispanic	5165 (6.2)	1898 (7.5)
White, non-Hispanic	35704 (42.8)	10772 (42.3)
Unknown	2230 (2.7)	724 (2.8)
Newborn hospital length of stay (d) ^a		
Mean (SD)	1.7 (0.8)	2.8 (0.9)
Median	1.6	2.6
Interquartile range	(1.2–2.0)	(2.1–3.2)

^a There were missing length of stay data for 210 vaginally delivered and 87 Cesarean-delivered newborns.

A newborn was included in the final analysis if any weight measurement was obtained during the birth hospitalization after 6 hours of age and before exclusive breastfeeding was discontinued. Between 6 and 72 hours for vaginal births and 6 and 96 hours for cesarean births, all weights obtained during the birth hospitalization were used if at the time the weight was obtained the newborn remained exclusively breastfed as defined by the World Health Organization (nothing other than breast milk, vitamins, minerals, and oral medications).^{25,26} Newborns were censored at the time of the first supplementary feeding. No weights obtained subsequent to censoring or after discharge from the birth hospitalization were used in this analysis. This study was approved by the University of California San Francisco Committee on Human Research and by the Institutional

Review Boards of Penn State Medical College and Kaiser Permanente Northern California.

Analyses

Quantile regression methods appropriate for data with repeated measures were used to estimate 50th (median), 75th, 90th, and 95th percentiles of weight loss as a function of time after birth. The penalized fixed-effects model in the R package “Regression Quantiles for Panel Data” was used to estimate the percentile curves.²⁷ The model is an extension of ordinary quantile regression²⁸ to longitudinal settings

in which newborns may be weighed at multiple time points. The model accounts for multiple (or repeated) weights from a newborn by including a separate intercept parameter for each newborn, with regularization used to estimate these intercepts by shrinking them toward a common value. The amount of regularization is controlled by a tuning parameter, which was set equal to 5 so that most intercepts were shrunk to zero, a prudent choice for sparse longitudinal data that includes many newborns contributing only a single data point (62% of newborns in the sample had only 1 weight included in the analysis). A B-spline basis with 4 degrees of freedom was used to generate nonlinear percentile curves.²⁹ Nonparametric bootstrapping with 500 resamples and the percentile method were used to obtain confidence intervals for each percentile curve.³⁰ To examine the reliability and robustness of the final estimated percentile curves, we generated separate curves for the cohort born 2009–2010 and the cohort born 2011–2013.

We examined whether our nomograms were affected by the inclusion of late preterm, postterm, and small or large-for-gestational-age infants by removing all newborns born at <37 weeks and ≥42 weeks from the cohort, and those with birth weights below the 10th percentile and above the 90th percentile for gestational age. The quantile regression model was refit on the basis of these exclusions, and we examined how the estimated percentile curves changed.

To determine whether censoring newborns after formula use may have caused bias to the estimated percentile curves, we undertook

TABLE 2 Number of Weights Used in the Final Analysis, *n* (%)

Weights Recorded After Birth Weight	Vaginal (<i>n</i> = 83 433), <i>n</i> (%)	Cesarean (<i>n</i> = 25 474), <i>n</i> (%)
1	59 953 (71.9)	6402 (25.1)
2	21 440 (25.7)	12 423 (48.8)
3	1825 (2.2)	5555 (21.8)
4	215 (0.3)	1094 (4.3)

TABLE 3 Number of Weights Included for Each 12-Hour Interval

Time, h	Vaginal (n = 109 190), n (%)	Cesarean (n = 52 443), n (%)
6–12	23 914 (21.9)	7221 (13.8)
13–24	45 394 (41.6)	14 377 (27.4)
25–36	26 185 (24)	9866 (18.8)
37–48	10 833 (9.9)	11 192 (21.3)
49–60	2168 (2.0)	5040 (9.6)
61–72	696 (0.6)	3535 (6.7)
73–84	—	780 (1.5)
85–96	—	432 (0.8)

—, not included.

a sensitivity analysis. If previously exclusively breastfed newborns began using formula supplementation primarily because of high levels of weight loss, then the estimated curves may have underestimated the actual weight loss in the population. To examine this possibility, the sensitivity analysis recreated the nomograms after using matching to impute subsequent weights for newborns who had 1 included weight recorded during the birth hospitalization and then were censored. First, each newborn censored due to formula use was matched to 1 uncensored newborn using a greedy matching algorithm.³¹ Matches were conditional on 3 variables: time of most recent weight measurement in which the censored newborn had been exclusively breastfed (± 2 hours), percent weight loss at that time (± 3 percentage points), and gestational age (± 3 weeks). The weight loss data from the uncensored newborn that occurred after formula use began for the censored newborn was then imputed as the actual weight loss data for the censored newborn. Last, the quantile regression model was refit using the original data plus the imputed data for censored newborns to examine how the estimated percentile curves changed. The main assumption of this sensitivity analysis is that the weight loss history of the matched uncensored newborn reasonably approximates the weight loss history that would have been observed for the censored newborn had formula never been introduced.

RESULTS

From 161 471 newborns in the cohort, 108 907 were included in the final analysis, of whom 83 433 (76.6%) were delivered vaginally and 25 474 (23.4%) were delivered by cesarean. Among 52 564 excluded newborns, 33 715 (64%) were excluded because no weight was obtained in the eligible period and before formula feeding, 18 368 (35%) were excluded due to either multiple birth or missing or discrepant birth data, and 481 (0.9%) were excluded because weight loss exceeded 10% in 24 hours or 15% at any time or because weight gain exceeded 5% (Fig 1). Data were available on race/ethnicity for 97.3% of mothers, and the cohort was racially diverse; 24% of mothers were Hispanic, 24% were Asian, 7% were black non-Hispanic, and 43% were white non-Hispanic. Other demographic and clinical characteristics are summarized in Table 1.

For neonates delivered vaginally, 109 190 weights subsequent to birth weight were recorded (1.3 per newborn), and for those delivered by cesarean, 52 443 were recorded (2.1 per newborn). A majority delivered vaginally (71.9%) had only 1 weight recorded during the follow-up, whereas a majority delivered via cesarean (74.9%) had ≥ 2 weights recorded (Table 2). The difference reflects the longer length of stay for cesarean newborns. See Table 3 for details on the distribution of age in hours for included weights.

Differences in weight loss by delivery type appeared early and were clearly

evident within 24 hours of birth. Figure 2A presents percentile curves for vaginally delivered newborns. Median percent weight loss for these neonates was 4.2%, 7.1%, and 6.4% at 24, 48, and 72 hours of age, respectively. By 48 hours, almost 5% of vaginally delivered newborns had lost at least 10% of their birth weight, and by shortly after 48 hours, the percentile curves for median weight loss had begun to rise. Figure 2B presents percentile curves for newborns delivered by cesarean. Median percent weight loss among these neonates was 4.9%, 8.0%, 8.6% and 5.8% at 24, 48, 72, and 96 hours after delivery. By 48 hours of age, almost 10% of newborns delivered by cesarean had lost at least 10% of their birth weight, and by 72 hours of age, the percentile curves for median weight loss had begun to rise, but $>25\%$ were at least 10% below their birth weight.

Percentile curves with 95% confidence intervals overlaid are shown in Supplementary Figure 3 A and B. Percentile curves estimated from the cohort of 34 809 newborns born 2009–2010 were similar to the percentile curves estimated from the cohort of 74 098 newborns born 2011–2013 (Supplemental Figure 4 A and B). Percentile curves generated after removing late preterm, postterm, and small- and large-for-gestational-age newborns were not visually distinguishable from the main results (Supplemental Fig 5 A and B).

In our main analysis depicted in Fig 2 A and B, 8457 newborns delivered vaginally (10.1%) and 8414 newborns delivered by cesarean (33.0%) had at least 1 weight included in the analysis and at least 1 weight censored from the analysis because it was obtained subsequent to a supplementary feeding. In sensitivity analysis, weights were imputed for censored newborns based on matching to uncensored newborns. More than 98% of censored newborns had exact





FIGURE 2
 A, Estimated percentile curves of percent weight loss by time after birth for vaginal deliveries. B, Estimated percentile curves of percent weight loss by time after birth for Cesarean deliveries.

matches for time and gestational age, and >90% had a match of percent weight loss within 0.20 percentage points (Table 4). Percentile curves estimated from the original data plus the imputed data for censored weights were similar to the curves of the original data for newborns delivered vaginally and by cesarean. See the online supplement for this comparison (Supplemental Figure 6 A and B).

DISCUSSION

These results provide the first graphical depiction of hourly weight loss for exclusively breastfed newborns from a large, diverse population. Because weight changes steadily throughout the birth hospitalization and is measured at varied intervals from the hour of birth, these new nomograms should substantially aid medical management by allowing clinicians and lactation support providers to categorize newborn weight loss and calibrate decision-making to reflect hour of age. The use of these nomograms may allow a more nuanced clinical approach to weight loss in a manner similar to that of the widely used nomogram for hyperbilirubinemia. Similar to the bilitool.org Web site, a new Web site, <http://www.newbornweight.org>, has been developed by the study team to allow clinicians to compare individual newborns and their weight loss against our large sample.

With nomograms presented separately for newborns delivered vaginally and by cesarean, it is clear that differential weight loss by delivery method materialized early and persisted over time. Factors contributing to lactation difficulty have been well documented among those undergoing cesarean delivery.^{32,33} In addition, the administration of large volumes of intravenous fluid before cesarean may be an independent risk factor for newborn weight loss.^{34,35} Excess weight loss $\geq 10\%$, which has been

TABLE 4 Summary of Matched Variables for Censored Newborns

	Vaginal (<i>n</i> = 8457), <i>n</i> (%)	Cesarean (<i>n</i> = 8414), <i>n</i> (%)
Time (h)		
Exact match	8420 (99.6)	8325 (98.9)
1 h difference	36 (0.4)	83 (1.0)
2 h difference	1 (<0.1)	6 (0.1)
Weight loss, percentage points		
<0.05	7373 (87.2)	5998 (71.3)
0.05–0.20	651 (7.7)	1445 (17.2)
0.20–0.50	253 (3)	566 (6.7)
0.50–1	146 (1.7)	295 (3.5)
1–2	34 (0.4)	101 (1.2)
>2	0	9 (0.1)
Gestational age		
Exact match	8389 (99.2)	8241 (97.9)
1 wk difference	67 (0.8)	163 (1.9)
2 wk difference	1 (<0.1)	8 (0.1)
3 wk difference	0	2 (<0.1)

associated with increased risk of both hyperbilirubinemia and hypernatremic dehydration,^{10,14,16,32} was common among both groups. Almost 5% of vaginally delivered newborns and almost 10% of those delivered by cesarean had lost $\geq 10\%$ of their birth weight by 48 hours. By 72 hours of age, $>25\%$ of newborns delivered by cesarean had lost $\geq 10\%$ of their birth weight. Weight gain typically began at 48 to 72 hours of age. Newborns with greater weight loss tended to have weight nadir at an older age.

Results from this study are unique because they contain detailed data by hour of life in a large, diverse population for which data were available on type and timing of all feedings. Therefore, our analysis included all weights obtained during the duration of exclusive breastfeeding and excluded any weight obtained after an infant received a supplementary feeding. Reliability of these nomograms is supported by the similarity between curves generated by the cohort born 2009–2010 and those born 2011–2013. Moreover, the similarity between the results presented in the main analysis and the results of sensitivity analysis in which weights were imputed for censoring due to formula use offers further evidence of the robust nature of the findings.

These analyses have several limitations. First, weights for this study were obtained in the course of routine care with various scales calibrated according to the guidelines of individual institutions. However, the results therefore reflect weight loss in a setting of routine clinical practice. Second, feeding reports in our study were obtained from the electronic medical record maintained by the nursing staff, and nursing staff may not have been aware of or documented all feedings. If some mothers fed formula without documentation in the medical record, our results might underestimate weight loss for exclusively breastfed newborns. Third, our study did not have access to accurate reports of reasons given for initiating formula feeding. If caregivers initiated formula because of weight loss, our results might underestimate weight loss for exclusively breastfed newborns. However, the similarity between our main model and our model obtained by imputing weights for censored newborns suggests that this was not a substantial source of bias. Fourth, our curves extend only to 72 hours for vaginal births and 96 hours for births by cesarean and therefore do not represent weight nadir for all newborns, although the majority of measured infants did begin weight

gain during this period. Fifth, the Northern California population studied was racially diverse with 43% being white non-Hispanic. Results could be different for populations with other racial and ethnic compositions. Additionally, we had no data on parity or previous maternal breastfeeding experience; weight loss may have been ameliorated for newborns of mothers multiparous or with previous breastfeeding experience.²³ Sixth, our cohort excluded newborns who received Level II or Level III care but may have included some newborns with congenital anomalies such as trisomy 21. It is possible that such well newborns with congenital anomalies not requiring Level II or Level III care had different weight loss patterns than newborns without congenital anomalies, but the small proportion of well newborns with congenital anomalies makes this group unlikely to affect the nomogram results. These limitations illustrate the potential importance of a future large, prospective cohort study that could rigorously measure feeding practices and weights at prespecified intervals while simultaneously assessing the effect of various weight trajectories on clinical outcomes.

CONCLUSIONS

The availability of detailed data on weight and feeding for a large cohort allows this study to present the first graphical depiction of hourly weight loss for exclusively breastfed newborns from a large, diverse population. Similar to the widely used bilirubin nomogram by Bhutani et al, our findings, available at <http://www.newbornweight.org>, have the potential for wide clinical applicability and may have strong generalizability because data are included from a large, racially diverse cohort delivered at multiple hospitals.¹⁸ Our curves demonstrate that expected weight loss differs

substantially by method of delivery and that this difference persists over time. Our results also show that

weight loss $\geq 10\%$ of birth weight is common and often occurs earlier in the postnatal course than previously

documented. These nomograms provide normative data that may inform clinical care.

Address correspondence to Valerie Flaherman, MD, MPH, University of California, San Francisco, 3333 California St, Box 0503, San Francisco, CA 94143-0503. E-mail: flahermanv@peds.ucsf.edu

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2015 by the American Academy of Pediatrics

FINANCIAL DISCLOSURE: The authors have indicated they have no financial relationships relevant to this article to disclose.

FUNDING: Supported by grant R40 MC 26811 through the US Department of Health and Human Services, Health Resources and Services Administration, Maternal and Child Health Research Program and grant K23 HD059818 from the National Institute of Child Health and Human Development. Funded by the National Institutes of Health.

POTENTIAL CONFLICT OF INTEREST: The authors have indicated they have no potential conflicts of interest to disclose.

COMPANION PAPER: A companion to this article can be found on page e174, online at www.pediatrics.org/cgi/doi/10.1542/peds.2014-3354.

REFERENCES

1. UNICEF/WHO. *Baby-Friendly Hospital Initiative: Revised, Updated and Expanded for Integrated Care, Section 1, Background and Implementation, Preliminary version*. 2006. Available at: http://www.who.int/nutrition/topics/BFHI_Revised_Section1.pdf. Accessed October 14, 2014
2. US Department of Health and Human Services. *Developing Healthy People 2020—Maternal, Infant and Child Health*. 2010. Available at: <http://www.healthypeople.gov/hp2020/Objectives/ViewObjective.aspx?id=177&TopicArea=Maternal%2c+Infant+and+Child+Health&Objective=MICH+HP2020%e2%80%9312&TopicAreaId=32>>. Accessed April 6, 2010
3. American Academy of Pediatrics Section on Breastfeeding. Breastfeeding and the use of human milk. *Pediatrics*. 2012; 129(3). Available at: www.pediatrics.org/cgi/content/full/129/3/e827
4. Centers for Disease Control and Prevention. *Maternal, infant and child health. Data 2020*. 2013. Available at: <http://www.healthypeople.gov/2020/topics-objectives/topic/maternal-infant-and-child-health>. Accessed March 20, 2014
5. Slusher TM, Slusher IL, Keating EM, et al. Comparison of maternal milk (breastmilk) expression methods in an African nursery. *Breastfeed Med*. 2012;7(2):107–111
6. Flaherman VJ, Gay B, Scott C, Avins A, Lee KA, Newman TB. Randomised trial comparing hand expression with breast pumping for mothers of term newborns feeding poorly. *Arch Dis Child Fetal Neonatal Ed*. 2012;97(1):F18–F23
7. Saint L, Smith M, Hartmann PE. The yield and nutrient content of colostrum and milk of women from giving birth to 1 month post-partum. *Br J Nutr*. 1984;52(1):87–95
8. Macdonald PD, Ross SR, Grant L, Young D. Neonatal weight loss in breast and formula fed infants. *Arch Dis Child Fetal Neonatal Ed*. 2003;88(6):F472–F476
9. Chantry CJ, Nommsen-Rivers LA, Pearson JM, Cohen RJ, Dewey KG. Excess weight loss in first-born breastfed newborns relates to maternal intrapartum fluid balance. *Pediatrics*. 2011;127(1). Available at: www.pediatrics.org/cgi/content/full/127/1/e171
10. van Dommelen P, van Wouwe JP, Breuning-Boers JM, van Buuren S, Verkerk PH. Reference chart for relative weight change to detect hypernatraemic dehydration. *Arch Dis Child*. 2007;92(6):490–494
11. Paul IM, Lehman EB, Hollenbeck CS, Maisels MJ. Preventable newborn readmissions since passage of the Newborns' and Mothers' Health Protection Act. *Pediatrics*. 2006;118(6):2349–2358
12. Escobar GJ, Greene JD, Hulac P, et al. Rehospitalisation after birth hospitalisation: patterns among infants of all gestations. *Arch Dis Child*. 2005; 90(2):125–131
13. O'Donnell HC, Colman G, Trachtman RA, Velazco N, Racine AD. Impact of newborn follow-up visit timing on subsequent ED visits and hospital readmissions: an instrumental variable analysis. *Acad Pediatr*. 2014;14(1):84–91
14. Salas AA, Salazar J, Burgoa CV, De-Villegas CA, Quevedo V, Soliz A. Significant weight loss in breastfed term infants readmitted for hyperbilirubinemia. *BMC Pediatr*. 2009;9:82
15. Chen CF, Hsu MC, Shen CH, et al. Influence of breast-feeding on weight loss, jaundice, and waste elimination in neonates. *Pediatr Neonatol*. 2011;52(2):85–92
16. Chang RJ, Chou HC, Chang YH, et al. Weight loss percentage prediction of subsequent neonatal hyperbilirubinemia in exclusively breastfed neonates. *Pediatr Neonatol*. 2010;53(1):41–44
17. Chen HL, Wang YH, Tseng HI, Lu CC. Neonatal readmission within 2 weeks after birth. *Acta Paediatr Taiwan*. 2005; 46(5):289–293
18. Bhutani VK, Johnson L, Sivieri EM. Predictive ability of a predischarge hour-specific serum bilirubin for subsequent significant hyperbilirubinemia in healthy term and near-term newborns. *Pediatrics*. 1999;103(1):6–14
19. Longhurst C, Turner S, Burgos AE. Development of a Web-based decision support tool to increase use of neonatal hyperbilirubinemia guidelines. *Jt Comm J Qual Patient Saf*. 2009;35(5):256–262

20. American Academy of Pediatrics Subcommittee on Hyperbilirubinemia. Management of hyperbilirubinemia in the newborn infant 35 or more weeks of gestation. *Pediatrics*. 2004;114(1):297–316
21. Flaherman VJ, Bokser S, Newman TB. First-day newborn weight loss predicts in-hospital weight nadir for breastfeeding infants. *Breastfeed Med*. 2010;5(4):165–168
22. Flaherman VJ, Kuzniewicz MW, Li S, Walsh E, McCulloch CE, Newman TB. First-day weight loss predicts eventual weight nadir for breastfeeding newborns. *Arch Dis Child Fetal Neonatal Ed*. 2013;98(6):F488–F492
23. Dewey KG, Nommsen-Rivers LA, Heinig MJ, Cohen RJ. Risk factors for suboptimal infant breastfeeding behavior, delayed onset of lactation, and excess neonatal weight loss. *Pediatrics*. 2003;112(3 Pt 1):607–619
24. Datar A, Sood N. Impact of postpartum hospital-stay legislation on newborn length of stay, readmission, and mortality in California. *Pediatrics*. 2006;118(1):63–72
25. Labbok MH, Starling A. Definitions of breastfeeding: call for the development and use of consistent definitions in research and peer-reviewed literature. *Breastfeed Med*. 2012;7(6):397–402
26. World Health Organization. Nutrition: exclusive breastfeeding. 2011. Available at: http://www.who.int/nutrition/topics/exclusive_breastfeeding/en. Accessed March 14, 2011
27. Koenker R. Quantile regression for longitudinal data. *J Multivariate Anal*. 2004;91:74–89
28. Koenker R, Hoallock KF. Quantile regression. *J Econ Perspect*. 2001;15:143–156
29. Koenker R, Ng P, Portnoy S. Quantile smoothing splines. *Biometrika*. 1994;81(4):673–680
30. Efron B, Tibshirani RJ. An introduction to the bootstrap. New York, NY: Springer Science; 1994
31. Bergsralh DJ, Kosanke JL. *Computerized Matching of Controls* (Technical Report Series No. 56). Rochester, MN: Mayo Clinic; 1995
32. Caglar MK, Ozer I, Altugan FS. Risk factors for excess weight loss and hypernatremia in exclusively breast-fed infants. *Braz J Med Biol Res*. 2006;39(4):539–544
33. Chapman DJ, Pérez-Escamilla R. Identification of risk factors for delayed onset of lactation. *J Am Diet Assoc*. 1999;99(4):450–454, quiz 455–456
34. Noel-Weiss J, Woodend AK, Peterson WE, Gibb W, Groll DL. An observational study of associations among maternal fluids during parturition, neonatal output, and breastfed newborn weight loss. *Int Breastfeed J*. 2011;6:9
35. Watson J, Hodnett E, Armson BA, Davies B, Watt-Watson J. A randomized controlled trial of the effect of intrapartum intravenous fluid management on breastfed newborn weight loss. *J Obstet Gynecol Neonatal Nurs*. 2012;41(1):24–32

Early Weight Loss Nomograms for Exclusively Breastfed Newborns
Valerie J. Flaherman, Eric W. Schaefer, Michael W. Kuzniewicz, Sherian X. Li,
Eileen M. Walsh and Ian M. Paul
Pediatrics 2015;135:e16
DOI: 10.1542/peds.2014-1532 originally published online December 1, 2014;

Updated Information & Services	including high resolution figures, can be found at: http://pediatrics.aappublications.org/content/135/1/e16
References	This article cites 22 articles, 11 of which you can access for free at: http://pediatrics.aappublications.org/content/135/1/e16#BIBL
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Nutrition http://www.aappublications.org/cgi/collection/nutrition_sub Breastfeeding http://www.aappublications.org/cgi/collection/breastfeeding_sub
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.aappublications.org/site/misc/Permissions.xhtml
Reprints	Information about ordering reprints can be found online: http://www.aappublications.org/site/misc/reprints.xhtml

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Early Weight Loss Nomograms for Exclusively Breastfed Newborns

Valerie J. Flaherman, Eric W. Schaefer, Michael W. Kuzniewicz, Sherian X. Li,
Eileen M. Walsh and Ian M. Paul

Pediatrics 2015;135:e16

DOI: 10.1542/peds.2014-1532 originally published online December 1, 2014;

The online version of this article, along with updated information and services, is
located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/135/1/e16>

Data Supplement at:

<http://pediatrics.aappublications.org/content/suppl/2014/11/25/peds.2014-1532.DCSupplemental>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2015 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

