

The Recommendation for Rest Following Acute Concussion

William P. Meehan III, MD^{a,b,c,d}, Richard G. Bachur, MD^c

FREE

^aThe Micheli Center for Sports Injury Prevention, Waltham, Massachusetts; and ^bSports Concussion Clinic, Division of Sports Medicine, ^cDivision of Emergency Medicine, and ^dBrain Injury Center, Boston Children's Hospital, Boston, Massachusetts

Opinions expressed in these commentaries are those of the author and not necessarily those of the American Academy of Pediatrics or its Committees.

www.pediatrics.org/cgi/doi/10.1542/peds.2014-3665

DOI: 10.1542/peds.2014-3665

Accepted for publication Nov 21, 2014

Address correspondence to William P. Meehan III, MD, The Micheli Center for Sports Injury Prevention, 9 Hope Ave, Ste 100, Waltham, MA 02453. E-mail: william.meehan@childrens.harvard.edu

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2015 by the American Academy of Pediatrics

FINANCIAL DISCLOSURE: Dr Meehan receives royalties from ABC-Clío publishing for the sale of his book, *Kids, Sports, and Concussion: A Guide for Coaches and Parents*, and royalties from Wolters Kluwer for working as an author for *UpToDate*. His research is funded, in part, by a grant from the National Football Players Association. Dr Bachur receives royalties from Wolters Kluwer for his work as editor for *UpToDate* and from Lippincott Williams & Wilkins publishing for his work as editor of *Textbook of Pediatric Emergency Medicine*. His research on biomarkers for appendicitis receives industry sponsorship from Astute Medical.

FUNDING: No external funding

POTENTIAL CONFLICT OF INTEREST: Dr Meehan has received grant funding from the National Football League Players Association and philanthropic support from the National Hockey League Alumni Association. Dr Bachur has no potential conflicts to disclose.

COMPANION PAPER: A companion to this article can be found on page 213, and online at www.pediatrics.org/cgi/doi/10.1542/peds.2014-0966.

The acute management of concussion is an important area of investigation and an area of clinical importance to the pediatrician. Recent studies suggest that concussions account for more than 10% of all sport-related injuries sustained by high school athletes and reveal an increase in the number of children with concussions being cared for in emergency departments.¹⁻³ At present, the mainstay of concussion management calls for both cognitive and physical rest until the acute symptoms resolve, followed by a graded return to activity.⁴ As noted by several medical societies, the evidence on which the recommendations for rest are based is sparse.⁴⁻⁶ This relative lack of evidence is due, in part, to the difficult nature of quantifying and tracking levels of physical and, particularly, cognitive activity. In this issue of *Pediatrics*, Thomas et al⁷ take this challenge head on; they should be applauded for their efforts.

The authors describe the results of a randomized trial of 88 patients cared for in a pediatric emergency department who were prescribed either strict rest or standard care as treatment of a concussion. Those participants in the intervention group received recommendations for 5 days of strict rest at home, which specified no work, no school, and no physical activity. Those participants in the control group were prescribed usual care, treating physicians prescribed whatever activity restrictions they saw fit. Study participants who were prescribed strict rest had a slower resolution of their symptoms and had a higher symptom burden during the first 10 days after their emergency

department visit than those prescribed usual care. There were no significant differences between the intervention group (prescribed strict rest) and the control group (prescribed usual care) with regard to balance scores, computerized neurocognitive assessments, or most paper-based neuropsychological tests after injury. Interestingly, the intervention group performed better than the control group on the symbol digit modalities test 3 days after injury, but worse than the control group on day 10 after injury. The authors postulate that 5 days of strict rest requiring restrictions in activities may cause deleterious effects, namely an increase in emotional symptoms.⁷ This sentiment is consistent with our clinical impression, as well as emerging evidence suggesting that emotional symptoms increase over the course of recovery from concussion.^{8,9}

Although this study adds some data on which to base recommendations for rest after a concussion, the optimal duration of rest after concussion remains unknown. Indeed, the optimal period of rest may vary, depending on age, gender, point in the calendar year, initial symptom level, the particular symptoms that predominate, the level of cognitive function, or other variables. As clinicians, we are forced to use the existing evidence, however limited, to develop a plan for our patients. In light of current consensus recommendations, previous investigations, and the study by Thomas et al,⁷ a recommendation of reasonable rest for the first few days after a concussion followed by a gradual resumption of cognitive

activities seems prudent. The resumption of cognitive activities by patients who remain symptomatic may require instituting academic accommodations.^{5,10,11} In addition, symptomatic patients may benefit, after a period of rest, from a gradual resumption of aerobic exercise as tolerated by symptoms, provided no increased risk of trauma to the head.¹² Once again, the timing of the initiation of exercise after injury remains unknown. Given the variability of forces involved in different concussive injuries, the different symptom clusters and burdens experienced by patients who sustain concussions, and the observed variability of recovery patterns after concussion, the entire plan for managing a concussion should not be determined in the emergency department. Rather, a few days of rest followed by prompt follow-up with the pediatrician, sports medicine physician, or other capable provider should be recommended, and each management plan should be tailored to each individual patient.

REFERENCES

1. Marar M, McIlvain NM, Fields SK, Comstock RD. Epidemiology of concussions among United States high school athletes in 20 sports. *Am J Sports Med.* 2012;40(4):747–755
2. Colvin JD, Thurm C, Pate BM, Newland JG, Hall M, Meehan WP III. Diagnosis and acute management of patients with concussion at children's hospitals. *Arch Dis Child.* 2013;98(12):934–938
3. Bakhos LL, Lockhart GR, Myers R, Linakis JG. Emergency department visits for concussion in young child athletes. *Pediatrics.* 2010;126(3). Available at: www.pediatrics.org/cgi/content/full/e550
4. McCrory P, Meeuwisse W, Aubry M, et al. Consensus statement on concussion in sport—the 4th International Conference on Concussion in Sport held in Zurich, November 2012. *Clin J Sport Med.* 2013; 23(2):89–117
5. Halstead ME, Walter KD; Council on Sports Medicine and Fitness. American Academy of Pediatrics. Clinical report— sport-related concussion in children and adolescents. *Pediatrics.* 2010;126(3): 597–615
6. Giza CC, Kutcher JS, Ashwal S, et al. Summary of evidence-based guideline update: evaluation and management of concussion in sports: report of the Guideline Development Subcommittee of the American Academy of Neurology. *Neurology.* 2013;80(24):2250–2257
7. Thomas DG, Apps JA, Hoffmann RG, McCrea M, Hammeke T. Benefits of strict rest after acute concussion: a randomized controlled trial. *Pediatrics.* 2015;135(2):213–223
8. Eisenberg MA, Meehan WP III, Mannix R. Duration and course of post-concussive symptoms. *Pediatrics.* 2014;133(6):999–1006
9. Joyce AS, LaBella CR, Carl RL, Lai JS, Zelko FA. The Postconcussion Symptom Scale: utility of a three-factor structure [published online ahead of print September 29, 2014]. *Med Sci Sports Exerc.* 2014. doi: 10.1249/MSS.0000000000000534
10. Master CL, Gioia GA, Leddy JJ, Grady MF. Importance of 'return-to-learn' in pediatric and adolescent concussion. *Pediatr Ann.* 2012;41(9):1–6
11. McGrath N. Supporting the student-athlete's return to the classroom after a sport-related concussion. *J Athl Train.* 2010;45(5):492–498
12. Leddy JJ, Kozlowski K, Donnelly JP, Pendergast DR, Epstein LH, Willer B. A preliminary study of subsymptom threshold exercise training for refractory post-concussion syndrome. *Clin J Sport Med.* 2010;20(1):21–27

The Recommendation for Rest Following Acute Concussion

William P. Meehan III and Richard G. Bachur

Pediatrics 2015;135;362

DOI: 10.1542/peds.2014-3665 originally published online January 5, 2015;

Updated Information & Services

including high resolution figures, can be found at:
<http://pediatrics.aappublications.org/content/135/2/362>

References

This article cites 10 articles, 5 of which you can access for free at:
<http://pediatrics.aappublications.org/content/135/2/362#BIBL>

Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):
Sports Medicine/Physical Fitness
http://www.aappublications.org/cgi/collection/sports_medicine:physical_fitness_sub
Concussion
http://www.aappublications.org/cgi/collection/concussion_sub

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
<http://www.aappublications.org/site/misc/Permissions.xhtml>

Reprints

Information about ordering reprints can be found online:
<http://www.aappublications.org/site/misc/reprints.xhtml>

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

The Recommendation for Rest Following Acute Concussion

William P. Meehan III and Richard G. Bachur

Pediatrics 2015;135;362

DOI: 10.1542/peds.2014-3665 originally published online January 5, 2015;

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/135/2/362>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2015 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

