

Early Exposure to Dietary Sugar and Salt

Susan S. Baker, MD, PhD, Robert D. Baker, MD, PhD

Obesity, with its comorbidities such as cardiovascular heart disease (CVD), is one of the most common chronic diseases in childhood.¹ It probably begins early in life, and as the prevalence continues to rise the diets of young children assume increasing significance. Cogswell et al² provide useful information on the salt and sugar contents of commercial food products that can be used to improve the diets of young children.

Taste preferences for sugar and salt are mostly innate. Sugar and salt contribute to the overall pleasure and enjoyment of food.^{3,4} Sweet taste receptors respond to all classes of sweet tastants including natural sugars, artificial sweeteners, d-amino acids, and intensely sweet proteins. Pleasure from sweet taste promotes adequate caloric intake for times when food is scarce. Salty taste supports the proper dietary electrolyte balance. This happy relationship becomes a problem when there is a surfeit of food. Competitive food markets take advantage of these preferences to encourage consumption. Children are an important target because food preferences, eating behaviors, and decisions about pleasurable foods begin early and probably last throughout the life.⁵

Sugar is a caloric source, and as caloric intake increases, so does weight gain unless compensatory activity ensues. Most fruits, vegetables, and grains have naturally occurring sugars that are simply part of the food and carry with them nutrients such as vitamin C, folate, and B vitamins. The problem comes when sugars are added to

increase consumption. Because sweetness can induce reward and craving similar to that seen in drug addiction, some have argued that the concept of addiction could apply to the intake of sweet foods.⁶

Salt is naturally occurring in most foods. Little information is available on dietary salt or recommendations for salt intake for children ≤ 2 years of age. However, the Institute of Medicine⁷ and the World Health Organization⁸ recommend that, in general, dietary sodium be limited to avoid CVD. Because preferences for sodium intake later in life are related to early experiences,⁹ infants and young children should not be exposed to high-sodium foods.

Cogswell et al² sought to quantify the amount of sugar and sodium added to infant and young child foods. They used a commercial nutrition database, food labels, and the Food and Drug Administration reference amounts customarily consumed per eating occasion (RACC) to estimate the sugar and sodium content of foods commercially prepared for and consumed by this age group. The good news is that little sugar and sodium is added to food intended for infants. The bad news is that many foods prepared for toddlers contain added sugar and salt. For example, the mean sodium content in toddler meals and dinners is 361 mg/RACC, but sodium content varied by 836 mg/RACC. Savory snacks and sides also have a range of sodium, and some have the same amount of sodium as salted potato chips. The adequate intake for toddlers is only 1000 mg/day,¹⁰

FREE

Digestive Diseases and Nutrition Center, University at Buffalo, Buffalo, New York

www.pediatrics.org/cgi/doi/10.1542/peds.2014-4028

DOI: 10.1542/peds.2014-4028

Accepted for publication Dec 19, 2014

Address correspondence to Susan S. Baker, MD, PhD, Digestive Diseases and Nutrition Center, University at Buffalo, 219 Bryant St, Buffalo, NY 14222. E-mail: sbaker@upa.chob.edu

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2015 by the American Academy of Pediatrics

FINANCIAL DISCLOSURE: The authors have indicated they have no financial relationships relevant to this article to disclose.

FUNDING: No external funding.

POTENTIAL CONFLICT OF INTEREST: The authors have indicated they have no potential conflicts of interest to disclose.

COMPANION PAPER: A companion to this paper can be found on page 416, and online at www.pediatrics.org/cgi/doi/10.1542/peds.2014-3251.

so young children could consistently consume too much sodium. Similarly, many toddler foods, especially convenience snacks, have added sugar, and often the form is a concentrated juice, which may be interpreted as a healthful part of the food. The lack of specificity in labeling made it difficult for the investigators to be precise in the amount of added sugar because natural and added sugars are reported as 1 number. Nevertheless, some commercially prepared toddler desserts contain more sugar than vanilla ice cream. This study shows that toddlers are exposed to foods that have an unnecessarily high content of salt and sugar. This could lead them to develop a desire for these tastes for the rest of their lives. The continued consumption of foods with high sodium and sugar content places children at risk for obesity and CVD, risks that they should not experience and risks that we, as pediatricians, have an obligation to mitigate.

This report is important for several reasons. It points out the variability of added sugars and salt in toddler

foods and the possibility that with guidance parents might make better choices. It provides information that pediatricians can use to guide parents and policymakers can use to set standards for commercial foods intended for children. Alternatively, of course, pediatricians could encourage families to provide home-prepared foods that have no additions and set the stage for lifelong healthy eating.

REFERENCES

1. Ogden CL, Carroll MD, Kit BK, Flegal KM. Prevalence of obesity in the United States, 2009–2010. NCHS data brief no. 82. Hyattsville, MD: National Center for Health Statistics; 2012:1–8
2. Cogswell ME, Gunn JP, Yuan K, Park S, Merritt R. Sodium and sugar in complementary infant and toddler foods sold in the United States. *Pediatrics*. 2015;135(3):416–423
3. Chandrashekar J, Hoon MA, Ryba NJ, Zuker CS. The receptors and cells for mammalian taste. *Nature*. 2006; 444(7117):288–294
4. Yarmolinsky DA, Zuker CS, Ryba NJ. Common sense about taste: from mammals to insects. *Cell*. 2009;139(2): 234–244
5. Birch LL. Development of food preferences. *Annu Rev Nutr*. 1999;19: 41–62
6. Ahmed SH, Guillem K, Vandaele Y. Sugar addiction: pushing the drug–sugar analogy to the limit. *Curr Opin Clin Nutr Metab Care*. 2013;16(4): 434–439
7. In: Strom BL, Yaktine AL, Oria M, eds. *Sodium Intake in Populations: Assessment of Evidence*. Washington, DC: National Academies Press; 2013
8. World Health Organization. *Guideline: Sodium Intake for Adults and Children*. Geneva, Switzerland: World Health Organization; 2012
9. Stein LJ, Cowart BJ, Beauchamp GK. The development of salty taste acceptance is related to dietary experience in human infants: a prospective study. *Am J Clin Nutr*. 2012;95(1):123–129
10. IOM (Institute of Medicine). *Dietary Reference Intakes for Water, Potassium, Sodium, Chloride, and Sulfate*. Washington, DC: National Academies Press; 2005

Early Exposure to Dietary Sugar and Salt

Susan S. Baker and Robert D. Baker

Pediatrics 2015;135;550

DOI: 10.1542/peds.2014-4028 originally published online February 2, 2015;

Updated Information & Services	including high resolution figures, can be found at: http://pediatrics.aappublications.org/content/135/3/550
References	This article cites 6 articles, 2 of which you can access for free at: http://pediatrics.aappublications.org/content/135/3/550#BIBL
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Nutrition http://www.aappublications.org/cgi/collection/nutrition_sub Obesity http://www.aappublications.org/cgi/collection/obesity_new_sub
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.aappublications.org/site/misc/Permissions.xhtml
Reprints	Information about ordering reprints can be found online: http://www.aappublications.org/site/misc/reprints.xhtml

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Early Exposure to Dietary Sugar and Salt

Susan S. Baker and Robert D. Baker

Pediatrics 2015;135;550

DOI: 10.1542/peds.2014-4028 originally published online February 2, 2015;

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/135/3/550>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2015 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

