

Primary Care Interventions for Pediatric Obesity: Need for an Integrated Approach

Cara B. Ebbeling, PhD^a, Richard C. Antonelli, MD, MS^b

Recommendations for preventing and treating pediatric obesity suggest a staged approach with escalating intensity of care.¹ Stage 1 is Prevention Plus, requiring sufficient capacity in a primary care setting to provide basic messages pertaining to healthful dietary and physical activity behaviors. Stages 2 and 3 are Structured Weight Management and Comprehensive Multidisciplinary Intervention, respectively, requiring more frequent clinical encounters and involving a registered dietitian (RD) and a behavioral medicine provider. Stage 2 may be appropriate in a primary care setting, although inadequate training and financial support for structured intervention often are limiting factors. Stage 3 requires primary care providers (PCPs) to coordinate patient care with a pediatric weight management clinic and, in some cases, community partners that provide opportunities for physical activity. Stage 4 is Tertiary Care Intervention and may include a very low-calorie diet, medication, and surgery.

Outcomes of clinical interventions for pediatric obesity are variable. In primary care settings, some interventions promote changes in diet, physical activity, or television viewing but do not achieve reductions in BMI^{2,3}; others have favorable, albeit sometimes small, effects on BMI.⁴⁻⁶ Although effective for certain patients/families, weight management clinics often are not feasible because of barriers associated with accessibility, transportation, and cost.⁷⁻⁹ Clearly, novel approaches are necessary to

enhance interventions in primary care settings, extending the impact of weight management clinics and thereby avoiding progression to tertiary care referrals.

The trial reported by Resnicow et al¹⁰ in this issue of *Pediatrics* provides impressive data on the effectiveness of motivational interviewing (MI) to reduce BMI in primary care. The trial involved 42 practices in the American Academy of Pediatrics' Pediatric Research in Office Settings network, with enrollment of overweight and obese children aged 2 to 8 years. Interventions included usual care, MI from a PCP (4 sessions), and MI from both a PCP (4 sessions) and an RD (6 sessions). Usual care comprised standard educational materials presented during routine visits. The "PCP only" and "PCP+RD" interventions augmented usual care with counseling directed toward building motivation for behavior change and collaborating with families to set action-oriented goals around discrete behaviors. The PCPs and RDs who implemented these interventions received MI training from a psychologist and booster training via DVD.¹¹ Reductions in BMI percentile over 2 years were 1.8, 3.8, and 4.9 units for the usual care, "PCP only," and "PCP+RD" interventions, respectively, suggesting that intervention intensity may be an important consideration. However, although PCPs completed >75% of their expected sessions, RDs completed <50%. Resnicow et al^{10,11} allude to a gap in care coordination between PCPs and RDs, leading to lack of care integration, as a partial

FREE

^aNew Balance Foundation Obesity Prevention Center, and
^bDivision of General Pediatrics, Department of Medicine,
Boston Children's Hospital, Boston, Massachusetts

Opinions expressed in these commentaries are those of the authors and not necessarily those of the American Academy of Pediatrics or its Committees.

www.pediatrics.org/cgi/doi/10.1542/peds.2015-0495

DOI: 10.1542/peds.2015-0495

Accepted for publication Feb 10, 2015

Address correspondence to Cara B. Ebbeling, PhD, New Balance Foundation Obesity Prevention Center, Boston Children's Hospital, 300 Longwood Ave (AU307), Boston, MA 02115. E-mail: cara.ebbeling@childrens.harvard.edu

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2015 by the American Academy of Pediatrics

FINANCIAL DISCLOSURE: The authors have indicated they have no financial relationships relevant to this article to disclose.

FUNDING: Supported by the New Balance Foundation. The content is solely the responsibility of the authors and does not necessarily represent the official views of the New Balance Foundation.

POTENTIAL CONFLICT OF INTEREST: The authors have indicated they have no potential conflicts of interest to disclose.

COMPANION PAPER: A companion to this article can be found on page 649, and online at www.pediatrics.org/cgi/doi/10.1542/peds.2014-1880.

explanation for the low session completion rate among RDs.

Integrated care has been defined as being coordinated across professionals, facilities, and support systems; continuous over time and between visits; tailored to needs and preferences of patients/families; and based on shared responsibility between patients/families and caregivers for optimizing health.¹² This has been further elaborated for pediatrics to emphasize the crucial role of service integration across multiple sectors impacting child health outcomes.¹³ So-called wrap-around services involving community partners (such as recreational/fitness facilities, schools, day care and after-school programs, grocery stores, or restaurants) likely would enhance an integrated, team-based approach for the most vulnerable children.

Several components of the trial reported by Resnicow et al¹⁰ provide impetus for asking important questions when developing integrated care models for treating pediatric obesity.

- Patients were from a large network of primary care practices. What linkages can be established between primary care practices and weight management clinics to coordinate care with obesity specialists (psychologist, dietitian, physicians) when necessary? Linkages must extend beyond referrals and notes in electronic medical records to systems for training,¹⁴ ongoing consultation, and possible comanagement.
- A psychologist trained the PCPs and RDs in MI. As such, the psychologist was part of the multidisciplinary team. However, timely and effective communication between psychologists and PCPs or RDs, to manage specific cases, was not part of the intervention. How can PCPs facilitate access to specialty colleagues for collaboration? Emerging communication technology could support such enhanced access.

- Parents completed a questionnaire pertaining to diet, physical activity, and sedentary time, and PCPs shared responses with RDs. However, a form for recording patient encounters and thereby promoting communication between PCPs and RDs was apparently inadequate. How can we ensure adequate and efficient “handoffs” among members of a team in different settings?
- Assisting patients/families to develop action plans involving utilization of community programs is within the purview of MI.¹¹ How can interaction with community partners be extended, beyond PCPs advising patients to use programs, to obtaining updates on progress of individual patients?

Clearly, answers to these questions require collaboration among numerous stakeholders. Substantial effort must be directed toward coordinating care across settings to integrate services centered on the comprehensive needs of patients/families¹³ and measuring performance of care-coordination activities.¹⁵

REFERENCES

1. Barlow SE; Expert Committee. Expert committee recommendations regarding the prevention, assessment, and treatment of child and adolescent overweight and obesity: summary report. *Pediatrics*. 2007;120(suppl 4):S164–S192
2. McCallum Z, Wake M, Gerner B, et al. Outcome data from the LEAP (Live, Eat and Play) trial: a randomized controlled trial of a primary care intervention for childhood overweight/mild obesity. *Int J Obes (Lond)*. 2007;31(4):630–636
3. Taveras EM, Gortmaker SL, Hohman KH, et al. Randomized controlled trial to improve primary care to prevent and manage childhood obesity: the High Five for Kids study. *Arch Pediatr Adolesc Med*. 2011;165(8):714–722
4. Wald ER, Moyer SC, Eickhoff J, Ewing LJ. Treating childhood obesity in primary care. *Clin Pediatr (Phila)*. 2011;50(11):1010–1017
5. DeBar LL, Stevens VJ, Perrin N, et al. A primary care-based, multicomponent

- lifestyle intervention for overweight adolescent females. *Pediatrics*. 2012; 129(3). Available at: www.pediatrics.org/cgi/content/full/129/3/e611
6. Dolinsky DH, Armstrong SC, Walter EB, Kemper AR. The effectiveness of a primary care-based pediatric obesity program. *Clin Pediatr (Phila)*. 2012;51(4):345–353
 7. Ambler KA, Hagedorn DW, Ball GD. Referrals for pediatric weight management: the importance of proximity. *BMC Health Serv Res*. 2010;10:302
 8. Hampf S, Paves H, Laubscher K, Eneli I. Patient engagement and attrition in pediatric obesity clinics and programs: results and recommendations. *Pediatrics*. 2011;128(suppl 2):S59–S64
 9. Skelton JA, Beech BM. Attrition in paediatric weight management: a review of the literature and new directions. *Obes Rev*. 2011;12(5):e273–e281
 10. Resnicow K, McMaster F, Bocian A, et al. Motivational interviewing and dietary counseling for obesity in primary care: an RCT. *Pediatrics*. 2015;135(4):649–657
 11. Resnicow K, McMaster F, Woolford S, et al. Study design and baseline description of the BMI2 trial: reducing paediatric obesity in primary care practices. *Pediatr Obes*. 2012;7(1):3–15
 12. Singer SJ, Burgers J, Friedberg M, Rosenthal MB, Leape L, Schneider E. Defining and measuring integrated patient care: promoting the next frontier in health care delivery. *Med Care Res Rev*. 2011;68(1):112–127
 13. Council on Children with Disabilities and Medical Home Implementation Project Advisory Committee. Patient- and family-centered care coordination: a framework for integrating care for children and youth across multiple systems. *Pediatrics*. 2014;133(5). Available at: www.pediatrics.org/cgi/content/full/133/5/e1451
 14. Rhodes ET, Ebbeling CB, Meyers AF, et al. Pediatric obesity management: variation by specialty and awareness of guidelines. *Clin Pediatr (Phila)*. 2007;46(6):491–504
 15. Antonelli RC, Stille CJ, Antonelli DM. Care coordination for children and youth with special health care needs: a descriptive, multisite study of activities, personnel costs, and outcomes. *Pediatrics*. 2008; 122(1). Available at: www.pediatrics.org/cgi/content/full/122/1/e209

Primary Care Interventions for Pediatric Obesity: Need for an Integrated Approach

Cara B. Ebbeling and Richard C. Antonelli

Pediatrics 2015;135;757

DOI: 10.1542/peds.2015-0495 originally published online March 30, 2015;

Updated Information & Services	including high resolution figures, can be found at: http://pediatrics.aappublications.org/content/135/4/757
References	This article cites 12 articles, 3 of which you can access for free at: http://pediatrics.aappublications.org/content/135/4/757#BIBL
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Administration/Practice Management http://www.aappublications.org/cgi/collection/administration:practice_management_sub Interdisciplinary Teams http://www.aappublications.org/cgi/collection/interdisciplinary_teams_sub Obesity http://www.aappublications.org/cgi/collection/obesity_new_sub
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.aappublications.org/site/misc/Permissions.xhtml
Reprints	Information about ordering reprints can be found online: http://www.aappublications.org/site/misc/reprints.xhtml

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Primary Care Interventions for Pediatric Obesity: Need for an Integrated Approach

Cara B. Ebbeling and Richard C. Antonelli

Pediatrics 2015;135:757

DOI: 10.1542/peds.2015-0495 originally published online March 30, 2015;

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/135/4/757>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2015 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

