

Active Play Opportunities at Child Care

Pooja S. Tandon, MD, MPH^{a,b}, Brian E. Saelens, PhD^{a,b}, Dimitri A. Christakis, MD, MPH^{a,b}

abstract

BACKGROUND AND OBJECTIVES: Physical activity (PA) is important for children's health and development, yet preschoolers are not meeting PA recommendations. The objective of this study was to examine different PA opportunities at child care and how variation in indoor versus outdoor and free versus teacher-led opportunities relate to children's PA.

METHODS: An observational study of 98 children (mean age 4.5 years, 49% girls) from 10 child care centers. Classrooms were observed for at least 4 full days per center (total 50 days) to categorize time into (1) not an active play opportunity (APO); (2) naptime; (3) APO, outdoor free play; (4) APO, outdoor teacher-led; (5) APO, indoor free play; and (6) APO, indoor teacher-led. Children wore accelerometers during observations. Linear regression models examined the influence of APO categories on moderate-vigorous physical activity (MVPA) and sedentary time.

RESULTS: Children's activity was 73% sedentary, 13% light, and 14% MVPA. For 88% of time children did not have APOs, including 26% time as naptime. On average, 48 minutes per day were APOs (41% sedentary, 18% light, and 41% MVPA), 33 minutes per day were outdoors. The most frequent APO was outdoor free play (8% of time); outdoor teacher-led time was <1%. Children were more active and less sedentary outdoors versus indoors and during the child-initiated APOs (indoors and outdoors) versus teacher-led APOs.

CONCLUSIONS: Preschoolers were presented with significantly fewer than recommended opportunities for PA at child care. More APOs are needed for children to meet recommendations, particularly those that encourage more outdoor time, more teacher-led and child-initiated active play, and flexibility in naptime for preschoolers.



^aCenter for Child Health, Behavior, and Development, Seattle Children's Research Institute, Seattle, Washington; and ^bDepartment of Pediatrics, University of Washington, Seattle, Washington

Dr Tandon conceptualized and designed the study, oversaw the data analysis, and drafted the initial manuscript; Drs Saelens and Christakis advised Dr Tandon on the study design and data collection procedures, and critically reviewed and revised the manuscript; and all authors approved the final manuscript as submitted.

www.pediatrics.org/cgi/doi/10.1542/peds.2014-2750

DOI: 10.1542/peds.2014-2750

Accepted for publication Mar 12, 2015

Address correspondence to Pooja S. Tandon, MD, MPH, M/S CW8-6, PO Box 5371, Seattle Children's Research Institute, Seattle, WA 98145-5005. E-mail: pooja@uw.edu

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2015 by the American Academy of Pediatrics

FINANCIAL DISCLOSURE: The authors have indicated they have no financial relationships relevant to this article to disclose.

FUNDING: This research was supported by the American Heart Association's Beginning Grant-in-Aid program and a Career Development Award from the National Heart, Lung, and Blood Institute (K23 HL112950-01A1). Funded by the National Institutes of Health (NIH).

POTENTIAL CONFLICT OF INTEREST: The authors have indicated they have no potential conflicts of interest to disclose.

WHAT'S KNOWN ON THIS SUBJECT: Physical activity (PA) of preschoolers has been found to be highly correlated with their child care environment. Preschool-aged children are sedentary for most of their time at child care and most are not meeting PA recommendations.

WHAT THIS STUDY ADDS: Preschoolers were presented with significantly fewer than recommended PA opportunities at child care. More active play opportunities are needed to increase PA, including more outdoor time, more teacher-led and child-initiated active play, and flexibility in naptime for preschoolers.

Physical activity (PA) is important for children's health and development¹⁻³ and excessive sedentary behaviors are associated with poor health outcomes.⁴ Physical and sedentary activity have been found to track through childhood^{5,6} and there is a strong protective effect of PA on body fat accumulation from preschool age to adolescence.⁷ Whereas 3- to 5-year-olds tend to be more active than older children,⁸ they are not sufficiently active, as nearly half are not meeting minimal recommendations for daily PA.⁹⁻¹¹

Increasing children's PA in child care settings is critical given that approximately 5 million US children attend preschools or structured child care programs, with the average child spending more than 30 hours per week in this care.¹² Best practice guidelines encourage 60 minutes per day of teacher-led structured PA time, 60 minutes per day or more of unstructured free play time, and daily outdoor time, although how much each contributes to desirable levels of PA is not known.¹³ Pediatric health care providers have an important role in addressing and promoting active play opportunities (APOs), including those at child care.^{14,15}

The PA of preschoolers has been found to be highly correlated with their child care environment, thus providing policy and environment opportunities for intervention.¹⁶⁻¹⁸ In one study, the preschool attended explained 27% of the variance in activity levels, and accounted for more variance than did factors such as age, gender, and race/ethnicity.¹⁹ Certain child care characteristics have been positively associated with more PA, such as open space, portable play equipment, outdoor time, structured physical activities, and staff supportive of PA.^{17,20-23} A review of existing interventions to increase PA at child care found studies of varying quality, with 4 of 8 studies that assessed PA outcomes having had significant intervention effect,

typically through structured PA programs.²⁴ Although the child care setting provides multiple targets for intervention, the research on how to best organize child care time to optimize PA and decrease sedentary behavior is in its nascence.

The overall aim of this project was to examine different types of PA opportunities at child care and how variation in indoor versus outdoor and free versus teacher-led opportunities relate to children's PA. Such evidence could inform the targeting of policy and practices to increase PA in early childhood education environments.

METHODS

We conducted an observational study of 3- to 5-year-olds from a convenience sample of 10 licensed child care centers between 2012 and 2014. This study was approved by the Seattle Children's Hospital Institutional Review Board. All participating centers were in the Seattle area, where state licensing standards require a 10:1 child:teacher ratio for preschool classrooms. None were Head Start programs or funded by the city or state. Each center provided full day programming (at least 10 hours per day) and served infants through school-aged children. In 8 of the 10 programs, the directors had more than 10 years of child care experience. All of the centers had posted daily schedules that had at least 60 minutes per day scheduled for outdoor play time, divided into 2 sessions per day, but that time was not specified as "unstructured" or "teacher-led" active play time. All centers had outdoor play areas with fixed play structures and variable amounts of portable play equipment. They each had a dedicated indoor space (separate room from classroom) available for PA, including running and jumping, and assorted indoor play equipment. No center had any specific mention of weather-related outdoor play policies in either

their staff or parent handbooks, but teachers did have the option of using the indoor play area if the weather was not considered favorable for outdoor time. We recruited 8 to 10 children (ages 3-5) from each center to participate.

Measures

Location and Type of Activity

Data collection occurred in nonsummer months, as many preschools change their schedules to offer summer camp programming. A trained research assistant observed 1 classroom from each participating center for a minimum of 4 full days, but sometimes more frequently if we did not get 4 days of data on many children. The observer was present between 8 AM and 5 PM, which is when most children are in care and most of the structured curriculum occurs. Observation days were the same as the days when children wore accelerometers. The observer noted precise times when children were in various locations (eg, classroom, playroom, playground) and specifically noted whether they were indoors or outdoors. Before beginning data collection, all 3 research assistants simultaneously observed a nonparticipating preschool classroom for 2 days to establish consistency in coding.

Based on classroom activities and location, during the observation, the observers coded time into 6 mutually exclusive categories: (1) not an APO (eg, circle time, crafts, seated learning activities, meals); (2) naptime (children are required to sleep or lay on a mat for "quiet" activities); (3) APO, outdoor free play (children outdoors and could choose activities); (4) APO, outdoor teacher-led (teacher-initiated activities, such as running laps or active games, in which all children were expected to participate); (5) APO, indoor free play (active play encouraged indoors with children initiating activities; for example, climbing equipment, balls, in an indoor playroom); and (6) APO,

indoor teacher-initiated (teacher-led exercises [eg, yoga] in which all children were expected to participate). Any uncertainty about categorization was discussed soon after the visit with the research team to reach consensus. A category 7 was created for the few activities/times that still could not be clearly classified. This category comprised 1% of observed time.

Interrater reliability was assessed for 20% of the study observation days (1–2 days per center). This was done by having all 3 research assistants and the principal investigator independently recode the same days of original observation notes into the 7 categories mentioned previously. When the few category 7 times were removed, 100% concordance was seen among the observers.

Activity Levels

After obtaining parental consent and child assent, children wore Actigraph GT3X+ accelerometers (Actigraph, Pensacola, FL) on their right hip by using an adjustable belt during child care time on observation days. The research assistant placed the belt on the children soon after they arrived at the center and removed it at pick up, and this information was used to determine accelerometer wear time. The Actigraph has been validated and calibrated for use among children.²⁵ Data were collected in 15-second epochs, which is the current standard for preschool-aged children. Time-stamped accelerometer data were scored as sedentary, light, and moderate-to-vigorous (MVPA) by using Pate criteria.²⁶ Accelerometer data were then matched to observed categories of classroom activities. Although observation times and accelerometer wear time were similar, they were not equal (for example, if a child took the accelerometer off for naptime). Therefore, the sedentary, light, and MVPA percentages reflect a percentage of accelerometer wear time among children who were

observed to have any time in that category. Of note, the observed type of activity (APO or not) reflects the expectation of the preschoolers, and the activity level is quantified based on the children's actual movements. It is possible, for example, to have some sedentary activity during an APO and some MVPA during a non-APO or even naptime.

Demographics

At the time of consent, the date of birth and gender of children were collected. Parents also were given a short survey on their child's race/ethnicity, the highest educational attainment of adults at home, and household income.

Analysis

We summarized continuous measures by using means and SDs, and categorical variables by using frequencies and proportions. By using average time per day in each category for each child as the unit of analysis, we created separate mixed-effects linear regression models for the 2 primary outcomes (MVPA and sedentary time) to evaluate the influence of the 4 APO categories (outdoor free play, outdoor teacher-led, indoor free play, indoor teacher-led). We controlled for age, gender, and accelerometer wear time and accounted for clustering of children within child care centers and among observations from a single child as random effects. We also created separate mixed-effects linear regression models to test the effect of total outdoor versus indoor time and total outdoor APOs (free play plus teacher-led) versus indoor APOs on the outcomes of sedentary and MVPA, including the same covariates and random effects.

RESULTS

We conducted 50 total days of observations from 10 centers (with a minimum of 4 days per center) for an average of 7.2 hours per day. Our sample consisted of 98 children who

wore accelerometers with a mean age of 4.5 (range 3.1–5.8) years and 49% were girls. Additional demographic information was available for 82 participants and is presented in Table 1.

Sixty-two percent of the child care day (268.6 ± 44.8 minutes) could be categorized as not providing APOs for children, excluding naptime (see Table 2). Naptime constituted an additional 26% of the child care day (112.3 ± 32.2 minutes). On average, 48.5 ± 14.4 minutes of the day (12% of child care time) were considered APOs. Of the APOs, outdoor child-initiated free play was most common, followed by indoor teacher-led, indoor child-initiated, and then outdoor teacher-led. Children spent an average of 32.7 ± 13.8 minutes outdoors daily, with most of that time (99%) in child-initiated, free play activities. Children's mean MVPA was 55.2 ± 20.3 minutes per day with 34% attaining ≥60 minutes per day.

Regression analyses revealed that during the teacher-led indoor active time, children were more sedentary than during indoor or outdoor child-initiated free play times ($P < .001$). Children attained less MVPA during teacher-led indoor active time

TABLE 1 Demographic Characteristics of Participating Children ($n = 82$)

Characteristic	%
Race	
White	70
African American/black	6
>1 race	15
Asian/Pacific Islander/ Native American	6
Hispanic	13
Highest educational attainment in household	
Less than high school	2
Completed high school	24
Completed college	51
Graduate/professional degree	20
Household income, \$	
<29 000	29
30 000–49 000	16
50 000–69 000	5
70 000–89 000	12
≥90 000	32

TABLE 2 Characterization of Children's Opportunities for Active Play at Child Care

Category	By Observation		By Accelerometer ^a		
	Min/d, Mean (SD)	Mean Percentage of Day	Sedentary Min/d (%)	Light Min/d (%)	MVPA Min/d (%)
Total ^b	431.5 (57.4)		279.3 (73)	49.7 (13)	55.2 (14)
No opportunity for PA, total	381.0 (53.4)	88	258.0 (77)	40.7 (12)	35.1 (11)
Nonactive time	268.6 (44.8)	62	200.8 (75)	35.3 (13)	30.7 (12)
Naptime	112.3 (29.0)	26	57.2 (85)	5.4 (8)	4.5 (7)
APOs, total	48.5 (14.4)	12	19.8 (41)	8.7 (18)	19.8 (41)
Outdoor child-initiated	32.3 (13.6)	8	12.3 (38)	6.1 (19)	14.2 (44)
Indoor teacher-led	8.4 (8.9)	2	4.9 (49)	1.6 (17)	3.3 (34)
Indoor child-initiated	7.4 (8.4)	2	4.9 (39)	1.8 (17)	4.3 (44)
Outdoor teacher-led	0.4 (1.0)	0.1	1.1 (42)	0.3 (16)	0.5 (42)

^a Observation time and accelerometer wear time were similar but not equal. The sedentary, light, and moderate-vigorous PA are percentages of accelerometer wear time, among children who were observed to have any time in that category.

^b Total minutes include 2 minutes of a "category 7," which were activities that were not able to be classified into the other categories.

compared with the other active opportunities ($P < .001$). When comparing indoor and outdoor APOs (including both child-initiated and teacher-led), children spent less time sedentary (38% vs 44%, $P = .006$) and more time in MVPA (44% vs 39%, $P = .056$) outdoors. For indoor versus outdoor time in total (including APOs and non-APOs), children were less sedentary (38% vs 76%, $P < .001$) and spent more time in MVPA (44% vs 12%, $P < .001$) outdoors. Boys were more active and less sedentary than girls in all of the models (data not shown), but age and accelerometer wear time were not statistically significantly related to the outcomes.

DISCUSSION

This study found that for 88% of child care time, children were not presented opportunities for active play, so the finding that more than 70% of children's time was sedentary is not surprising. Compared with the recommended 120 minutes per day of PA time,¹³ our average of 48 minutes per day of APOs is considerably suboptimal. APOs did result in higher proportion of time in MVPA and less sedentary time, but constituted a small proportion of the day. Teacher-led PA opportunities were rare, and when present, children achieved lower percentages of time in MVPA compared with free play. Outdoor time was also low, but children were more active and less

sedentary when they were outdoors compared with indoors. These findings highlight the continued disconnection between PA recommendations and actual practices in early learning environments, which results in undesirable levels of sedentary behavior and PA.

Our findings suggest that child-initiated activity, indoors or outdoors, may result in less sedentary and more active play compared with teacher-led PA. Encouraging child-initiated free play of course has a myriad of additional benefits for young children's social, emotional, cognitive, and physical well-being.²⁷ Our results, however, seem inconsistent with previous studies, which found that structured, adult-led activities were associated with greater PA in children at child care.²⁴ Many of these programs used trained professionals, however (ie, not the classroom teacher), and implemented the intervention in a way that the structured activity was delivered with consistent frequency.^{28,29} Studies that use existing classroom staff to deliver interventions have been less successful and have reported challenges with intervention fidelity.^{30,31} Using professional physical education teachers at child care is not widely feasible and our results highlight what a limited role structured PA currently plays in many child care settings. On average, daily teacher-led active play time

amounted to 8.4 minutes indoors and 24 seconds outdoors, highlighting a significant gap from the recommended 60 minutes of teacher-led active time. Given that teacher-led activities also could have other potential benefits (eg, greater inclusion of all children, role modeling, integrating educational content), increasing child care providers' skills and comfort level in leading PA would be important to increasing APOs. In addition, it would be important to consider indoor and outdoor environments and equipment, which may better support teachers in encouraging movement both as part of the curriculum and during recess. How to best schedule and support APOs in terms of teacher-led or unstructured, frequency and duration, to maximize MVPA, needs to be better understood.

Our study also suggests that increasing outdoor time at child care is another route for promoting healthier activity levels. The National Association for the Education of Young Children recommends "daily" outdoor time³² as an accreditation criteria, but most states do not mandate this.³³ In our study, children had an average of 33 minutes of outdoor time per day (8% of their child care day), which was considerably less than the 60 minutes per day scheduled in all of the participating centers. A previous study found that preschoolers in 2 preschools in Sweden spent 46% of

their attendance time outdoors compared with 18% for children in 2 US preschools.³⁴ PA counts per minute was significantly higher outdoors versus indoors in both settings and the time spent in MVPA at preschool was very limited and predominantly outdoors. The investigators pointed out that there were more expectations for preschoolers in the United States to be sedentary (line up, sit down for circle time, nap, or engage in quiet activities during naptime). Those expectations plus the large difference in time indoors versus outdoors are likely reasons for higher MVPA in the Swedish children. A previous study using nationally representative data found that most preschool-aged children are not being taken outdoors daily by their parent(s).³⁵ Because many preschoolers spend considerable time in child care, usually during the daytime when outdoor play is most feasible, increasing outdoor time there is a strategy for increasing their overall PA. Children's outdoor time is a modifiable, environmental correlate of PA, although increasing outdoor time would likely require strategies to overcome perceived barriers around weather. Present findings suggest that both child-initiated and teacher-led outdoor time is conducive to children's PA.

We also found that children were less sedentary outdoors overall, and also during outdoor APOs compared with indoor APOs. The study that compared preschools in the United States and Sweden also found that children spent significantly less time engaged in sedentary behavior when they were outdoors compared with indoors. However, 2 previous studies did not find significant association between outdoor time and sedentary behavior.^{36,37} There is growing interest in targeting reductions in sedentary time as a distinct approach in health promotion,³⁸ as sedentary behavior is independently associated with adiposity in children and adults.³⁹⁻⁴²

There are several postulated mechanisms for the independent relationship between sedentary behavior and poor health outcomes, including physiologic changes that occur from loss of skeletal muscle contraction that lead to elevated glucose, triglycerides and free fatty acids and create a biochemical milieu conducive to cardiovascular risk factors.⁴³ Shifting some of this excessive sedentary behavior even to light PA, could be beneficial.

Naptime constituted a notable 26% of child care time in our study and is worthy of discussion with regard to age-appropriate expectations for napping. Daily rest time is often a licensing regulation in the United States, whereas other countries do not have a such a strict mandate, especially for older preschoolers.³⁴ Although there are certainly benefits to daytime sleep for young children⁴⁴ and many children may not be getting adequate nighttime sleep, some older preschoolers may no longer need daily scheduled long naps. Perhaps a shorter duration of rest time or the option to engage in different activities for some children would allow some of that large proportion of sedentary time to be redirected to APOs.

Excessive sedentary time and suboptimal active play may be having negative consequences on numerous aspects of children's development, health, and well-being. The paucity of APOs in child care is likely driven by multiple factors, including parent and teacher priorities. Parental preferences for academics and concerns about safety have been reported by child care providers as barriers they face in promoting PA in child care.⁴⁵ Teacher training and self-efficacy in implementing APOs is another possible barrier. Although the relationship between PA and learning is gaining considerable support through research in school-aged children,⁴⁶ it is not yet as well established in preschoolers. Increased

focus on the idea that active play and learning are not mutually exclusive may help reframe the importance of daily APOs to promote more supportive early learning practices and policies.

There are some limitations to our study. First, different levels of outdoor time and/or PA may occur on days when an observer is present compared with nonobserved days. However, relying on teacher report of indoor/outdoor time and PA opportunities also could include bias.⁴⁷ Second, this study was conducted at a limited number of licensed child care centers within 1 geographic area and therefore has limited generalizability to other child care settings and geographic regions. For our geographic area, we did capture a range of seasons and weather conditions over a 2-year period. Future studies could more precisely examine the relationship of seasonality and weather to outdoor play and PA.

CONCLUSIONS

Preschoolers were presented with significantly fewer than recommended opportunities for PA at child care. More APOs and fewer sedentary expectations at child care are needed for children to meet PA recommendations. Strategies to increase preschoolers' PA should be informed by research on which modifications would have the greatest benefit, including possibly encouraging more outdoor time, more teacher-led and child-initiated active play, and even some flexibility in mandated naptime for older preschoolers. In an environment in which sedentary behavior is increasingly common, pediatric health care providers can and should address PA opportunities of children within the context of all settings where children spend time.

ACKNOWLEDGMENTS

We acknowledge the research associates on the team: Albert Hsu,

Sefanit Ghezaghen, and Kelly Walters. We appreciate the statistical support provided by Wren Haaland and Chuan Zhou. Finally, we are grateful to the participating child care center staff, families, and children.

REFERENCES

1. Trost S. Active Education: Physical Education, Physical Activity and Academics Performance. Princeton, NJ: Robert Wood Johnson Foundation; 2009
2. Strong WB, Malina RM, Blimkie CJ, et al. Evidence based physical activity for school-age youth. *J Pediatr.* 2005;146(6): 732–737
3. Janssen I, Leblanc AG. Systematic review of the health benefits of physical activity and fitness in school-aged children and youth. *Int J Behav Nutr Phys Act.* 2010; 7(1):40
4. LeBlanc AG, Spence JC, Carson V, et al. Systematic review of sedentary behaviour and health indicators in the early years (aged 0-4 years). *Appl Physiol Nutr Metab.* 2012;37(4):753–772
5. Janz KF, Dawson JD, Mahoney LT. Tracking physical fitness and physical activity from childhood to adolescence: the Muscatine study. *Med Sci Sports Exerc.* 2000;32(7):1250–1257
6. Janz KF, Burns TL, Levy SM; Iowa Bone Development Study. Tracking of activity and sedentary behaviors in childhood: the Iowa Bone Development Study. *Am J Prev Med.* 2005;29(3):171–178
7. Moore LL, Gao D, Bradlee ML, et al. Does early physical activity predict body fat change throughout childhood? *Prev Med.* 2003;37(1):10–17
8. Pellegrini AD, Smith PK. Physical activity play: the nature and function of a neglected aspect of playing. *Child Dev.* 1998;69(3):577–598
9. Pate RR, Pfeiffer KA, Trost SG, Ziegler P, Dowda M. Physical activity among children attending preschools. *Pediatrics.* 2004;114(5):1258–1263
10. Fisher A, Reilly JJ, Kelly LA, et al. Fundamental movement skills and habitual physical activity in young children. *Med Sci Sports Exerc.* 2005; 37(4):684–688
11. Tucker P. The physical activity levels of preschool-aged children: A systematic review. *Early Child Res Q.* 2008;23: 547–558
12. Iruka IU, Carver PR. National Center for Education Statistics National Household Education Surveys Program of 2005: Initial Results From the 2005 NHES Early Childhood Program Participation Survey. Washington, DC: National Center for Education Statistics, Institute of Education Sciences, US Department of Education; 2006
13. National Association for Sport and Physical Education. *Active Start: A Statement of Physical Activity Guidelines for Children Birth to Five Years.* Reston, VA: National Association for Sport and Physical Education; 2002
14. American Academy of Pediatrics Committee on Early Childhood, Adoption, and Dependent Care. Quality early education and child care from birth to kindergarten. *Pediatrics.* 2005;115(1): 187–191
15. Hagan JF, Shaw JS, Duncan PM. *Bright Futures: Guidelines for Health Supervision of Infants, Children, and Adolescents.* Elk Grove Village, IL: American Academy of Pediatrics; 2008
16. Ward DS. Physical activity in young children: the role of child care. *Med Sci Sports Exerc.* 2010;42(3):499–501
17. Bower JK, Hales DP, Tate DF, Rubin DA, Benjamin SE, Ward DS. The childcare environment and children's physical activity. *Am J Prev Med.* 2008;34(1):23–29
18. Trost SG, Ward DS, Senso M. Effects of child care policy and environment on physical activity. *Med Sci Sports Exerc.* 2010;42(3):520–525
19. Pate RR, Mclver K, Dowda M, Brown WH, Addy C. Directly observed physical activity levels in preschool children. *J Sch Health.* 2008;78(8):438–444
20. Dowda M, Pate RR, Trost SG, Almeida MJ, Sirard JR. Influences of preschool policies and practices on children's physical activity. *J Community Health.* 2004;29(3):183–196
21. Willenberg LJ, Ashbolt R, Holland D, et al. Increasing school playground physical activity: a mixed methods study combining environmental measures and children's perspectives. *J Sci Med Sport.* 2010;13(2):210–216
22. Ridgers ND, Stratton G, Fairclough SJ, Twisk JWR. Long-term effects of playground markings and physical structures on children's recess physical activity levels. *Prev Med.* 2007;44(5): 393–397
23. Brown WH, Googe HS, Mclver KL, Rathel JM. Effects of teacher-encouraged physical activity on preschool playgrounds. *J Early Interv.* 2009;31(2): 126–145
24. Ward DS, Vaughn A, McWilliams C, Hales D. Interventions for increasing physical activity at child care. *Med Sci Sports Exerc.* 2010;42(3):526–534
25. Pate RR, Almeida MJ, Mclver KL, Pfeiffer KA, Dowda M. Validation and calibration of an accelerometer in preschool children. *Obesity (Silver Spring).* 2006; 14(11):2000–2006
26. Pate RR, O'Neill JR, Mitchell J. Measurement of physical activity in preschool children. *Med Sci Sports Exerc.* 2010;42(3):508–512
27. Milteer RM, Ginsburg KR; Council on Communications and Media; Committee on Psychosocial Aspects of Child and Family Health. The importance of play in promoting healthy child development and maintaining strong parent-child bond: focus on children in poverty. *Pediatrics.* 2012;129(1). Available at: www.pediatrics.org/cgi/content/full/129/1/e204
28. Eliakim A, Nemet D, Balakirski Y, Epstein Y. The effects of nutritional-physical activity school-based intervention on fatness and fitness in preschool children. *J Pediatr Endocrinol Metab.* 2007;20(6):711–718
29. Specker B, Binkley T, Fahrenwald N. Increased periosteal circumference remains present 12 months after an exercise intervention in preschool children. *Bone.* 2004;35(6):1383–1388
30. Alhassan S, Whitt-Glover MC. Intervention fidelity in a teacher-led program to promote physical activity in preschool-age children. *Prev Med.* 2014;69(suppl 1): S34–S36
31. Finch M, Wolfenden L, Morgan PJ, Freund M, Jones J, Wiggers J. A cluster randomized trial of a multi-level intervention, delivered by service staff, to increase physical activity of children attending center-based childcare. *Prev Med.* 2014;58:9–16
32. Center for Science in the Public Interest. Menu labeling. Updated February 4,

2009. Available at: www.cspinet.org/menulabeling. Accessed March 9, 2009
33. Benjamin SE, Craddock A, Walker EM, Slining M, Gillman MW. Obesity prevention in child care: a review of US state regulations. *BMC Public Health*. 2008;8:188
 34. Raustorp A, Pagels P, Boldemann C, Cosco N, Söderström M, Mårtensson F. Accelerometer measured level of physical activity indoors and outdoors during preschool time in Sweden and the United States. *J Phys Act Health*. 2012;9(6):801–808
 35. Tandon PS, Zhou C, Christakis DA. The frequency of outdoor play for preschool age children cared for at home-based child care settings. *Acad Pediatr*. 2012; 12(6):475–480
 36. Dolinsky DH, Brouwer RJ, Evenson KR, Siega-Riz AM, Østbye T. Correlates of sedentary time and physical activity among preschool-aged children. *Prev Chronic Dis*. 2011;8(6):A131
 37. King AC, Parkinson KN, Adamson AJ, et al; Gateshead Millennium Study Core Team. Correlates of objectively measured physical activity and sedentary behaviour in English children. *Eur J Public Health*. 2011;21(4): 424–431
 38. Marshall SJ, Ramirez E. Reducing sedentary behavior: A new paradigm in physical activity promotion. *Am J Lifestyle Med*. 2011;5(6):518–530
 39. Marshall SJ, Biddle SJ, Gorely T, Cameron N, Murdey I. Relationships between media use, body fatness and physical activity in children and youth: a meta-analysis. *Int J Obes Relat Metab Disord*. 2004;28(10):1238–1246
 40. Marshall SJ, Biddle SJH, Sallis JF, McKenzie TL, Conway TL. Clustering of sedentary behaviors and physical activity among youth: a cross-national study. *Pediatr Exerc Sci*. 2002;14: 401–417
 41. Fairclough SJ, Boddy LM, Hackett AF, Stratton G. Associations between children's socioeconomic status, weight status, and sex, with screen-based sedentary behaviours and sport participation. *Int J Pediatr Obes*. 2009; 4(4):299–305
 42. Thorp AA, Owen N, Neuhaus M, Dunstan DW. Sedentary behaviors and subsequent health outcomes in adults: a systematic review of longitudinal studies, 1996-2011. *Am J Prev Med*. 2011; 41(2):207–215
 43. Hamilton MT, Hamilton DG, Zderic TW. Exercise physiology versus inactivity physiology: an essential concept for understanding lipoprotein lipase regulation. *Exerc Sport Sci Rev*. 2004;32(4):161–166
 44. Kurdziel L, Duclos K, Spencer RMC. Sleep spindles in midday naps enhance learning in preschool children. *Proc Natl Acad Sci U S A*. 2013;110(43): 17267–17272
 45. Copeland KA, Sherman SN, Kendeigh CA, Kalkwarf HJ, Saelens BE. Societal values and policies may curtail preschool children's physical activity in child care centers. *Pediatrics*. 2012;129(2):265–274
 46. Institute of Medicine, Kohl HW III, Cook HD, eds. *Educating the Student Body: Taking Physical Activity and Physical Education to School*. Washington, DC: The National Academies Press; 2013
 47. Elgethun K, Yost MG, Fitzpatrick CTE, Nyerges TL, Fenske RA. Comparison of global positioning system (GPS) tracking and parent-report diaries to characterize children's time-location patterns. *J Expo Sci Environ Epidemiol*. 2007;17(2):196–206

Active Play Opportunities at Child Care

Pooja S. Tandon, Brian E. Saelens and Dimitri A. Christakis

Pediatrics 2015;135:e1425

DOI: 10.1542/peds.2014-2750 originally published online May 18, 2015;

Updated Information & Services

including high resolution figures, can be found at:
<http://pediatrics.aappublications.org/content/135/6/e1425>

References

This article cites 39 articles, 4 of which you can access for free at:
<http://pediatrics.aappublications.org/content/135/6/e1425#BIBL>

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
<http://www.aappublications.org/site/misc/Permissions.xhtml>

Reprints

Information about ordering reprints can be found online:
<http://www.aappublications.org/site/misc/reprints.xhtml>

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Active Play Opportunities at Child Care

Pooja S. Tandon, Brian E. Saelens and Dimitri A. Christakis

Pediatrics 2015;135:e1425

DOI: 10.1542/peds.2014-2750 originally published online May 18, 2015;

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/135/6/e1425>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2015 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

