



Skin-to-Skin Care for Term and Preterm Infants in the Neonatal ICU

Jill Baley, MD, COMMITTEE ON FETUS AND NEWBORN

abstract

“Kangaroo mother care” was first described as an alternative method of caring for low birth weight infants in resource-limited countries, where neonatal mortality and infection rates are high because of overcrowded nurseries, inadequate staffing, and lack of equipment. Intermittent skin-to-skin care (SSC), a modified version of kangaroo mother care, is now being offered in resource-rich countries to infants needing neonatal intensive care, including those who require ventilator support or are extremely premature. SSC significantly improves milk production by the mother and is associated with a longer duration of breastfeeding. Increased parent satisfaction, better sleep organization, a longer duration of quiet sleep, and decreased pain perception during procedures have also been reported in association with SSC. Despite apparent physiologic stability during SSC, it is prudent that infants in the NICU have continuous cardiovascular monitoring and that care be taken to verify correct head positioning for airway patency as well as the stability of the endotracheal tube, arterial and venous access devices, and other life support equipment.

FREE

This document is copyrighted and is property of the American Academy of Pediatrics and its Board of Directors. All authors have filed conflict of interest statements with the American Academy of Pediatrics. Any conflicts have been resolved through a process approved by the Board of Directors. The American Academy of Pediatrics has neither solicited nor accepted any commercial involvement in the development of the content of this publication.

Clinical reports from the American Academy of Pediatrics benefit from expertise and resources of liaisons and internal (American Academy of Pediatrics) and external reviewers. However, clinical reports from the American Academy of Pediatrics may not reflect the views of the liaisons or the organizations or government agencies that they represent.

The guidance in this report does not indicate an exclusive course of treatment or serve as a standard of medical care. Variations, taking into account individual circumstances, may be appropriate.

All clinical reports from the American Academy of Pediatrics automatically expire 5 years after publication unless reaffirmed, revised, or retired at or before that time.

www.pediatrics.org/cgi/doi/10.1542/peds.2015-2335

DOI: 10.1542/peds.2015-2335

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2015 by the American Academy of Pediatrics

BACKGROUND

“Kangaroo mother care” (KMC) was first described as an alternative method of caring for low birth weight infants in resource-limited countries, where neonatal mortality and infection rates are high because of overcrowded nurseries, inadequate staffing, and lack of equipment. In the original version of KMC, the infant is placed in continuous skin-to-skin contact in a vertical position between the mother’s breasts and beneath her clothes and is exclusively (or nearly exclusively) breastfed. A meta-analysis of 988 infants enrolled in 3 randomized controlled trials of continuous KMC begun in the first postnatal week in low- or middle-income countries found a 51% reduction in mortality among infants with a birth weight <2000 g (relative risk: 0.49 [95% confidence interval: 0.29–0.82]).¹ Although the methods of this review have come under question,² a Cochrane meta-analysis of 18 trials of continuous KMC begun before postnatal day 10 in infants with a birth weight <2500 g also showed significantly reduced mortality and morbidity at discharge or 40

to 41 weeks' postmenstrual age and at follow-up; it also found a decreased incidence of health care–related sepsis and an improvement in some measures of infant growth, breastfeeding, and mother–infant attachment.³ Thirteen of these 18 studies were conducted in low- to middle-income countries.

Intermittent skin-to-skin care (SSC) in NICUs in resource-rich countries differs from traditional KMC in that it is usually used for varying, shorter periods of time; can be offered to less stable and technology-supported infants; and can be performed by both parents. Intermittent SSC in resource-rich countries has not been associated with decreased mortality, although data are currently insufficient to determine an effect.³ However, it is widely offered to parents for other perceived benefits, such as enhancing attachment, parental self-esteem, and breastfeeding.^{4,5}

EVIDENCE

Benefits

The most substantial evidence of benefit from SSC is for breastfeeding. Individual randomized controlled trials and a systematic review have shown that intermittent SSC is associated with longer and more exclusive breastfeeding and higher volumes of expressed milk.^{6,7} The systematic review reported that short periods of SSC (up to 1 hour at all visits) increased the duration of any breastfeeding, variably reported by different studies as 1 month after discharge (relative risk: 4.76 [95% confidence interval: 1.19–19.10]) or for more than 6 weeks (relative risk: 1.95 [95% confidence interval: 1.03–3.70]) among clinically stable infants in industrialized nations.⁷ A number of studies have also indicated that SSC may improve a mother's attachment or bonding and her feeling of being needed by or comfortable with her infant.^{3,8–12}

In addition, SSC promotes the participation of the mother and father in the infant's care, strengthens the family role in the care of a fragile infant, and decreases feelings of helplessness.¹⁰ Mothers report less stress and more satisfaction with NICU care, and both parents are more responsive to their infant's cues.^{3,8–12}

The evidence is less clear for a beneficial effect regarding sleep and neurobehavioral maturation. One report found increased frontal brain activity during both quiet and active sleep, which is thought to be predictive of improved neurobehavioral outcomes.¹³ Other studies using electroencephalography and polysomnography data indicate that preterm infants who receive SSC have more mature sleep organization, with increased total and quiet sleep, decreased REM sleep and arousals from sleep, and an improvement in sleep cycling.^{14,15} They also appeared more alert and observant and spent less time crying. Two cohort studies found that infants receiving SSC demonstrated better autonomic regulation and maternal–infant interactions at term gestation, as well as higher scores on the Bayley Scales of Infant Development–Second Edition at 6 or 12 months of age.^{8,16} Of the infants enrolled in the second study, 117 were followed up to 10 years of age, and the authors reported that those who received SSC showed attenuated stress response, improved autonomic functioning, better-organized sleep, and better cognitive control.¹⁷

SSC has also been advocated for the nonpharmacologic management of procedural pain. A Cochrane review of the effect of SSC for relief of procedural pain concluded that it seemed to be effective for a single painful procedure such as a heel lance, as measured by using composite pain indicators.¹⁸ The review found that behavioral indicators of pain tended to favor SSC, whereas physiologic indicators were

generally not affected, suggesting possible observer bias in scoring behavioral indicators. However, small studies have reported reduced cortisol concentrations and decreased autonomic indicators of pain in preterm infants during SSC.^{19,20} The authors of the Cochrane review recommend confirmatory studies of previous findings and call for new studies examining optimal duration of SSC, use in different gestational age groups, effects of repeated use, and long-term effects.¹⁸

Risks

Investigators initially postulated that continuous KMC would promote colonization with maternal flora rather than resistant hospital flora. Consistent with this hypothesis, meta-analyses of randomized controlled trials in resource-limited countries have exhibited fewer episodes of sepsis, necrotizing enterocolitis, and pneumonia.^{1,3} However, infections may be spread among mothers, infants, and caregivers, particularly in multiple-bed units, as has been reported for respiratory syncytial virus and tuberculosis.^{21,22} Although a recent report described an association between SSC and development of methicillin-resistant *Staphylococcus aureus* infections among infants in 1 NICU (particularly those with very low birth weights), the authors did not believe that there was a causal relationship.²³ Parents should be monitored for skin infections and might need cleansing of the skin before infant contact. Some experts consider infants with open lesions (eg, open neural tube defects, abdominal wall defects) to be particularly at risk.

Most studies of physiologic stability during SSC have been performed on stable, nonintubated infants. One meta-analysis reported a statistically but not clinically significant increase in body temperature (0.22°C) and a decrease in oxygen saturation (0.60%) in 190 term and 326

preterm infants receiving SSC compared with incubator care.²⁴ These effects were most pronounced in nurseries in low- and middle-income settings and in cold environments. There was no change in heart rate before, during, or after SSC, and no difference was noted between preterm and term infants. Although 1 study of 22 infants reported an increase in desaturation and bradycardia during SSC,²⁵ other studies have shown no significant increase in desaturation, bradycardic or apneic events, or in oxygen consumption.^{26–28} Despite apparent physiologic stability during SSC, it is prudent that infants in the NICU be continuously monitored and that care be taken to verify correct head positioning for airway patency as well as the stability of the endotracheal tube, arterial and venous access devices, and other life support equipment. Any infant who requires careful temperature regulation or a high-humidity environment might have SSC delayed until he or she is more stable.

There may be resistance among health care providers regarding offering SSC. This resistance could stem from fear of harm to the infant or from lack of experience, time, or assistance to transfer the infant to the parent and/or monitor the infant's well-being. A nursing simulation training program may help promote acceptance of SSC.²⁹ Multiple guidelines for the provision of SSC have been published,^{30–33} and each facility needs to consider staffing, experience, and resources in the development of its institutional guidelines. Because SSC has been shown to be feasible and safe in the NICU in infants as young as 26 weeks' gestation,³⁴ with benefits for both parents and infants, facilities are encouraged to offer this care when possible.

IMPLICATIONS FOR CLINICAL PRACTICE

1. It has been shown that skin-to-skin care results in improved

- breastfeeding, milk production, parental satisfaction, and bonding.
2. Both parents can be encouraged to provide skin-to-skin care, with appropriate guidelines and protocols, for both preterm and term infants in the NICU.
3. Despite apparent physiologic stability during skin-to-skin care, it is prudent that infants in the NICU have continuous cardiovascular monitoring and that care be taken to monitor correct head positioning for airway patency as well as the stability of the endotracheal tube, arterial and venous access devices, and other life support equipment.

LEAD AUTHOR

Jill Baley, MD

COMMITTEE ON FETUS AND NEWBORN, 2014–2015

Kristi Watterberg, MD, Chairperson
James Cummings, MD
Eric Eichenwald, MD
Brenda Poindexter, MD
Dan L. Stewart, MD
Susan W. Aucott, MD
Karen M. Puopolo, MD
Jay P. Goldsmith, MD

LIAISONS

William Benitz, MD – *AAP Section on Perinatal Pediatrics*
Kasper S. Wang, MD – *AAP Section on Surgery*
Thierry Lacaze, MD – *Canadian Pediatric Society*
Jeffrey L. Ecker, MD – *American College of Obstetricians and Gynecologists*
Tonse N.K. Raju, MD, DCH – *National Institutes of Health*
Wanda Barfield, MD, MPH – *Centers for Disease Control and Prevention*
Erin Keels, MS, APRN, NNP-BC – *National Association of Neonatal Nurses*

STAFF

Jim Couto, MA
Address correspondence to: KWatterberg@salud.unm.edu

ABBREVIATIONS

KMC: kangaroo mother care
SSC: skin-to-skin care

REFERENCES

1. Lawn JE, Mwansa-Kambafwile J, Horta BL, Barros FC, Cousens S. 'Kangaroo mother care' to prevent neonatal deaths due to preterm birth complications. *Int J Epidemiol.* 2010;39(suppl 1):i144–i154
2. Sloan NL, Ahmed S, Anderson GC, Moore E. Comment on: 'kangaroo mother care' to prevent neonatal deaths due to preterm birth complications. *Int J Epidemiol.* 2011;40(2):521–525
3. Conde-Agudelo A, Díaz-Rossello JL. Kangaroo mother care to reduce morbidity and mortality in low birthweight infants. *Cochrane Database Syst Rev.* 2014;4(4):CD002771
4. Franck LS, Bernal H, Gale G. Infant holding policies and practices in neonatal units. *Neonatal Netw.* 2002; 21(2):13–20
5. Field T, Hernandez-Reif M, Feijo L, Freedman J. Prenatal, perinatal and neonatal stimulation: a survey of neonatal nurseries. *Infant Behav Dev.* 2006;29(1):24–31
6. Hake-Brooks SJ, Anderson GC. Kangaroo care and breastfeeding of mother-preterm infant dyads 0-18 months: a randomized, controlled trial. *Neonatal Netw.* 2008;27(3):151–159
7. Renfrew MJ, Craig D, Dyson L, et al. Breastfeeding promotion for infants in neonatal units: a systematic review and economic analysis. *Health Technol Assess.* 2009;13(40):1–146, iii–iv
8. Ohgi S, Fukuda M, Moriuchi H, et al. Comparison of kangaroo care and standard care: behavioral organization, development, and temperament in healthy, low-birth-weight infants through 1 year. *J Perinatol.* 2002;22(5):374–379
9. Charpak N, Ruiz JG, Zupan J, et al. Kangaroo mother care: 25 years after. *Acta Paediatr.* 2005;94(5):514–522
10. Nyqvist KH, Anderson GC, Bergman N, et al. Towards universal kangaroo mother care: recommendations and report from the first European conference and Seventh International Workshop on kangaroo mother care. *Acta Paediatr.* 2010;99(6):820–826
11. Johnson AN. The maternal experience of kangaroo holding. *J Obstet Gynecol Neonatal Nurs.* 2007;36(6):568–573

12. Tessier R, Charpak N, Giron M, Cristo M, de Calume ZF, Ruiz-Peláez JG. Kangaroo mother care, home environment and father involvement in the first year of life: a randomized controlled study. *Acta Paediatr*. 2009;98(9):1444–1450
13. Welch MG, Myers MM, Grieve PG, et al; FNI Trial Group. Electroencephalographic activity of preterm infants is increased by Family Nurture Intervention: a randomized controlled trial in the NICU. *Clin Neurophysiol*. 2014;125(4):675–684
14. Ludington-Hoe SM, Johnson MW, Morgan K, et al. Neurophysiologic assessment of neonatal sleep organization: preliminary results of a randomized, controlled trial of skin contact with preterm infants. *Pediatrics*. 2006;117(5). Available at: www.pediatrics.org/cgi/content/full/117/5/e909
15. Feldman R, Eidelman AI. Skin-to-skin contact (kangaroo care) accelerates autonomic and neurobehavioural maturation in preterm infants. *Dev Med Child Neurol*. 2003;45(4):274–281
16. Feldman R, Eidelman AI, Sirota L, Weller A. Comparison of skin-to-skin (kangaroo) and traditional care: parenting outcomes and preterm infant development. *Pediatrics*. 2002;110(1 pt 1):16–26
17. Feldman R, Rosenthal Z, Eidelman AI. Maternal-preterm skin-to-skin contact enhances child physiologic organization and cognitive control across the first 10 years of life. *Biol Psychiatry*. 2014;75(1):56–64
18. Johnston C, Campbell-Yeo M, Fernandes A, Inglis D, Streiner D, Zee R. Skin-to-skin care for procedural pain in neonates. *Cochrane Database Syst Rev*. 2014;1(1):CD008435
19. Cong X, Ludington-Hoe SM, Walsh S. Randomized crossover trial of kangaroo care to reduce biobehavioral pain responses in preterm infants: a pilot study. *Biol Res Nurs*. 2011;13(2):204–216
20. Cong X, Cusson RM, Walsh S, Hussain N, Ludington-Hoe SM, Zhang D. Effects of skin-to-skin contact on autonomic pain responses in preterm infants. *J Pain*. 2012;13(7):636–645
21. Visser A, Delpont S, Venter M. Molecular epidemiological analysis of a nosocomial outbreak of respiratory syncytial virus associated pneumonia in a kangaroo mother care unit in South Africa. *J Med Virol*. 2008;80(4):724–732
22. Heyns L, Gie RP, Goussard P, Beyers N, Warren RM, Marais BJ. Nosocomial transmission of Mycobacterium tuberculosis in kangaroo mother care units: a risk in tuberculosis-endemic areas. *Acta Paediatr*. 2006;95(5):535–539
23. Sakaki H, Nishioka M, Kanda K, Takahashi Y. An investigation of the risk factors for infection with methicillin-resistant Staphylococcus aureus among patients in a neonatal intensive care unit. *Am J Infect Control*. 2009;37(7):580–586
24. Mori R, Khanna R, Pledge D, Nakayama T. Meta-analysis of physiological effects of skin-to-skin contact for newborns and mothers. *Pediatr Int*. 2010;52(2):161–170
25. Bohnhorst B, Gill D, Dördelmann M, Peter CS, Poets CF. Bradycardia and desaturation during skin-to-skin care: no relationship to hyperthermia. *J Pediatr*. 2004;145(4):499–502
26. Bauer J, Sontheimer D, Fischer C, Linderkamp O. Metabolic rate and energy balance in very low birth weight infants during kangaroo holding by their mothers and fathers. *J Pediatr*. 1996;129(4):608–611
27. Heimann K, Vaessen P, Peschgens T, Stanzel S, Wenzl TG, Orlikowsky T. Impact of skin to skin care, prone and supine positioning on cardiorespiratory parameters and thermoregulation in premature infants. *Neonatology*. 2010;97(4):311–317
28. Ludington-Hoe SM, Anderson GC, Swinth JY, Thompson C, Hadeed AJ. Randomized controlled trial of kangaroo care: cardiorespiratory and thermal effects on healthy preterm infants. *Neonatal Netw*. 2004;23(3):39–48
29. Hendricks-Munoz KD, Mayers RM. A neonatal nurse training program in kangaroo mother care (KMC) decreases barriers to KMC utilization in the NICU. *Am J Perinatol*. 2014;31(11):987–992
30. Kledzik T. Holding the very low birth weight infant: skin-to-skin techniques. *Neonatal Netw*. 2005;24(1):7–14
31. DiMenna L. Considerations for implementation of a neonatal kangaroo care protocol. *Neonatal Netw*. 2006;25(6):405–412
32. Welch MG, Hofer MA, Stark RI, et al; FNI Trial Group. Randomized controlled trial of family nurture intervention in the NICU: assessments of length of stay, feasibility and safety. *BMC Pediatr*. 2013;13:148–162
33. Ludington-Hoe SM, Ferreira C, Swinth J, Ceccardi JJ. Safe criteria and procedure for kangaroo care with intubated preterm infants. *J Obstet Gynecol Neonatal Nurs*. 2003;32(5):579–588
34. Bier JA, Ferguson AE, Morales Y, et al. Comparison of skin-to-skin contact with standard contact in low-birth-weight infants who are breast-fed. *Arch Pediatr Adolesc Med*. 1996;150(12):1265–1269

Skin-to-Skin Care for Term and Preterm Infants in the Neonatal ICU
Jill Baley and COMMITTEE ON FETUS AND NEWBORN
Pediatrics 2015;136;596
DOI: 10.1542/peds.2015-2335 originally published online August 31, 2015;

Updated Information & Services

including high resolution figures, can be found at:
<http://pediatrics.aappublications.org/content/136/3/596>

References

This article cites 34 articles, 6 of which you can access for free at:
<http://pediatrics.aappublications.org/content/136/3/596#BIBL>

Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):

Current Policy

http://www.aappublications.org/cgi/collection/current_policy

Committee on Fetus & Newborn

http://www.aappublications.org/cgi/collection/committee_on_fetus_newborn

Fetus/Newborn Infant

http://www.aappublications.org/cgi/collection/fetus:newborn_infant_sub

Neonatology

http://www.aappublications.org/cgi/collection/neonatology_sub

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:

<http://www.aappublications.org/site/misc/Permissions.xhtml>

Reprints

Information about ordering reprints can be found online:

<http://www.aappublications.org/site/misc/reprints.xhtml>

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®



PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Skin-to-Skin Care for Term and Preterm Infants in the Neonatal ICU

Jill Baley and COMMITTEE ON FETUS AND NEWBORN

Pediatrics 2015;136;596

DOI: 10.1542/peds.2015-2335 originally published online August 31, 2015;

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/136/3/596>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 345 Park Avenue, Itasca, Illinois, 60143. Copyright © 2015 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®

