

Hospital Utilization Among Children With the Highest Annual Inpatient Cost

Alon Peltz, MD, MBA,^{a,b,c} Matt Hall, PhD,^d David M. Rubin, MD, MSCE,^{e,f} Kenneth D. Mandl, MD, MPH,^{g,h} John Neff, MD,ⁱ Mark Brittan, MD, MPH,^j Eyal Cohen, MD, MSc,^k David E. Hall, MD,^l Dennis Z. Kuo, MD, MHS,^m Rishi Agrawal, MD, MPH,^{n,o} Jay G. Berry, MD, MPH^{a,p}

abstract

BACKGROUND AND OBJECTIVES: Children who experience high health care costs are increasingly enrolled in clinical initiatives to improve their health and contain costs. Hospitalization is a significant cost driver. We describe hospitalization trends for children with highest annual inpatient cost (CHIC) and identify characteristics associated with persistently high inpatient costs in subsequent years.

METHODS: Retrospective study of 265 869 children age 2 to 15 years with ≥ 1 admission in 2010 to 39 children's hospitals in the Pediatric Health Information System. CHIC were defined as the top 10% of total inpatient costs in 2010 ($n = 26\ 574$). Multivariate regression and regression tree modeling were used to distinguish individual characteristics and interactions of characteristics, respectively, associated with persistently high inpatient costs (≥ 80 th percentile in 2011 and/or 2012).

RESULTS: The top 10% most expensive children (CHIC) constituted 56.9% (\$2.4 billion) of total inpatient costs in 2010. Fifty-eight percent ($n = 15\ 391$) of CHIC had no inpatient costs in 2011 to 2012, and 27.0% ($n = 7180$) experienced persistently high inpatient cost. Respiratory chronic conditions (odds ratio [OR] = 3.0; 95% confidence interval [CI], 2.5–3.5), absence of surgery in 2010 (OR = 2.0; 95% CI, 1.8–2.1), and technological assistance (OR = 1.6; 95% CI, 1.5–1.7) were associated with persistently high inpatient cost. In regression tree modeling, the greatest likelihood of persistence (65.3%) was observed in CHIC with ≥ 3 hospitalizations in 2010 and a chronic respiratory condition.

CONCLUSIONS: Most children with high children's hospital inpatient costs in 1 year do not experience hospitalization in subsequent years. Interactions of hospital use and clinical characteristics may be helpful to determine which children will continue to experience high inpatient costs over time.

^aDepartment of Medicine, Boston Children's Hospital, Boston, Massachusetts; and ^bComputational Health Informatics Program, Boston Children's Hospital, Boston, Massachusetts; and ^cDepartment of Pediatrics, Boston Medical Center, Boston, Massachusetts; and ^dRobert Wood Johnson Foundation Clinical Scholars Program, Yale University, New Haven, Connecticut; ^eChildren's Hospital Association, Overland Park, Kansas; ^fPolicyLab at The Children's Hospital of Philadelphia, Philadelphia, Pennsylvania; and ^gDepartment of Pediatrics, Perelman School of Medicine at the University of Pennsylvania, Philadelphia, Pennsylvania; ^hDepartment of Biomedical Informatics, Harvard Medical School, Boston, Massachusetts; ⁱDepartment of Pediatrics, Harvard Medical School, Boston, Massachusetts; ^jDepartment of Pediatrics, Seattle Children's Hospital, University of Washington, Seattle, Washington; ^kDepartment of Pediatrics, Children's Hospital Colorado, University of Colorado School of Medicine, Aurora, Colorado; ^lDepartment of Pediatrics, Hospital for Sick Children, University of Toronto, Toronto, Ontario, Canada; ^mDepartment of Pediatrics, Monroe Carell Jr. Children's Hospital at Vanderbilt, Vanderbilt University School of Medicine, Nashville, Tennessee; ⁿDepartment of Pediatrics, Center for Applied Research and Evaluation, University of Arkansas for Medical Sciences, Arkansas Children's Hospital, Little Rock, Arkansas; and ^oDepartment of Pediatrics, Northwestern University, Feinberg School of Medicine, Chicago, Illinois; and ^pDivision of Hospital-Based Medicine, Lurie Children's Hospital of Chicago, Chicago, Illinois;

Drs Peltz and Rubin participated in the conceptualization and design of the study, made a substantial contribution to the analysis and interpretation of data, and drafted, critically

WHAT'S KNOWN ON THIS SUBJECT: Children who experience high health care costs are increasingly enrolled in clinical initiatives to improve their quality of care and contain their health care costs over time. However, little is known about longitudinal trends in hospital costs of these children.

WHAT THIS STUDY ADDS: Most children with high inpatient costs in 1 year are not hospitalized in subsequent years. Children with a respiratory chronic condition and multiple hospitalizations were most likely to experience persistently high inpatient costs.

To cite: Peltz A, Hall M, Rubin DM, et al. Hospital Utilization Among Children With the Highest Annual Inpatient Cost. *Pediatrics*. 2016;137(2):e20151829

Pediatric health care costs are highly concentrated among a small group of children with chronic conditions.¹⁻⁶

In recent years, there has been a significant rise in innovative clinical programs targeting this population of children.⁷ Health plans, children's hospitals, and other entities are enrolling high-cost children into care management programs to improve their health.⁸ Some of these programs have reported a major impact on health care utilization and costs, suggesting that they may help interrupt the cycle of persistent health care spending that the children would have experienced otherwise.⁹⁻¹³

As for their adult counterparts,¹⁴ hospital care is a significant contributor to the overall cost accrued by children with chronic conditions.^{2,5,6} Many children who experience high health care costs incur frequent hospital admissions and lengthy hospitalizations.¹⁵⁻¹⁸ Previous studies have reported that hospital care accounts for as much as 80% of total health care spending in children with severe chronic conditions.^{2,5,6} Spending on other health services, such as emergency department care, medications, and primary care, is much smaller.¹⁹ Hospital costs and how the costs are distributed (eg, to ICU or surgical care) vary significantly by the chronic conditions experienced by the children.¹⁷

The financial value proposition of many care management programs for patients with chronic conditions centers on containing and reducing inpatient utilization. It is believed that some of the hospital costs incurred by these patients might have been avoided with better quality of care and better care coordination.^{20,21} However, there is little information on longitudinal trends in hospital costs for children and on the predictors of persistently high hospital costs.^{2,21} Without this information, it is difficult to

target the correct patients to help and to discern the true impact of interventions.

A better understanding of the characteristics of children who incur high inpatient costs and of when the high costs are most likely to persist may enable care management initiatives to focus enrollment on children who are most likely to benefit from them. Therefore, the objectives of this study are to describe trends in hospital use for children with the highest annual inpatient cost (CHIC) in children's hospitals throughout the United States and to distinguish characteristics of children who experience persistently high inpatient costs over time.

METHODS

Study Design, Setting, and Population

We conducted a retrospective cohort analysis of 265 869 children ages 2 to 15 years who were hospitalized at least once between January 1, 2010 and December 31, 2010 in 1 of 39 freestanding children's hospitals from 24 states and the District of Columbia in the Pediatric Health Information System (PHIS) data set. All hospitals participating in PHIS are members of the Children's Hospital Association.²² PHIS includes a unique patient identifier that permits the measurement of repeated hospitalizations for the same patient within the same hospital but not at different PHIS hospitals. The Boston Children's Hospital Institutional Review Board approved this study with a waiver for informed consent.

Identifying Children With the Highest Inpatient Costs in 2010

We categorized each child into 1 of 10 cost deciles based on their total inpatient costs accrued in 2010. Hospital costs included room and board, laboratory, imaging, pharmacy, medical supplies, and

ancillary services. We did not include professional fees (eg, physician consultation) because PHIS does not contain these data. Costs were calculated from charges based on annual hospital specific cost-to-charge ratios. CHIC were the top 10% of children who incurred the highest total inpatient costs in 2010. All other children (non-CHIC) were the remaining 90% of children in the cohort. We used the most expensive 10% as a threshold based on previous surveys of high-cost children,²³ previous evaluations of adult Medicare beneficiaries,²⁴ and the distribution of costs across hospitalized children (Fig 1).

Main Outcome Measures

The main outcome measures were trends in hospital use for CHIC and non-CHIC and the persistence of high inpatient cost. For each child who was hospitalized at least once in 2010, we measured their hospital utilization back to January 1, 2008 and forward to December 31, 2012. Hospital utilization included the number of admissions, hospital days, and hospital costs. Children aged <2 years on January 1, 2010 were excluded from the analysis because they were not able to experience 2 full years of exposure for hospitalization before 2010. Children aged >15 years on January 1, 2010 were excluded to create an age ceiling of <18 years at the end of the study period. Children aged ≥18 years may be more likely to transfer their inpatient care to an adult hospital. We excluded children who experienced in-hospital mortality during any year of the study period, because they did not have the opportunity to experience hospitalization throughout the study period.

Characteristics Associated With Persistent High Inpatient Cost

We assessed patients' demographic, clinical, and hospitalization characteristics from 2008 to 2010

that might correlate with persistent high inpatient cost in 2011 and/or 2012.

Demographic Characteristics

Demographic characteristics included age, gender, race or ethnicity, and insurance (public, private, and other).

Clinical Characteristics

Clinical characteristics included the number, type, and complexity of patients' chronic conditions and assistance with medical technology. These characteristics were identified with International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM) codes. To evaluate for the number of comorbid chronic health conditions, we used the Chronic Condition Indicator classification system, developed by the Agency for Healthcare Research and Quality, to identify the presence of a chronic condition.²⁵ We used complex chronic conditions (CCCs) to identify children with a complex medical condition expected to last ≥ 12 months and involve either severe single-organ disease or multiple-organ dysfunction.²⁶ Patients assisted with medical technology, defined as a device used to overcome a severe limitation in physiologic functioning (eg, tracheostomy tube, gastrostomy tube), were identified with an established set of ICD-9-CM codes known to correlate with hospital resource use.²⁷⁻²⁹

Hospitalization Characteristics

We assessed the reasons for hospitalization in 2010, distinguishing medical from surgical admissions by using 3M's All Patient Refined Diagnosis Related Groups (3M Health Information Systems, Wallingford, CT).³⁰ We assessed for hospitalizations related to ambulatory care sensitive conditions, defined as conditions (eg, asthma, dehydration) for which hospitalization may be avoided with high-quality outpatient care.³¹




FIGURE 1

Hospital utilization in 2010 by patients' inpatient cost decile. Median annual hospitalizations, hospital bed days, and aggregate inpatient costs in 2010, and percentage of total, for each of the 10 cost groups (ie, deciles) based on their total inpatient costs accrued in 2010.

Statistical Analysis

In bivariate analysis, we used Wilcoxon rank sum and χ^2 tests to distinguish the demographic, clinical, and hospitalization characteristics of CHIC. In multivariate analysis, we assessed these relationships by using a generalized estimating equation (GEE) clustered by hospital. We derived the GEE model by simultaneously entering all the covariates and retaining them regardless of statistical significance.

We also used classification and regression tree (CART) modeling with binary split and postpruning goodness-of-fit rules to assess the most statistically significant interactions between multiple characteristics and to assess which unique patients had the highest likelihood of persistent high inpatient cost.³² The CART model complements the GEE model by assessing patient-level probabilities of persistent high inpatient costs for every possible interaction of characteristics (eg, 833 possible 3-level interactions of characteristics assessed in the current study). Statistical significance was set at $P < .05$. Statistical Analysis Software version 9.3 (SAS Institute, Inc, Cary, NC) and R version

3.0.2 (Vienna, Austria), with the PARTY package, were used for the analyses.³³

We conducted sensitivity analyses by excluding 2 populations who might be likely to experience episodic high-cost hospitalization and therefore limit our ability to distinguish children with persistently high inpatient costs: children undergoing a single surgery in 2010 and children with oncologic diagnoses.

RESULTS

Study Population

There were 265 869 children ages 2 to 15 years hospitalized in 2010 across the 39 children's hospitals in the cohort. In total, these children experienced 344 050 hospital admissions, accounted for 1.3 million hospital bed days, and incurred \$4.2 billion in hospital costs. Median age at admission was 8 years (interquartile range [IQR] 4-12). Fifty-four percent ($n = 144 685$) were male, 49.8% ($n = 132 362$) were non-Hispanic white, and 53.2% ($n = 141 468$) used public insurance. Sixty-eight percent ($n = 180 662$) had ≥ 1 chronic condition, 30.6%

TABLE 1 Characteristics of CHIC Compared With Children Without High Inpatient Costs in 2010

Characteristics ^a	All Children	CHIC in 2010 ^b	Children Without Highest Inpatient Cost in 2010
Number of patients, <i>N</i> (%)	265 869	26 574 (10.0)	239 295 (90.0)
Age, median (IQR), y	8 (4, 12)	10 (5, 13)	8 (4, 12)
Male gender, <i>N</i> (%)	144 685 (54.4)	13 948 (52.5)	130 737 (54.6)
Public insurance, <i>N</i> (%)	141 468 (53.2)	14 215 (53.5)	127 253 (53.2)
Race or ethnicity, <i>N</i> (%)			
Non-Hispanic white	132 362 (49.8)	14 145 (53.2)	118 217 (49.4)
Non-Hispanic black	55 575 (20.9)	4 797 (18.1)	50 778 (21.2)
Hispanic	48 257 (18.2)	4 495 (16.9)	43 762 (18.3)
Asian	5 399 (2.0)	589 (2.2)	4 810 (2.0)
Other	24 276 (9.1)	2 548 (9.6)	21 728 (9.1)
No. of chronic conditions, <i>N</i> (%)			
None	85 207 (32.0)	1 123 (4.2)	84 084 (35.1)
1	93 916 (35.3)	4 782 (18.0)	89 134 (37.2)
2	44 127 (16.6)	5 368 (20.2)	38 759 (16.2)
3+	42 619 (16.0)	15 301 (57.6)	27 318 (11.4)
Chronic condition type, <i>N</i> (%)			
Noncomplex, chronic	100 535 (37.8)	4 289 (16.1)	96 246 (40.2)
Complex, chronic	81 450 (30.6)	21 217 (79.8)	60 233 (25.2)
Complex Chronic Conditions, <i>N</i> (%) ^c			
Cardiology	15 421 (5.8)	5 457 (20.5)	9 964 (4.2)
Gastrointestinal	4 673 (1.8)	1 329 (5.0)	3 344 (1.4)
Hematology and immunology	9 042 (3.4)	1 786 (6.7)	7 256 (3.0)
Malignancy	12 159 (4.6)	5 430 (20.4)	6 729 (2.8)
Metabolic	6 060 (2.3)	2 564 (9.6)	3 496 (1.5)
Neurology	31 981 (12.0)	7 181 (27.0)	24 800 (10.4)
Renal	4 389 (1.7)	1 326 (5.0)	3 063 (1.3)
Respiratory	5 437 (2.0)	1 908 (7.2)	3 529 (1.5)
Other	15 801 (5.9)	5 692 (21.4)	10 109 (4.2)
Technological assistance, <i>N</i> (%)	26 242 (9.9)	9 067 (34.1)	17 175 (7.2)

^a All *P* values <.001 when we compared CHIC in 2010 and children without highest inpatient costs in 2010 as calculated by using either Wilcoxon rank sum or χ^2 tests.

^b CHIC are defined as the 10% most expensive children hospitalized between January 1, 2010 and December 31, 2010.

^c Patients may have zero, one, or more than one CCC. Column totals will not add up to 100%.

(*n* = 81 450) had a CCC, and 9.9% (*n* = 26 242) had technological assistance (Table 1).

Children With Highest Inpatient Cost in 2010

In 2010 the top 10% most expensive children (CHIC) accounted for 20.8% (*n* = 71 616) of all hospital admissions, 48.8% (*n* = 625 832 days) of all hospital days, and 56.9% (*n* = \$2.4 billion) of all hospital costs (Fig 1).

CHIC experienced a median of 2 hospitalizations (IQR 1–3) in 2010. Seventy percent (*n* = 51 097) of hospitalizations experienced by CHIC were for medical treatment, and 30.2% (*n* = 22 068) were for surgery. Chemotherapy, sickle cell crisis, and seizure were among the most common medical reasons for admission when all causes for

hospitalization were considered. Spinal fusion for scoliosis and cerebrospinal fluid ventricular shunt operations were the most common surgeries. Only 8.5% (*n* = 6206) of hospitalizations experienced by CHIC were for ambulatory care sensitive conditions. Seizures and pneumonia were the most common ambulatory care sensitive reasons for hospitalization.

Compared with all other children, CHIC had an older median age (10 years [IQR 5–13] vs 8 years [IQR 4–12], *P* < .001) and a higher percentage of non-Hispanic white race and ethnicity (53.2% vs 49.4%, *P* < .001). CHIC had a higher prevalence of ≥ 1 comorbid chronic health condition (95.8% vs 64.9%, *P* < .001) and ≥ 3 comorbid chronic health conditions (57.6% vs 11.4%, *P* < .001). CHIC also had a higher

prevalence of CCCs (79.8% vs 25.2%, *P* < .001) and technological assistance (34.1% vs 7.2%, *P* < .001) (Table 1).

Trends in Hospital Utilization (2008–2012)

Both CHIC and children with less inpatient cost in 2010 experienced significantly fewer median admissions, hospital bed days, and hospital costs in the 2 preceding and subsequent years (Fig 2): The median number of admissions, hospital days, and hospital costs were 0 in 2008 to 2009 and in 2011 to 2012 for both groups. Regarding inpatient costs for CHIC, 57.9% (*n* = 15 391) had no inpatient costs in 2011 or 2012 (because they had no hospital admissions); 15.1% (*n* = 4003) had inpatient costs <80th percentile in 2011 or 2012. More than one-quarter (27.0%, *n* = 7180) of CHIC




FIGURE 2 Trends in hospital utilization of CHIC and children without highest inpatient cost ($n = 265\,869$). Median annual hospitalizations, bed days, and aggregate inpatient costs in 2008 to 2012 for all children hospitalized in 2010. Black circles denote CHIC. Gray squares denote all other children. Vertical bars denote the IQRs.

experienced persistent high inpatient costs (≥ 80 th percentile) in 2011 or 2012 (Fig 3). Similar longitudinal trends in utilization were observed when we excluded children with an oncologic diagnosis and children undergoing a high-cost surgery in 2010.

Distinguishing Children With Persistent High Inpatient Cost

In bivariate analysis, CHIC in 2010 with persistent high inpatient costs in 2011 or 2012 had a higher




FIGURE 3 Comparison of hospital utilization in 2008 to 2012 for persistent high-cost and nonpersistent high-cost children ($n = 26\,574$). Median annual hospitalizations, bed days, and aggregate inpatient costs in 2008 to 2012 for CHIC in 2010. Black circles denote CHIC with persistently high cost. Gray squares denote CHIC with nonpersistently high cost. Vertical bars denote IQRs.

percentage of public insurance (60.1% vs 51.1%, $P < .001$), CCCs (90.6% vs 75.9%, $P < .001$), and technological assistance (49.4% vs 28.5%, $P < .001$) when compared with CHIC in 2010 without persistent high inpatient cost. CHIC with persistent high inpatient cost also had a higher percentage of ≥ 3 comorbid chronic health conditions (77.9% vs 50%) compared with CHIC with no persistent high inpatient cost (Table 2).

In multivariable regression, CHIC with the highest likelihood of persistent high cost were those with a respiratory CCC (adjusted odds ratio [OR] 3.0; 95% confidence interval [CI], 2.5–3.5; $P < .001$), those who did not undergo surgery in 2010 (OR 2.0; 95% CI, 1.8–2.1; $P < .001$), and those with technological assistance (OR 1.6; 95% CI, 1.5–1.7; $P < .001$) (Table 2). The presence of ≥ 3 comorbid chronic health conditions was not significantly associated with persistent high inpatient costs in 2011 or 2012 ($P = .1$).

In CART analysis we assessed 833 potential interactions of patient characteristics. The greatest likelihood (65.3%) of persistent high costs was observed in CHIC with a respiratory chronic condition and ≥ 3 hospitalizations in 2010 (Fig 4). This combination of characteristics was observed in 3.4% ($n = 930$) of CHIC with persistently high inpatient cost. The second greatest likelihood (63.9%) of persistently high inpatient cost was observed in CHIC with a respiratory chronic condition, with < 3 hospitalizations in 2010 and no surgery in 2010. Cystic fibrosis and bronchopulmonary dysplasia were the most common respiratory CCCs experienced by these children.

DISCUSSION

Consistent with previous studies,^{14,23,24,34} our findings suggest that hospital costs are concentrated among a small group of patients: 10% of children with highest inpatient costs in children's hospitals in 2010 accounted for more than one-half of total hospital expenditures. Most children with the highest inpatient costs in 2010 were not admitted again to the same children's hospital over the next 2 years, nor had they been admitted in the 2 previous years. Only one-fourth of the children with highest annual inpatient costs in 2010 went on to experience inpatient costs in the top 20% in

TABLE 2 Characteristics of CHIC in 2010 With Persistent High Cost in 2011 or 2012 (*N* = 26 574)

Characteristics	CHIC in 2010		Likelihood of Persistent High Cost		
	No Persistent High Cost	Persistent High Cost	OR ^a	95% CI ^a	<i>P</i> ^a
Number of patients (%)	19 394 (73.0)	7 180 (27.0)	—	—	—
Age, median (IQR), y	10 [5, 14]	9 [4, 13]	1.0	[1.0–1.0]	1.0
Male gender, no. (%) of patients	10 175 (52.5)	3 773 (52.5)	1.0	[0.9–1.0]	.2
Public insurance ^b	9 902 (51.1)	4 313 (60.1)	1.2	[1.1–1.4]	<.001
Race or ethnicity					
Non-Hispanic white	10 394 (53.6)	3 751 (52.2)	Reference	—	—
Non-Hispanic black	3 449 (17.8)	1 348 (18.8)	1.0	[0.9–1.1]	.9
Hispanic	3 112 (16)	1 383 (19.3)	1.1	[0.9–1.2]	.3
Asian	455 (2.3)	134 (1.9)	0.8	[0.6–1.0]	.03
Other	1 984 (10.2)	564 (7.9)	0.8	[0.7–1.0]	.007
No. of chronic conditions					
1	4 236 (21.8)	546 (7.6)	Reference	—	—
2	4 375 (22.6)	993 (13.8)	1.7	[0.6–4.4]	.3
3+	9 706 (50)	5 595 (77.9)	2.3	[0.8–6.0]	.1
Chronic condition grouping					
None	1 026 (5.3)	42 (0.6)	Reference	—	—
Noncomplex, chronic	3 654 (18.8)	635 (8.8)	1.3	[0.5–3.6]	.6
Complex, chronic	14 714 (75.9)	6 503 (90.6)	2.0	[0.7–5.5]	.2
Complex Chronic Conditions, <i>N</i> (%) ^c					
Cardiology	4 162 (21.5)	1 295 (18.0)	0.8	[0.7–0.9]	<.001
Gastrointestinal	836 (4.3)	493 (6.9)	1.2	[1.1–1.4]	.005
Hematology/immunology	1 038 (5.4)	748 (10.4)	1.4	[1.3–1.6]	<.001
Malignancy	3 556 (18.3)	1 874 (26.1)	1.0	[0.9–1.2]	.6
Metabolic	1 545 (8.0)	1 019 (14.2)	1.0	[0.9–1.1]	.6
Neurology	4 665 (24.1)	2 516 (35.0)	1.2	[1.1–1.3]	.002
Renal	749 (3.9)	577 (8.0)	1.5	[1.4–1.7]	<.001
Respiratory	781 (4.0)	1 127 (15.7)	3.0	[2.5–3.5]	<.001
Technological assistance	5 523 (28.5)	3 544 (49.4)	1.6	[1.5–1.7]	<.001
Number of hospitalizations (2010)	1 [1–3]	3 [2–5]	1.1	[1.1–1.2]	<.001
Hospital bed days (2010)	13 [7–24]	22 [13–39]	1.0	[1.0–1.1]	<.001
No surgical admissions in 2010 ^d	5 732 [29.6]	3 489 [48.6]	2.0	[1.8–2.1]	<.001

Non-applicable denoted by em dash (—).

^a Obtained from multivariate regression by using GEEs clustered on hospital representing the association between the clinical, demographic, and utilization characteristic and the persistence of high inpatient costs as defined by cost \geq 80th percentile in 2011 or 2012.

^b Reference group is private insurance.

^c Reference group is children without the CCC

^d Reference group is children with at least one surgical admission in 2010

subsequent years. Children with a respiratory CCC (eg, cystic fibrosis, bronchopulmonary dysplasia) who experienced multiple hospitalizations in 2010 had the highest likelihood of persistent high inpatient costs in subsequent years.

Our findings complement a growing body of literature suggesting that many high-cost patients experience substantially less health care expenditures after their high-cost year. For example, a majority (ie, up to 75%) of highest-cost adult Medicare beneficiaries in 1 year may not remain highest cost in consecutive years.^{14,24,35} One recent study found that few adult “super-utilizers” experienced consistently

high levels of utilization over time.³⁶ Data from the Medical Expenditure Panel Survey reveal that approximately half of the most expensive children in 1 year will not remain in the highest-cost decile in the next year. For these children, the rates of hospitalization and the number of hospital days decreased by 75% and 84%, respectively, after their high-cost year.²³ These data, and ours, suggest transiency in hospital costs for most children after a year in which they experienced high inpatient costs.

This study also contributes to the understanding of the reasons why some children might accrue high inpatient costs. We excluded patients

prone to particularly expensive episodes of inpatient care: infants (including those using neonatal intensive care³⁷) and children dying in the hospital.³⁸ These children did not contribute to the findings. Acute illness (eg, infection, trauma) in previously healthy children probably was not a major factor either, because a small minority (4%) of children with the highest inpatient costs did not have a chronic health condition. Undergoing surgery in the high-cost year was associated with lower likelihood of persistent high costs in subsequent years, although we observed similar longitudinal trends in hospital utilization after exclusion of children

who experienced a single high-cost surgical-related hospitalization in 2010. Additional investigation is necessary to determine which surgical procedures might portend persistently high costs and whether the clinical effectiveness of select surgeries may contribute to lower inpatient costs in subsequent years.

We found that children with a respiratory CCC who experienced ≥ 3 hospitalizations in their high-cost year were the most likely to experience persistent high costs. Many respiratory CCCs may be incurable. A cluster of multiple hospitalizations experienced by these children may indicate that their condition is unstable or becoming more severe with time.³⁹ These multiple hospitalizations may also represent interventions to optimize the quality of life for children whose chronic illness is becoming more severe. A better understanding of the health and chronic illness trajectories for these children and other children with CCCs may be very helpful when estimating their future health care expenditures and developing interventions to improve their quality of care.

Although we identified combinations of clinical and demographic characteristics that were associated with persistent high inpatient costs, the highest rate of persistence in patients with these characteristics was 65%. That is, nearly 1 in 3 of these children did not experience persistent high inpatient costs in subsequent years. Clinical initiatives to contain costs may prefer to target children with a higher specificity of persistence. There are several potential explanations for why higher rates of persistence were not observed in the current study. First, we aggregated costs on an annual basis. Shorter time frames (eg, monthly) or time frames not fixed to the calendar year might allow more precise modeling; however, they may also introduce




FIGURE 4

Patient characteristics and interactions of patient characteristics associated with greatest likelihood of persistently high inpatient cost in 2011 and/or 2012. The CART model included all characteristics outlined in Table 2. Individual characteristics, and interactions between multiple characteristics, were assessed. Persistence denotes patient-level probabilities of inpatient costs ≥ 80 th percentile in 2011 and/or 2012. The gray shading denotes the terminal nodes in the decision tree analysis.

other confounders (eg, seasonality). Second, clinical assessment may be more accurate than the ICD-9 coding-based algorithms we used to identify children with different types of chronic conditions. Third, we did not assess functional status,^{40,41} disease severity,⁴² or other patient attributes that may influence trends in health care utilization because this information is not contained in PHIS. Lastly, we did not evaluate family socioeconomic status and the availability of outpatient and community health services, both of which may be associated with challenges accessing health services. Additional evaluations incorporating these elements into predictive models for future hospital expenditures should be considered.

This study has several other limitations. A gold standard definition of persistent, high health care costs does not exist. In a post hoc analysis, when the definition of high-cost patients was limited to a more or less strict percentile (eg, 90th

or 70th percentile) for persistence, the findings remained the same. We converted hospital charges to costs, which could have underestimated or overestimated costs for some hospitalizations. Generalizability may be limited to patients hospitalized in children's hospitals because the PHIS database does not contain data from non-children's hospitals. Some children could have used non-PHIS hospitals during the study period, which may have led to undercounting of hospital use. A recent study found that 13% of readmissions in children occur at a different hospital.⁴³ The likelihood of using different hospitals is significantly lower when children have a chronic condition, which most children had in our study. Health care utilization and costs for community, emergency, outpatient, and other nonhospital health services are not available in PHIS. Subsequent investigation of longitudinal spending across the care continuum in high-cost children is necessary; such investigation using comprehensive payer claim data

may help contextualize the trends in hospital care observed in the current study.

We were unable to identify children who died outside the hospital during the study period. Some children experience high hospital use at their end of life.⁴⁴ The study is not positioned to adequately assess longitudinal trends in hospital use by infants and children <2 years of age. We used ICD-9-CM codes to identify technological assistance, which may underreport the prevalence of certain technologies that do not have an ICD-9-CM code (eg, wheelchairs). We were unable to validly distinguish planned from unplanned hospitalizations, so this information was not assessed in the study. Using PHIS, we could not distinguish which children were exposed to clinical services that might affect their hospital use (eg, a care coordination or managed care program).

Despite these limitations, the findings from this study have implications for initiatives striving to improve quality of care and reduce costs. The transiency of hospital expenditures for children with highest inpatient costs may make it difficult to distinguish the effectiveness of

initiatives that target children who have already accrued high costs. Absent the use of research methods (eg, a contemporaneous control group) that account for an expected transiency in inpatient costs, some initiatives may find difficulty correlating their clinical interventions with a reduction in hospital expenditures, even when the interventions have been effective. As children's hospitals, pediatric practices, and payers continue to explore alternative payment models (eg, accountable care organizations), the unpredictability and nonpersistence of patients with high inpatient costs should be taken into account in setting cost containment goals. Future investigation is needed to identify high-cost children in advance of their cost accumulation, determine how to optimize the quality of care for these children, and understand how to intervene with targeted care management ahead of time.

CONCLUSIONS

Hospital costs are concentrated in a small number of high-cost, high-need children. Most children with high inpatient costs in 1 year may experience transient costs, with

few hospitalizations in preceding and subsequent years. Only one-fourth of these high-cost children experience persistent high children's hospital costs in subsequent years. Children with a respiratory CCC, who experienced multiple hospitalizations in their high-cost year, had the strongest correlation with persistently high inpatient costs. Clinical initiatives striving to contain hospital use in children may find this information useful.

ABBREVIATIONS

CART: classification and regression tree

CCC: complex chronic condition

CHIC: children with highest annual inpatient cost

CI: confidence interval

GEE: generalized estimating equations

ICD-9-CM: International Classification of Diseases, Ninth Revision, Clinical Modification

IQR: interquartile range

OR: odds ratio

PHIS: Pediatric Health Information System

reviewed, and revised the manuscript; Dr Matt Hall participated in the conceptualization and design of the study, coordinated acquisition of study data, carried out the statistical analysis, and drafted, critically reviewed, and revised the manuscript; Drs Neff, Mandl, Cohen, Kuo, and Agrawal participated in the conceptualization and design of the study, made a substantial contribution to the analysis and interpretation of data, and critically reviewed and revised the manuscript; Drs Brittan and David Hall made a substantial contribution to the analysis and interpretation of data and critically reviewed and revised the manuscript; Dr Berry participated in the conceptualization and design of the study, participated in the statistical analysis, and drafted, critically reviewed, and revised the manuscript; and all authors approved the final manuscript as submitted.

DOI: 10.1542/peds.2015-1829

Accepted for publication Oct 27, 2015

Address correspondence to Alon Peltz, MD, MBA, Robert Wood Johnson Foundation Clinical Scholars Program, Yale University School of Medicine, 333 Cedar St, PO Box 208088, New Haven, Connecticut, 06520. E-mail: alon.peltz@yale.edu

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2016 by the American Academy of Pediatrics

FINANCIAL DISCLOSURE: Dr Berry was supported by the Eunice Kennedy Shriver National Institute for Child Health and Human Development (K23HD058092) and the Agency for Healthcare Research and Quality (R21HS23092). These funders were not involved in the design and conduct of the study, in the collection, analysis, and interpretation of the data, and in the preparation, review, or approval of the manuscript. The other authors have indicated they have no financial relationships relevant to this article to disclose.

FUNDING: No external funding.

POTENTIAL CONFLICT OF INTEREST: The authors have indicated they have no potential conflicts of interest to disclose.

REFERENCES

- Newacheck PW, Inkelas M, Kim SE. Health services use and health care expenditures for children with disabilities. *Pediatrics*. 2004;114(1):79–85
- Berry JG AR, Cohen E, Kuo DZ. *The Landscape of Medical Care for Children With Medical Complexity*. Overland Park, KS: Children's Hospital Association; 2013
- Kuo DZ, Melguizo-Castro M, Goudie A, Nick TG, Robbins JM, Casey PH. Variation in child health care utilization by medical complexity. *Matern Child Health J*. 2014; (Apr):17
- Simon TD, Berry J, Feudtner C, et al. Children with complex chronic conditions in inpatient hospital settings in the United States. *Pediatrics*. 2010;126(4):647–655
- Cohen E, Berry JG, Camacho X, Anderson G, Wodchis W, Guttman A. Patterns and costs of health care use of children with medical complexity. *Pediatrics*. 2012;130(6). Available at: www.pediatrics.org/cgi/content/full/130/6/e1463
- Neff JM, Sharp VL, Muldoon J, Graham J, Myers K. Profile of medical charges for children by health status group and severity level in a Washington State health plan. *Health Serv Res*. 2004;39(1):73–89
- Centers for Medicare and Medicaid Innovation. Health Care Innovation Awards, round 1 and round 2. Available at: <http://innovation.cms.gov/initiatives/Health-Care-Innovation-Awards>. Accessed May 18, 2015
- Berry JG, Agrawal R, Kuo DZ, et al. Characteristics of hospitalizations for patients who use a structured clinical care program for children with medical complexity. *J Pediatr*. 2011;159(2):284–290
- Casey PH, Lyle RE, Bird TM, et al. Effect of hospital-based comprehensive care clinic on health costs for Medicaid-insured medically complex children. *Arch Pediatr Adolesc Med*. 2011;165(5):392–398
- Berman S, Rannie M, Moore L, Elias E, Dryer LJ, Jones MD Jr. Utilization and costs for children who have special health care needs and are enrolled in a hospital-based comprehensive primary care clinic. *Pediatrics*. 2005;115(6). Available at: www.pediatrics.org/cgi/content/full/115/6/e637
- Gordon JB, Colby HH, Bartelt T, Jablonski D, Krauthoefer ML, Havens P. A tertiary care–primary care partnership model for medically complex and fragile children and youth with special health care needs. *Arch Pediatr Adolesc Med*. 2007;161(10):937–944
- Liptak GS, Burns CM, Davidson PW, McAnarney ER. Effects of providing comprehensive ambulatory services to children with chronic conditions. *Arch Pediatr Adolesc Med*. 1998;152(10):1003–1008
- Mosquera RA, Avritscher EB, Samuels CL, et al. Effect of an enhanced medical home on serious illness and cost of care among high-risk children with chronic illness: a randomized clinical trial. *JAMA*. 2014;312(24):2640–2648
- Riley GF. Long-term trends in the concentration of Medicare spending. *Health Aff (Millwood)*. 2007;26(3):808–816
- Berry JG, Hall DE, Kuo DZ, et al. Hospital utilization and characteristics of patients experiencing recurrent readmissions within children's hospitals. *JAMA*. 2011;305(7):682–690
- Feudtner C, Levin JE, Srivastava R, et al. How well can hospital readmission be predicted in a cohort of hospitalized children? A retrospective, multicenter study. *Pediatrics*. 2009;123(1):286–293
- Berry JG, Hall M, Hall DE, et al. Inpatient growth and resource use in 28 children's hospitals: a longitudinal, multi-institutional study. *JAMA Pediatr*. 2013;167(2):170–177
- Burns KHC, Casey PH, Lyle RE, Bird TM, Fussell JJ, Robbins JM. Increasing prevalence of medically complex children in US hospitals. *Pediatrics*. 2010;126(4):638–646
- Berry JG, Hall M, Neff J, et al. Children with medical complexity and Medicaid: spending and cost savings. *Health Aff (Millwood)*. 2014;33(12):2199–2206
- Billings J, Mijanovich T. Improving the management of care for high-cost Medicaid patients. *Health Aff (Millwood)*. 2007;26(6):1643–1654
- Cohen E, Jovcevska V, Kuo DZ, Mahant S. Hospital-based comprehensive care programs for children with special health care needs: a systematic review. *Arch Pediatr Adolesc Med*. 2011;165(6):554–561
- Childrens Hospital Association. <http://www.childrenshospitals.org>. Accessed on December 8, 2015
- Liptak GS, Shone LP, Auinger P, Dick AW, Ryan SA, Szilagyi PG. Short-term persistence of high health care costs in a nationally representative sample of children. *Pediatrics*. 2006;118(4). Available at: www.pediatrics.org/cgi/content/full/118/4/e1001
- Joynt KE, Gawande AA, Orav EJ, Jha AK. Contribution of preventable acute care spending to total spending for high-cost Medicare patients. *JAMA*. 2013;309(24):2572–2578
- Chronic condition indicator: Healthcare Cost and Utilization Project (HCUP). Rockville, MD: Agency for Healthcare Research and Quality; 2011,
- Feudtner C, Christakis DA, Connell FA. Pediatric deaths attributable to complex chronic conditions: a population-based study of Washington State, 1980–1997. *Pediatrics*. 2000;106(1 pt 2):205–209
- Feudtner C, Villareale NL, Murray B, Sharp V, Hays RM, Neff JM. Technology-dependency among patients discharged from a children's hospital: a retrospective cohort study. *BMC Pediatr*. 2005;5(1):8
- Palfrey JS, Haynie M, Porter S, et al. Prevalence of medical technology assistance among children in Massachusetts in 1987 and 1990. *Public Health Rep*. 1994;109(2):226–233
- Palfrey JS, Walker DK, Haynie M, et al. Technology's children: report of a statewide census of children dependent on medical supports. *Pediatrics*. 1991;87(5):611–618
- 3M Health Information Systems. *All Patient Refined Diagnosis Related Groups (APR-DRGs). Version 20.0*

Methodology Overview. Wallingford, CT: 3M Health Information Systems; 2003

31. AHRQ Quality Indicators—Guide to Prevention Quality Indicators: Hospital Admission for Ambulatory Care Sensitive Conditions. Rockville, MD: Agency for Healthcare Research and Quality, 2001. AHRQ pub. no. 02-R0203
32. Breiman L, Friedman J, Stone CJ, Olshen RA. *Classification and Regression Trees*. Monterey, CA: Wadsworth & Brooks/Cole Advanced Books & Software; 1984
33. R Core Team. *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R Foundation for Statistical Computing; 2013. Available at: www.R-project.org/
34. Garber AMMT, McClellan MB. Persistence of Medicare expenditures among elderly beneficiaries. In: Garber AM, ed. *Frontiers in Health Policy Research*, vol. 1. Cambridge, MA: National Bureau of Economic Research, Massachusetts Institute of Technology; 1998:153–180
35. Congressional Budget Office. High-cost Medicare beneficiaries: a CBO paper. May 2005. Available at: <https://www.cbo.gov/sites/default/files/cbofiles/ftpdocs/63xx/doc6332/05-03-medispending.pdf>
36. Johnson TL, Rinehart DJ, Durfee J, et al. For many patients who use large amounts of health care services, the need is intense yet temporary. *Health Aff (Millwood)*. 2015;34(8):1312–1319
37. Institute of Medicine (US) Committee on Understanding Premature Birth and Assuring Healthy Outcomes; Behrman RE, Butler AS, eds. Preterm birth: causes, consequences, and prevention. Available at: www.ncbi.nlm.nih.gov/books/NBK11391/. Accessed May 18, 2015
38. Lindley LC, Lyon ME. A profile of children with complex chronic conditions at end of life among Medicaid beneficiaries: implications for health care reform. *J Palliat Med*. 2013;16(11):1388–1393
39. Feudtner C. Collaborative communication in pediatric palliative care: a foundation for problem-solving and decision-making. *Pediatr Clin North Am*. 2007;54(5):583–607, ix
40. Fleishman JA, Cohen JW. Using information on clinical conditions to predict high-cost patients. *Health Serv Res*. 2010;45(2):532–552
41. DeSalvo KB, Jones TM, Peabody J, et al. Health care expenditure prediction with a single item, self-rated health measure. *Med Care*. 2009;47(4):440–447
42. Omachi TA, Gregorich SE, Eisner MD, et al. Risk adjustment for health care financing in chronic disease: what are we missing by failing to account for disease severity? *Med Care*. 2013;51(8):740–747
43. Khan A, Nakamura MM, Zaslavsky AM et al. Same-hospital readmission rates as a measure of pediatric quality of care. *JAMA Pediatr*. 2015;169(10):905–912
44. Feudtner C, DiGiuseppe DL, Neff JM. Hospital care for children and young adults in the last year of life: a population-based study. *BMC Med*. 2003;1:3

Hospital Utilization Among Children With the Highest Annual Inpatient Cost
Alon Peltz, Matt Hall, David M. Rubin, Kenneth D. Mandl, John Neff, Mark Brittan,
Eyal Cohen, David E. Hall, Dennis Z. Kuo, Rishi Agrawal and Jay G. Berry
Pediatrics 2016;137;
DOI: 10.1542/peds.2015-1829 originally published online January 18, 2016;

Updated Information & Services	including high resolution figures, can be found at: http://pediatrics.aappublications.org/content/137/2/e20151829
References	This article cites 32 articles, 12 of which you can access for free at: http://pediatrics.aappublications.org/content/137/2/e20151829#BIBL
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Administration/Practice Management http://www.aappublications.org/cgi/collection/administration:practice_management_sub System-Based Practice http://www.aappublications.org/cgi/collection/system-based_practice_sub Advocacy http://www.aappublications.org/cgi/collection/advocacy_sub Child Health Financing http://www.aappublications.org/cgi/collection/child_health_financing_sub
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.aappublications.org/site/misc/Permissions.xhtml
Reprints	Information about ordering reprints can be found online: http://www.aappublications.org/site/misc/reprints.xhtml

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Hospital Utilization Among Children With the Highest Annual Inpatient Cost
Alon Peltz, Matt Hall, David M. Rubin, Kenneth D. Mandl, John Neff, Mark Brittan,
Eyal Cohen, David E. Hall, Dennis Z. Kuo, Rishi Agrawal and Jay G. Berry
Pediatrics 2016;137;

DOI: 10.1542/peds.2015-1829 originally published online January 18, 2016;

The online version of this article, along with updated information and services, is
located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/137/2/e20151829>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2016 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

