

Pediatric Obesity-Related Asthma: The Role of Metabolic Dysregulation

Nandini Vijayakanthi, MBBS, DNB,^a John M. Grealley, MBBSCh, PhD,^{a,b,c} Deepa Rastogi, MBBS, MS^a

The burden of obesity-related asthma among children, particularly among ethnic minorities, necessitates an improved understanding of the underlying disease mechanisms. Although obesity is an independent risk factor for asthma, not all obese children develop asthma. Several recent studies have elucidated mechanisms, including the role of diet, sedentary lifestyle, mechanical fat load, and adiposity-mediated inflammation that may underlie the obese asthma pathophysiology. Here, we review these recent studies and emerging scientific evidence that suggest metabolic dysregulation may play a role in pediatric obesity-related asthma. We also review the genetic and epigenetic factors that may underlie susceptibility to metabolic dysregulation and associated pulmonary morbidity among children. Lastly, we identify knowledge gaps that need further exploration to better define pathways that will allow development of primary preventive strategies for obesity-related asthma in children.

abstract

Pediatric obesity is a major public health issue that has reached epidemic proportions, affecting ~18% of school-going children in the United States.¹ Although the overall prevalence of pediatric obesity has increased, prevalence rates differ by age, gender, and ethnicity¹ and are partly determined by sociodemographic factors.² Notably, obesity is more prevalent among Hispanic and African American children than their non-Hispanic white counterparts.¹

Asthma, another chronic pediatric disease with increasing prevalence over the past 3 decades, affects ~10% of all school-age children in the United States.³ Racial and ethnic differences evident in the prevalence of obesity overlap with those of asthma; namely, asthma is more common among African Americans and Hispanics, particularly Puerto Ricans, compared with non-Hispanic white children.³ The increase in asthma prevalence is likely multifactorial, ranging from environmental factors including

early-life exposures to allergens to caregiver issues including low caregiver asthma management self-efficacy and empowerment.⁴ Similar environmental factors including increased built area with decreased outdoor play, increased screen time, increased intake of high-calorie foods, and caregiver food choices play a role in high childhood obesity prevalence.⁵

Over the past decade, many cross-sectional and prospective epidemiologic studies have found an association between pediatric obesity and asthma.⁶⁻¹² A recent meta-analysis, including 6 prospective cohort studies on the effect of body weight on future risk of asthma, found a twofold increased risk in obese children compared with normal-weight children,¹² suggesting that obesity is an independent risk factor for childhood asthma.¹² Clinical studies also suggest that obesity-related asthma is distinct from normal-weight asthma. Obesity-related asthma is associated



Departments of ^aPediatrics, ^bGenetics, and ^cMedicine, Albert Einstein College of Medicine, Bronx, New York

Dr Vijayakanthi performed literature searches on the topic and subtopics for relevant articles for inclusion and drafted and edited the manuscript; Dr Grealley conceptualized the manuscript with Mr Rastogi and critically reviewed the manuscript; Mr Rastogi conceptualized the manuscript with Dr Grealley, performed literature searches on the topic and subtopics for relevant articles for inclusion in the manuscript, and edited the manuscript; and all authors approved the final manuscript as submitted.

DOI: 10.1542/peds.2015-0812

Accepted for publication Nov 19, 2015

Address correspondence to, Deepa Rastogi MBBS, MS, Children's Hospital at Montefiore, Albert Einstein College of Medicine, 3415 Bainbridge Ave, Bronx, NY 10467. E-mail: drastogi@montefiore.org

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

To cite: Vijayakanthi N, Grealley JM, Rastogi D. Pediatric Obesity-Related Asthma: The Role of Metabolic Dysregulation. *Pediatrics*. 2016;137(5): e20150812

with decreased medication responsiveness¹³ and poor disease control,^{14–16} particularly among ethnic minority children,^{14,15} contributing to increased health care expenditure. However, the mechanisms that underlie these epidemiologic and clinical associations are poorly understood, and this gap precludes scientific advancement toward development of targeted therapies. Here, we review recent scientific studies that begin to elucidate the potential role of metabolic dysregulation and its associated inflammation in pediatric obesity-related asthma. It is important to note that although the majority of studies included in this review defined obesity as BMI >95th percentile, some combined current definitions of overweight and obese (BMI >85th percentile). Moreover, this review focuses on mechanisms underlying obesity-related asthma, rather than incident obesity among children with asthma.

MECHANISMS LINKING OBESITY AND ASTHMA

Many mechanisms may underlie the obesity-asthma association (Fig 1). They range from obesity-mediated alteration of pulmonary function, due to the mechanical truncal fat load and/or inflammation,¹⁷ alteration in macro- and micronutrient intake,¹⁸ and a sedentary lifestyle and its associated obesogenic behaviors.¹⁹ Furthermore, immunomodulatory effects of obesity, mediated by adipocytokines including leptin, have also been postulated to underlie asthma in obese children.^{20,21} However, because not all obese children develop asthma, these factors may play a role, but do not explain the higher predisposition for pulmonary morbidity in some, but not other, obese children. Therefore, there is need for better identification of key molecules and biomarkers that may predict asthma

development among at-risk obese children.

Recently, asthma has been associated with insulin resistance,²² dyslipidemia,^{23,24} and metabolic syndrome,²⁵ measures of metabolic dysregulation that develop in some but not all obese children.^{26,27} Moreover, genetic and epigenetic differences in molecules involved in metabolic dysregulation and its associated inflammation have been found in the context of obesity-related asthma. In this review, we initially summarize the better-investigated mechanisms such as the mechanical effect of truncal adiposity on pulmonary function and the association of sedentary lifestyle and dietary intake with obesity-related asthma. We then discuss more recent literature on the association of obesity-mediated inflammation and metabolic dysregulation with pediatric obesity-related asthma and its genetic and epigenetic footprint (Fig 1), which together begin to identify key molecules that likely underlie the complex pathophysiology of obesity-related asthma.

MECHANICAL EFFECT OF OBESITY ON PULMONARY MECHANICS AND ASTHMA

Obesity, and its related measure, truncal adiposity,^{28–30} have been associated with asthma,^{29–31} and pulmonary function deficits,²⁸ including among ethnic minority children^{30,32} (Table 1). Excess truncal adiposity renders a mechanical disadvantage to the diaphragm due to mechanical fat load and is associated with decreased functional residual capacity (FRC),^{33–35} with reduced residual volume (RV) and expiratory reserve volume (ERV).^{33–37} Lower FRC influences bronchial smooth muscle stretch, especially at the end of tidal volume exhalation, which leads to perception of increased respiratory effort with normal inspiration.³⁸ Although obese children have higher forced

expiratory volume in 1 second (FEV₁) and forced vital capacity (FVC)^{39,40} than normal-weight children, the combination of mechanical restriction and low lung volumes predispose obese children to present with lower FEV₁/FVC ratio,³³ reduced even compared with their normal-weight counterparts^{32,39,41} (Table 1). However, unlike normal-weight asthma, obese asthmatic children have reduced FRC with low lung volumes.³³

These studies suggest an association among truncal adiposity, asthma, and altered lung mechanics. However, because truncal adiposity is a risk factor for metabolic dysregulation,⁵¹ we speculate that metabolic dysregulation, not investigated in these earlier studies, may have coexisted with truncal adiposity. In keeping with this speculation, insulin resistance and dyslipidemia were found to be predictors of FEV₁/FVC ratio and ERV,³⁴ the 2 pulmonary function indices that are decreased among obese asthmatics, and mediated the association of BMI and waist circumference with these indices,³⁴ suggesting that biological factors other than mechanical fat load may mediate the influence of obesity on pulmonary function.

Ethnicity and gender may also influence these associations. Hispanics and African Americans, who bear a higher burden of obesity-related asthma, have greater truncal adiposity for the same body weight than Caucasians.⁵² With regard to gender, although obese girls have more symptoms^{53,54} and nonatopic inflammation compared with boys,⁴⁷ boys have a higher prevalence of metabolic syndrome.²⁵ Moreover, whereas one study reported an association between truncal fat and FEV₁ and FVC only among boys,⁵⁰ another study found an association between lean mass, rather than fat mass, with FEV₁, FVC, and total lung capacity (TLC) among boys,




FIGURE 1

Mechanisms proposed to underlie pediatric obesity-related asthma. This figure summarizes the factors associated with obesity-related asthma in the context of obesity preceding asthma. Although several factors such as genetics and epigenetics are also associated with childhood asthma, the relationships shown in this figure are specific to those discussed in this review.

and only with TLC in girls.⁴⁸ The disparate results of these few studies highlight the need for gender as well as ethnicity-specific investigations of the associations among mechanical fat load, presence of metabolic dysregulation, and pulmonary function deficits linked with pediatric obesity-related asthma.

Weight Loss

Thus far, there are only 2 studies on the effect of weight loss on obesity-related asthma in children.^{48,55} Jensen et al reported an improvement in asthma control in obese asthmatics

following diet-induced weight loss.⁴⁸ ERV and RV/TLC ratio and Pediatric Asthma Quality of Life Questionnaire (PAQLQ) symptom and emotional domain scores also improved but did not differ significantly from the change in the control group.⁴⁸ Van Leeuwen et al reported decreased severity of exercise-induced bronchoconstriction and improved PAQLQ scores, particularly in the symptoms and activity domain, after weight loss.⁵⁵ Moreover, limiting caloric intake in the normal range in obese children was also associated with improvement in

asthma control and PAQLQ scores.⁵⁶ Together, this limited literature suggests that weight loss in children is associated with improvement in clinical and quality-of-life parameters. However, there are no studies on pulmonary effects of weight loss in ethnic minority children. Because diet-induced weight loss in children, particularly among those of minority ethnic background, is often modest,⁵⁷ other therapeutic options are needed to address obesity-mediated pulmonary morbidity in the pediatric populations most afflicted by these diseases.

TABLE 1 Summary of Pediatric Studies Reporting an Association Between General or Truncal Adiposity and Asthma

Study	Study Design	Obesity Definition/ BMI Analysis	Sample Size, <i>n</i>	Age, y, (Range/ Mean ± SD)	Ethnicity	Finding
Tantisira et al, 2003 ³⁹	CS	BMI >95th percentile for age and gender	1041	5–12	Whites (17%), Blacks (17.6%), Hispanic (18.8%) Others (16.6%)	Decrease in FEV ₁ /FVC ratio was associated with increase in BMI.
Perez-Padilla et al, 2006 ⁴²	CS	BMI >95th percentile or BMI above limits set by International Obesity Task Force	6784	8–20	Mexican	Increase in FEV ₁ and FVC but decrease in FEV ₁ /FVC ratio with increase in BMI. Increase more in preadolescents than adolescents.
Chu et al, 2009 ⁴⁰	CS	BMI >95th percentile for age and gender	14 654	14.3 ± 0.9	Taiwanese	Higher FEV ₁ and FVC but lower FEV ₁ /FVC ratio with higher BMI.
Musaad et al, 2009 ²⁸	CS	BMI >85th percentile for age and gender	1123	5–18	Caucasian (57.4%), African American (33.8%), others (8.8%)	BMI and waist circumference were inversely associated with FEV ₁ .
Spathopoulos et al, 2009 ⁴¹	CS	BMI >95th percentile for age and gender	2715	6–11	Caucasian (Greek)	High BMI is inversely correlated with FEV ₁ , FVC, FEV ₁ /FVC ratio, and FEF _{25%–75%} .
Chen et al, 2009 ⁴³	CS	BMI analyzed as a continuous variable	718	6–17	Caucasian (Canadian)	Waist circumference was inversely associated with FEV ₁ /FVC.
Consilvio et al, 2010 ⁴⁴	CS	BMI >2 SD for age and gender	118	6–9	Caucasian (Italian)	Obese asthmatic children had low FEV ₁ /FVC ratio.
Sidoroff et al, 2011 ⁴⁵	L	BMI >98th percentile for age and gender	100	4–12	Caucasian (Finnish)	Weight gain associated with decrease in FEV ₁ /FVC ratio.
Huang et al, 2012 ⁴⁶	CS	BMI >95th percentile for age and gender	140	10–16	Mexican	No association between FEV ₁ /FVC ratio and BMI in asthmatics.
Rastogi et al, 2012 ³⁵	CS	BMI >95th percentile for age and gender	120	7–11	African Americans (50%), Hispanics (50%)	FEV ₁ /FVC ratio, and FEF _{25%–75%} was lower in obese asthmatics
Vo et al, 2013 ³²	CS	BMI >95th percentile for age and gender	980	7–20	Whites (16%), African Americans (42%), and Hispanics (42%)	Higher FEV ₁ and FVC but decrease in FEV ₁ /FVC ratio with higher BMI. FEV ₁ /FVC ratio was lower in overweight and obese African Americans and Hispanics, and obese Whites.
Jensen et al, 2013 ⁴⁷	CS	BMI z score >1.64 SD	361	8–17	Caucasian (Australian)	Lung volumes reduced among obese children; ERV reduced in obese asthmatics. RV and RV/TLC ratio reduced in obese nonasthmatic children.
Jensen et al, 2014 ⁴⁸	CS	BMI z score analyzed as a continuous variable	48	11.9 ± 2.3 (Boys) 13.6 ± 2.2 (Girls)	Caucasian (Australian)	Lean mass, not fat mass, is associated with FEV ₁ , FVC, and TLC in boys and with TLC in girls.
Sanchez-Jimenez et al, 2014 ⁴⁹	CS	BMI >95th percentile for age and gender	153	4–15	Caucasian (Spanish)	Waist circumference was inversely associated with FEV ₁ and FVC.
Wang et al, 2014 ⁵⁰	CS	BMI z score analyzed as a continuous variable	646	11–12	Caucasian (British)	Higher FEV ₁ and FVC with higher BMI in girls. Percent truncal fat inversely correlated with FEV ₁ and FVC in boys but not girls.
Rastogi et al, 2014 ³⁴	CS	BMI >95th percentile	168	13–18	African Americans (42.1%), Hispanics (57.9%)	Truncal adiposity and general adiposity were associated with reduced FRC, RV, and RV/TLC ratio.

CS, cross-sectional; L, longitudinal.

DIET AND ASTHMA

Increased intake of processed food, high in fat and low in antioxidant content, has been associated with asthma.^{58,59} Conversely, consumption of the Mediterranean diet, high in fruits and vegetables, and omega-3

fatty acids, has been found to be protective.^{60–63} Therefore, the type of fat, in addition to total fat intake, may play a role in its association with asthma. Systemic inflammation linked to dietary fat intake may underlie these associations.^{64–66}

Omega-6 fatty acids, including arachidonic acid (20:4 n-6) and linoleic acid (18:2 n-6), mediate inflammation,⁶⁷ whereas omega-3 fatty acids, including eicosapentaenoic acid (20:5 n-3) and docosahexaenoic acid (22:6 n-3),

are protective.^{67,68} Prostaglandins and leukotrienes, both of which are arachidonic acid metabolites, have been quantified in exhaled breath condensates from children with asthma.⁶⁹ However, the extent to which asthma symptomatology and pulmonary function improve with increased intake of omega-3 or decreased intake of omega-6 fatty acids is not well known among obese children with asthma and is being investigated in ongoing randomized trials of omega-3 fatty acids supplementation.⁷⁰

Furthermore, intake of micronutrients such as vitamins A,^{59,71,72} C,^{59,73} and E⁷³ has been inversely associated with asthma, whereas vitamin D insufficiency has been associated with higher asthma disease burden⁷⁴ and lower lung function.^{75,76} Although the exact mechanism through which vitamin D influences asthma in obese children is not known, vitamin D does have immunomodulatory effects⁷⁷ and may influence intestinal microflora,⁷⁸ mechanisms that have been associated with asthma pathophysiology.^{79,80} There is also evidence to suggest that maternal diet may influence incident childhood asthma and obesity, an aspect that has been previously reviewed.⁸¹ Although these initial studies suggest that dietary intake may be linked to obesity-related asthma, more research is needed to explore the various effects of dietary macro- and micronutrients on asthma.

There is also a substantive role of parental choice⁸² and feeding practices⁸³ in a child's dietary intake⁸² and behavior,⁸³ including among Hispanics^{84,85} and African Americans.⁸⁶ For example, among Hispanic households, >50% of the parents reported having sugar-sweetened beverages, and >80% reported having energy-dense foods including potato chips, cookies, cake, or ice cream in their home.⁸⁴ In keeping with these findings,

weight-resilient African American adolescents were those who consumed more fruits and vegetables and whose parents were in the healthy weight range and provided supervision to physical activity and accessed grocery stores with better food availability.⁸⁶ Given the complex relationships between macro- and micronutrient intake and asthma and the role of parental dietary choices and feeding practices on dietary intake, future studies are needed to define the impact of each of these aspects of nutritional intake on childhood asthma, including obesity-related asthma. These findings may elucidate a role of dietary modification rather than restriction in the management of obesity-related asthma,^{18,87} particularly in those of minority ethnicities, given their higher disease burden and modest effectiveness of weight loss.

SEDENTARY LIFESTYLE, PHYSICAL FITNESS, AND ASTHMA

Obese children also tend to have a sedentary lifestyle. Increased use of electronic gadgets, television watching, and video games has decreased outdoor play time and been linked with overweight and obesity in children.^{88,89} The number of hours playing video games and watching TV directly correlate with asthma incidence and prevalence among children.^{19,90} Sedentary lifestyle and decreased physical fitness cause central obesity and thereby predispose children to asthma.^{91,92} Moreover, the association of functional exercise capacity among obese asthmatics with BMI and not with FEV₁/FVC ratio, suggests a larger role of adiposity in exercise limitation among obese asthmatics.⁹³ Together these associations highlight the importance of addressing such obesogenic behaviors early in life to prevent the development of obesity and its associated pulmonary morbidities.

OBESITY-MEDIATED INFLAMMATION AND ASTHMA

Obesity is recognized to be a low-grade inflammatory state. Obesity-mediated inflammation has been associated with asthma and pulmonary function deficits.³³ Adipocyte hypoxia due to delayed neovascularization of adipose tissue is the most potent known stimulus for initiation of adipose tissue inflammation⁹⁴ and release of leptin, a proinflammatory adipokine. The proinflammatory cascade comprises a shift in the macrophage pool from the antiinflammatory M2 macrophages to the proinflammatory M1 macrophages (Fig 1).⁹⁵ Additionally, there is enhanced CD4+ T lymphocyte proliferation and differentiation into Th1 cells (Fig 1), with increased interferon- γ (IFN- γ), interleukin (IL)-6, and tumor necrosis factor (TNF) production.⁹⁵ This correlates with suppression of Th2 cells and decrease in T regulatory cells.⁹⁶ To maintain homeostasis, the proinflammatory effect of leptin is offset by antiinflammatory adipokines, including adiponectin, and omentin and the related antiinflammatory cytokine IL-10.⁹⁷⁻⁹⁹

Clinical studies have demonstrated elevated leptin²¹ and reduced adiponectin levels in obese children^{33,100} compared with nonobese children with asthma, suggesting that obesity-induced changes in the systemic adipocytokine milieu may underlie asthma in children.¹⁰¹ Serum leptin levels correlate with higher Th1/Th2 cell ratio,^{33,102} and higher serum IFN- γ levels,²¹ indicative of nonatopic inflammation among obese asthmatic children compared with their nonobese counterparts, including in ethnic minority children.^{33,103} These nonatopic systemic inflammatory patterns correlate with lower airway obstruction³³ and exercise-induced bronchoconstriction¹⁰⁴ among obese asthmatic children and persist into

adulthood.¹⁰⁵ These findings support epidemiologic reports of a lack of association between childhood obesity-related asthma and atopy^{7,101,106} and higher prevalence of noneosinophilic asthma in obese children.⁴⁷ However, there are also reports of increased atopy among obese children,^{53,54,107} including associations among BMI, atopic sensitization, and bronchial hyperresponsiveness,⁵³ as well as among BMI, atopy, cough, and wheeze,⁵⁴ particularly among girls^{53,54} (Table 2). Similar disparate links among obesity, asthma, and atopy are also observed in investigations of allergic airway inflammation using fractional exhaled nitric oxide (FeNO) and obesity-related asthma. Whereas BMI was associated with asthma only among children with low FeNO,¹⁰⁸ BMI was associated with higher asthma disease burden among those with high FeNO¹⁰⁸ (Table 2). Furthermore, FeNO was not associated with asthma among obese children,¹⁰⁹ and a significant association between BMI and FeNO was observed only among nonasthmatic children.¹¹⁰

It is hypothesized that these disparate reports either support heterogeneity in the pathophysiology of obesity-related asthma¹¹⁶ or are reflective of inherent differences in disease severity.¹⁰⁸ As noted in normal-weight asthma, although classic asthma is atopic, involving eosinophils and Th2 cells, severe asthma, even among normal-weight individuals, is nonatopic, mediated by neutrophils.¹¹⁷ Whether similar variability in the involvement of innate immune pathways comprising Th1 cells, M1 macrophages, and neutrophils occurs in the pathogenesis of obesity-related asthma needs further investigation. Recent literature highlighting the role of metabolic dysregulation in obesity-related asthma¹¹⁸ may begin to clarify these issues because differential inflammation among

obese individuals with or without metabolic dysregulation may partly underlie the heterogeneity of the obese asthma phenotype.

Obese asthmatics are also less responsive to steroid treatments.^{13,119} Peripheral blood mononuclear cells from obese asthmatics had lower production of antiinflammatory enzymes in response to dexamethasone,¹¹⁹ and increased TNF production, which directly correlated with BMI.¹¹⁹ Similar trends were also observed in bronchoalveolar lavage cells obtained from obese asthmatics.¹¹⁹ On the basis of these reports, it can be speculated that obese asthmatics may respond to nonsteroidal antiinflammatory agents including montelukast or etanercept, a TNF inhibitor,^{120,121} an aspect that needs further investigation.

OBESITY-MEDIATED METABOLIC DYSREGULATION AND ASTHMA

Association With Insulin Resistance

Obese children, particularly those of ethnic minorities, are predisposed to develop insulin resistance,¹²² a precursor to diabetes,¹²³ that is associated with systemic hyperinsulinemia.¹²² Our review of the recent literature highlights that metabolic dysregulation plays a role in pediatric obesity-related asthma (Table 3). Higher prevalence¹²⁴ and degree of insulin resistance²² and higher prevalence of its surrogate marker, acanthosis nigricans,²³ and metabolic syndrome,²⁵ have been reported among children with asthma compared with their nonasthmatic counterparts. Insulin resistance correlates with the proinflammatory markers leptin and IL-6¹²⁴ and is found to be a predictor of both lower airway obstruction and reduced lung volumes, 2 distinct measures of lung function deficits, independent of general and truncal adiposity.³⁴

Systemic inflammation, associated with insulin resistance, may be one of the mechanisms through which insulin resistance contributes to impaired lung function and asthma phenotype. In addition to its role in glucose metabolism, insulin has antiinflammatory effects.¹²⁶ Insulin supplementation has been associated with attenuation of lipopolysaccharide-induced acute lung injury in a murine model, with decreased TNF, IL-1 β , and IL-6 in the bronchoalveolar lavage fluid.¹²⁷ Obesity-mediated inhibition of insulin signaling, a key mechanism underlying insulin resistance, is associated with adipose tissue inflammation with activation of Th1 cells and innate immune pathways, involving macrophages.¹²⁸ Recent studies have found that insulin resistance mediates the association of systemic Th1 polarization with obesity-mediated pulmonary function deficits among ethnic minority children.¹⁰³ Given these initial investigations into the association among insulin resistance, inflammation, and pulmonary function impairment, further study of the underlying immunometabolic pathways is needed to determine the mechanism through which insulin resistance contributes to the obese asthma phenotype.

Another mechanism is the influence of insulin on airway smooth muscle (ASM). Insulin resistance is associated with airway hyperreactivity due to increased ASM contractility.¹²⁹ Several mechanisms underlie this observation. Hyperinsulinemia increases laminin expression in bovine ASM cells via phospho-inositide-3 kinase (PI3K)/Akt dependent pathway.¹²⁹ Insulin resistance also increases free insulin-like growth factor, which is associated with ASM proliferation.¹³⁰ Furthermore, insulin may increase airway hyperresponsiveness by modulating parasympathetic stimulation, studied in an obese

TABLE 2 Summary of Pediatric Studies Reporting an Association Between Inflammatory Mediators of Obesity and Asthma

Study	Study Design	Obesity Definition/ BMI Analysis	Sample Size, <i>n</i>	Age Range, y	Ethnicity	Finding
Huang et al, 1999 ⁵³	CS	BMI analyzed in quintiles	1459	13.2–15.5	Taiwanese	BMI was a significant predictor of atopy, allergic symptoms, and airway hyperresponsiveness in teenage girls.
Von Mutius et al, 2001 ⁷	CS	BMI analyzed in quartiles	7370	4–17	Caucasians (26.3%), African Americans (34%), Mexican Americans (35%), others (4.8%)	BMI is associated with asthma, but not with atopy, among children sampled in NHANES III.
Schachter et al, 2003 ⁵⁴	CS	BMI >95th percentile for age and gender	5993	7–12	Caucasian	Higher BMI is a risk factor for atopy, wheeze and cough in girls only but not a risk factor for asthma or airway hyperresponsiveness in either boys or girls.
Leung et al, 2004 ¹¹¹	CS	Body wt >120% of the median wt for height	115	7–18	Hong Kong	Obesity is not associated with FeNO or airway leukotriene levels in asthmatic children
Santamaria et al, 2007 ¹⁰⁹	CS	BMI >95th percentile for age and gender	50	8–16	Caucasian (Italian)	No association between FeNO and obesity among asthmatic children.
Huang et al, 2008 ¹¹²	CS	BMI >95th percentile for age and gender	89	10–16	Mexican	Higher markers of endothelial inflammation (sICAM) among obese asthmatics. No difference in CRP levels between obese and normal-weight asthmatic children.
Michelson et al, 2009 ¹¹³	CS	BMI z score analyzed as a continuous variable	10 140	0–19	Caucasians (60.6%), African Americans (14.4%), Mexicans (12.4%), other Hispanics (6.4%), others (6.3%)	BMI z score and CRP levels were associated with asthma severity among children in NHANES 2001–2004.
Visness et al, 2010 ¹⁰⁶	CS	BMI >95th percentile for age and gender	16 074	2–19	Caucasians (59.9%), African Americans (14.7%), Mexicans (12.5%), others (12.9%)	Association between obesity and asthma greater among non-atopic children than atopic children. Association of CRP with asthma among nonatopics mediated by BMI.
Rastogi et al, 2012 ⁵³	CS	BMI >95th percentile for age and gender	120	7–11	African Americans (50%), Hispanics (50%)	Obese asthmatics had systemic Th1 polarization, which directly correlated with lower airway obstruction.
Huang et al, 2012 ⁴⁶	CS	BMI >95th percentile for age and gender	178	10–16	Mexican	Obese asthmatics and nonasthmatics had higher plasminogen activator inhibitor, fibrinogen, and BMI with inversely correlated with FEV ₁ /FVC ratio.
Khan et al, 2012 ¹¹⁴	CS	BMI >95th percentile for age and gender	124	12–20	African Americans (41%), Hispanics (59%)	hsCRP highest in obese asthmatics compared with obese nonasthmatics, normal-weight asthmatics, and healthy controls
Sah PK et al, 2013 ¹¹⁵	CS	BMI >95th percentile for age and gender	269	6–17	Whites (32.7%), nonwhites (67.3%)	Obese asthmatics with poor asthma control had lower serum levels of IL-5, IL-13, and IL-10.
Youssef et al, 2013 ¹⁰²	CS	BMI >95th percentile for age and gender	70	9.3 ± 2.5 (obese asthmatics) 10.4 ± 1.3 (nonobese asthmatics) 10.7 ± 2.9 (controls)	Caucasian (Egyptian)	Obese asthmatics had high asthma severity, lower FEV ₁ . Serum leptin levels correlated with serum IFN γ levels, which directly correlated with asthma symptoms and inversely correlated with FEV ₁ among obese asthmatic children.
Jensen et al, 2013 ⁴⁷	CS	BMI z score >1.64 SD	361	8–17	Caucasian (Australian)	Noneosinophilic asthma more prevalent in obese asthmatic girls than boys.
Han et al, 2014 ¹⁰⁸	CS	BMI >95th percentile for age and gender	2681	6–17	Caucasians (33.3%), African Americans (20.2%), Hispanics (40.2%), others (5.7%)	Adiposity indicators are associated with asthma among children with low FeNO. Adiposity indicators are associated with worse asthma morbidity in those with high FeNO among children in NHANES 2007–2010.

TABLE 2 Continued

Study	Study Design	Obesity Definition/ BMI Analysis	Sample Size, <i>n</i>	Age Range, y	Ethnicity	Finding
Rastogi et al, 2015 ¹⁰³	CS	BMI >95th percentile for age and gender	168	13–18	African Americans (42.1%), Hispanics (57.9%)	Th1 polarization and monocyte activation correlated with metabolic abnormalities in obese asthmatics. Association of monocyte activation with pulmonary function was mediated by BMI, whereas that of Th1 polarization was mediated by insulin resistance.

CRP, C-reactive protein; CS, cross-sectional; HsCRP, high-sensitivity C-reactive protein.

TABLE 3 Summary of Pediatric Studies Reporting an Association Between Metabolic Dysregulation and Asthma

Study	Study Design	Obesity Definition	Sample Size, <i>n</i>	Age, y (Range/ Mean ± SD)	Ethnicity	Finding
Al-Shawwa, et al, 2006 ²⁴	CS	BMI >95th percentile for age and gender	188	4–20	Not available	Hypercholesterolemia is associated with higher asthma frequency, independent of obesity.
Al-Shawwa et al, 2007 ²²	CS	BMI >95th percentile for age and gender	415	2–18	Caucasian (40.5%), African Americans (38.5%), others (21%)	Obese asthmatics had higher levels of insulin resistance compared with morbidly obese nonasthmatics. Asthma prevalence directly correlated with insulin levels.
Arshi et al, 2010 ¹²⁴	CS	BMI analyzed as a continuous variable	31	6–17.9	Caucasian (Australian)	Insulin resistance was present among atopic asthmatics, which correlated with leptin and IL-6 levels.
Del-Rio-Navarro et al, 2010 ²⁵	CS	BMI >95th percentile for age and gender	443	12–14.2	Mexican	Prevalence of metabolic syndrome was higher among obese asthmatic boys, not girls.
Cottrell et al, 2011 ²³	CS	BMI >95th–98.9th percentile for age and gender	17 994	4–12	Whites (90.7%), African American (2.3%), others (0.9%)	Asthma is associated with dyslipidemia and insulin resistance, independent of BMI.
Lee et al, 2012 ⁵⁸	CS	BMI analyzed as a continuous variable	2082	8.5 ± 1.7	Taiwanese	Diet with high intake of fat and simple sugars was associated with increased risk of asthma.
Chen et al, 2013 ¹²⁵	CS	BMI >95th percentile for age and gender	462	10–15	Taiwanese	Asthma was associated with higher levels of total cholesterol and low-density lipoprotein, particularly in overweight and obese children.
Rastogi et al, 2014 ³⁴	CS	BMI >95th percentile for age and gender	168	13–18	African Americans (42.1%), Hispanics (57.9%)	Dyslipidemia and insulin resistance were predictors of pulmonary function deficits, independent of adiposity
Sanchez-Jimenez et al, 2014 ⁴⁹	CS	BMI >95th percentile for age and gender	153	4–15	Caucasian (Spanish)	Insulin levels were associated with allergen sensitization among asthmatics.
Rastogi et al, 2015 ¹⁰³	CS	BMI >95th percentile for age and gender	168	13–18	African Americans (42.1%), Hispanics (57.9%)	Th1 polarization and monocyte activation correlated with metabolic abnormalities. Insulin resistance mediated the association of Th1 polarization with pulmonary function.

CS, cross-sectional.

rat model.¹³¹ These studies suggest that insulin resistance and associated hyperinsulinemia influence ASM cell function through different mechanisms, with the end result of increased bronchial hyperresponsiveness. Translational

studies are now needed to study the role of each of these pathways in ASM cells obtained from obese asthmatics. We believe that better understanding of these pathways will be paradigm changing by potentially extending the use of metformin, routinely

prescribed for insulin resistance, into management of obesity-related asthma.¹³²

Association With Dyslipidemia

Similar to insulin resistance, links have been described between

dyslipidemia and wheezing in adults¹³³ and asthma among children²³ (Table 3). High-density lipoprotein (HDL) has been found to have a protective effect on pulmonary function indices in obese urban adolescents.³⁴ There is preliminary evidence to suggest that this effect may be mediated in part by the protective effect of HDL on monocyte activation.¹⁰³ There is also increasing evidence to suggest that the origins of the association of diet-induced metabolic dysregulation and pulmonary morbidity may start as early as in utero. For instance, altered fat intake, with low intake of poly-unsaturated fatty acids in the mother, has been associated with an increased predisposition to asthma among the offspring.¹³⁴ Thus, altered fat intake and dyslipidemia, irrespective of BMI, may be risk factors for airway inflammation and hyperreactive airways.²³

Many mechanisms, including inflammation, may underlie the association between dyslipidemia and asthma. High fat intake in an adult cohort was associated with increased neutrophilic airway inflammation and an attenuated response to bronchodilators.¹³⁵ Similarly, a high-fat meal, associated with elevated triglycerides and reduced HDL after 2 hours, correlated with increased levels of FeNO.¹³⁶ Because high-fat diet is associated with decreased consumption of antioxidants, it may make the lung susceptible for oxidative damage and inflammation.¹⁸ These pathways have not been studied in children, and thus the mechanisms underlying the association of dyslipidemia with asthma in children are relatively unknown.

CROSSTALK BETWEEN GENES AND ENVIRONMENT IN OBESITY-RELATED ASTHMA

The associations of asthma with obesogenic lifestyles and

obesity-mediated metabolic dysregulation, inflammation, and mechanical fat load suggest that these environmentally mediated exposures and clinical states may influence the lungs via epigenetic mechanisms.^{137,138} However, it is also evident that not all obese children with metabolic dysregulation or inflammation develop asthma, suggesting that differences in genetic susceptibility may also underlie the development of pulmonary morbidity in only some obese children. Although few studies have investigated the genetics or epigenetics of obesity-related asthma, we discuss the existing literature and the direction of association observed in these initial investigations.

Epigenetics of Obesity-Related Asthma

Epigenetic differences have been identified in context of both obesity^{139,140} and asthma^{141,142} compared with healthy controls. Among obese children, differences in DNA methylation, an epigenetic regulatory mechanism, were identified at 5 sites at the *FTO* gene, variants of which are strongly associated with obesity.¹³⁹ Similarly, hypomethylation of DNA at the *IL-4* gene promoter and hypermethylation of the *IFN γ* promoter have been observed in children with atopic asthma.¹⁴³ Genome-wide studies have also identified differential methylation of several genes associated with atopic inflammation among asthmatics.¹⁴¹ However, only 1 study defined differences in DNA methylation among children with obesity-related asthma compared with children with normal-weight asthma, obesity without asthma, and healthy controls.¹⁴⁴ In this study, specific DNA methylation patterns were associated with childhood obesity-related asthma. Gene promoters encoding for molecules involved in Th1 polarization, chemokine (C-C motif) ligand 5 (CCL5), interleukin

2 receptor α chain (IL2RA), and T-box transcription factor (TBX21), were hypomethylated, whereas those encoding for receptors for immunoglobulin E and TGF β 1, involved in Th1 cell inhibition, were hypermethylated,¹⁴⁴ suggesting DNA methylation plays a role in Th1 polarized systemic inflammation. Additionally, molecules such as PI3K and PPAR γ , involved in glucose metabolism in T cells,¹⁴⁵ and lipid uptake, respectively, were hypomethylated in obese asthmatics relative to obese nonasthmatics. These findings suggest that molecules associated with both inflammation and metabolic dysregulation are differentially methylated among obese asthmatics. Because dietary intake and nutrients modify DNA methylation,¹³⁸ these pilot results highlight the need for additional studies to investigate the effect of diet modification and related weight loss on DNA methylation and its association with insulin resistance, dyslipidemia, and systemic inflammation among obese asthmatics.

Genetics of Obesity-Related Asthma

While few conclusive studies have identified susceptibility loci for development of asthma among obese children,¹⁴⁶ a common 16p11.2 inversion that may protect against susceptibility to asthma and obesity has been identified in adults.¹⁴⁷ This inversion, found in 10% of Africans and ~50% of Europeans, is associated with increased expression of obesity-associated proteins including apolipoprotein B (APOB48R) and SH2B1, which inhibit type 1 interferon and IL27. This inversion explains ~40% of the population-attributable risk for joint susceptibility to obesity and asthma. Additional genes including the β 2-adrenergic receptor gene (*ADRB2*),^{148,149} the TNF gene,^{150,151} and the lymphotoxin- α (*LTA*) gene^{152,153} have been associated in

both obesity and asthma in children. However, the limited number of these studies highlights the paucity of data on genetic susceptibility for obesity-related asthma. Moreover, given the ethnic differences in the prevalence of pediatric obesity-related asthma, studies are needed to identify the role, if any, of ancestry-specific genetic polymorphisms that may explain the greater disease burden among Hispanics and African Americans.

RECOMMENDATIONS FOR CLINICAL PRACTICE

Together, these studies on the mechanisms underlying obesity-related asthma suggest a complex interplay among mechanical fat load of truncal adiposity, metabolic dysregulation, and inflammation. On the basis of these studies, we suggest that pediatricians consider implementing the following in their clinical practice:

1. Routine evaluation for truncal adiposity by measuring waist circumference among their patients who are overweight/obese
2. Routine evaluation for metabolic dysregulation, specifically for insulin resistance and dyslipidemia in fasting blood among obese children,¹⁵⁴ particularly in those with truncal adiposity
3. Elucidation of respiratory symptoms among obese children, particularly those with truncal adiposity, and/or metabolic dysregulation
4. Testing for pulmonary function deficits among obese children, especially those with truncal

adiposity, and/or metabolic dysregulation

5. Ensure good asthma control and encourage physical activity for weight control because there is no therapy specific for obesity-related asthma, and these children are suboptimally responsive to inhaled steroids
6. Encourage parents to monitor dietary intake, with increased intake of foods included in a Mediterranean diet and decreased consumption of processed foods

ROAD MAP FOR FUTURE

In summary, obesity-related asthma is an emerging health problem among children. Although it appears to be distinct from normal-weight asthma, further investigations are needed to better define its pathophysiology. The association of obesity-related asthma with insulin resistance and dyslipidemia provides directionality to future investigations into underlying pathways that may be amenable to pharmacologic modification. Because these metabolic abnormalities are obesity-mediated but do not develop in all obese children, quantification of these metabolic biomarkers may help identify obese children at risk for developing obesity-mediated pulmonary morbidity. Moreover, the association of asthma with diet, particularly fat and vitamin intake, and the association of diet with DNA methylation highlights the need for studies to better define the links between diet and epigenetics of obesity-related asthma, in the presence of insulin resistance and/or dyslipidemia. Given the modest effect of weight loss interventions

among children and lack of studies on the pulmonary effects of bariatric surgery, these future studies will identify mechanisms underlying the beneficial effects of nutrients and thereby facilitate the development of targeted diets for obese children at risk for developing obesity-related asthma, specifically those of Hispanic and African American ancestry. Identification of ancestry-specific genetic susceptibility will not only shed light on the reasons underlying increased disease burden among certain populations but may facilitate the development of primary prevention strategies for those identified to be genetically susceptible to obesity and its associated morbidities. Because obese asthmatics are suboptimally responsive to current asthma medications, identification of mechanisms underlying obesity-related asthma will provide direction for development of both preventative strategies and targeted therapy.

ABBREVIATIONS

ASM: airway smooth muscle
ERV: expiratory reserve volume
FeNO: fractional exhaled nitric oxide
FEV₁: forced expiratory volume in 1 second
FRC: functional residual capacity
FVC: forced vital capacity
HDL: high-density lipoprotein
IFN- γ : interferon- γ
IL: interleukin
PAQLQ: Pediatric Asthma Quality of Life Questionnaire
RV: residual volume
Th cells: T helper cells
TLC: total lung capacity
TNF: tumor necrosis factor

Copyright © 2016 by the American Academy of Pediatrics

FINANCIAL DISCLOSURE: The authors have indicated they have no financial relationships relevant to this article to disclose.

FUNDING: Dr Rastogi (K23HL118773) and Dr Grealley (R01DA030317 and R21GM101880) are supported by the National Institutes of Health. Funded by the National Institutes of Health (NIH).

POTENTIAL CONFLICT OF INTEREST: The authors have indicated they have no potential conflicts of interest to disclose.

REFERENCES

- Ogden CL, Carroll MD, Kit BK, Flegal KM. Prevalence of obesity and trends in body mass index among US children and adolescents, 1999–2010. *JAMA*. 2012;307(5):483–490
- Rossen LM, Talih M. Social determinants of disparities in weight among US children and adolescents. *Ann Epidemiol*. 2014;24(10):705–713.e2
- Akinbami LJ, Moorman JE, Garbe PL, Sondik EJ. Status of childhood asthma in the United States, 1980–2007. *Pediatrics*. 2009;123 (suppl 3):S131–145
- Canino G, Vila D, Normand SL, et al. Reducing asthma health disparities in poor Puerto Rican children: the effectiveness of a culturally tailored family intervention. *J Allergy Clin Immunol*. 2008;121(3):665–670
- Dixon B, Peña MM, Taveras EM. Lifecourse approach to racial/ethnic disparities in childhood obesity. *Adv Nutr*. 2012;3(1):73–82
- Figueroa-Muñoz JI, Chinn S, Rona RJ. Association between obesity and asthma in 4–11 year old children in the UK. *Thorax*. 2001;56(2):133–137
- von Mutius E, Schwartz J, Neas LM, Dockery D, Weiss ST. Relation of body mass index to asthma and atopy in children: the National Health and Nutrition Examination Study III. *Thorax*. 2001;56(11):835–838
- Gold DR, Damokosh AI, Dockery DW, Berkey CS. Body-mass index as a predictor of incident asthma in a prospective cohort of children. *Pediatr Pulmonol*. 2003;36(6):514–521
- Garcia-Marcos L, Arnedo Pena A, Busquets-Monge R, et al. How the presence of rhinoconjunctivitis and the severity of asthma modify the relationship between obesity and asthma in children 6–7 years old. *Clin Exp Allergy*. 2008;38(7):1174–1178
- Flaherman V, Rutherford GW. A meta-analysis of the effect of high weight on asthma. *Arch Dis Child*. 2006;91(4):334–339
- Rzehak P, Wijga AH, Keil T, et al; GA²LEN-WP 1.5 Birth Cohorts. Body mass index trajectory classes and incident asthma in childhood: results from 8 European Birth Cohorts—a Global Allergy and Asthma European Network initiative. *J Allergy Clin Immunol*. 2013;131(6):1528–1536
- Chen YC, Dong GH, Lin KC, Lee YL. Gender difference of childhood overweight and obesity in predicting the risk of incident asthma: a systematic review and meta-analysis. *Obes Rev*. 2013;14(3):222–231
- Forno E, Lescher R, Strunk R, Weiss S, Fuhlbrigge A, Celedón JC; Childhood Asthma Management Program Research Group. Decreased response to inhaled steroids in overweight and obese asthmatic children. *J Allergy Clin Immunol*. 2011;127(3):741–749
- Belamarich PF, Luder E, Kattan M, et al. Do obese inner-city children with asthma have more symptoms than nonobese children with asthma? *Pediatrics*. 2000;106(6):1436–1441
- Kattan M, Kumar R, Bloomberg GR, et al. Asthma control, adiposity, and adipokines among inner-city adolescents. *J Allergy Clin Immunol*. 2010;125(3):584–592
- Quinto KB, Zuraw BL, Poon KY, Chen W, Schatz M, Christiansen SC. The association of obesity and asthma severity and control in children. *J Allergy Clin Immunol*. 2011;128(5):964–969
- Dixon AE, Holguin F, Sood A, et al; American Thoracic Society Ad Hoc Subcommittee on Obesity and Lung Disease. An official American Thoracic Society Workshop report: obesity and asthma. *Proc Am Thorac Soc*. 2010;7(5):325–335
- Wood LG, Gibson PG. Dietary factors lead to innate immune activation in asthma. *Pharmacol Ther*. 2009;123(1):37–53
- Corbo GM, Forastiere F, De Sario M, et al; Sidria-2 Collaborative Group. Wheeze and asthma in children: associations with body mass index, sports, television viewing, and diet. *Epidemiology*. 2008;19(5):747–755
- Guler N, Kirerlerli E, Ones U, Tamay Z, Salmayenli N, Darendeliler F. Leptin: does it have any role in childhood asthma? *J Allergy Clin Immunol*. 2004;114(2):254–259
- Mai XM, Böttcher MF, Leijon I. Leptin and asthma in overweight children at 12 years of age. *Pediatr Allergy Immunol*. 2004;15(6):523–530
- Al-Shawwa BA, Al-Huniti NH, DeMattia L, Gershan W. Asthma and insulin resistance in morbidly obese children and adolescents. *J Asthma*. 2007;44(6):469–473
- Cottrell L, Neal WA, Ice C, Perez MK, Piedimonte G. Metabolic abnormalities in children with asthma. *Am J Respir Crit Care Med*. 2011;183(4):441–448
- Al-Shawwa B, Al-Huniti N, Titus G, Abu-Hasan M. Hypercholesterolemia is a potential risk factor for asthma. *J Asthma*. 2006;43(3):231–233
- Del-Rio-Navarro BE, Castro-Rodríguez JA, Garibay Nieto N, et al. Higher metabolic syndrome in obese asthmatic compared to obese nonasthmatic adolescent males. *J Asthma*. 2010;47(5):501–506
- Lee JM, Okumura MJ, Davis MM, Herman WH, Gurney JG. Prevalence and determinants of insulin resistance among U.S. adolescents: a population-based study. *Diabetes Care*. 2006;29(11):2427–2432
- Korsten-Reck U, Kromeyer-Hauschild K, Korsten K, Baumstark MW, Dickhuth HH, Berg A. Frequency of secondary dyslipidemia in obese children. *Vasc Health Risk Manag*. 2008;4(5):1089–1094
- Musaad SM, Patterson T, Ericksen M, et al. Comparison of anthropometric measures of obesity in childhood allergic asthma: central obesity is most relevant. *J Allergy Clin Immunol*. 2009;123(6):1321–7.e12
- Chen YC, Tu YK, Huang KC, Chen PC, Chu DC, Lee YL. Pathway from central obesity to childhood asthma. Physical fitness and sedentary time are leading factors. *Am J Respir Crit Care Med*. 2014;189(10):1194–1203
- Vangeepuram N, Teitelbaum SL, Galvez MP, Brenner B, Doucette J, Wolff MS. Measures of obesity associated with asthma diagnosis in ethnic minority children. *J Obes*. 2011;2011:517417

31. Papoutsakis C, Chondronikola M, Antonogeorgos G, et al. Associations between central obesity and asthma in children and adolescents: a case-control study. *J Asthma*. 2015;52(2):128–134
32. Vo P, Makker K, Matta-Arroyo E, Hall CB, Arens R, Rastogi D. The association of overweight and obesity with spirometric values in minority children referred for asthma evaluation. *J Asthma*. 2013;50(1):56–63
33. Rastogi D, Canfield SM, Andrade A, et al. Obesity-associated asthma in children: a distinct entity. *Chest*. 2012;141(4):895–905
34. Rastogi D, Bhalani K, Hall CB, Isasi CR. Association of pulmonary function with adiposity and metabolic abnormalities in urban minority adolescents. *Ann Am Thorac Soc*. 2014;11(5):744–752
35. Davidson WJ, Mackenzie-Rife KA, Witmans MB, et al. Obesity negatively impacts lung function in children and adolescents. *Pediatr Pulmonol*. 2014;49(10):1003–1010
36. Li AM, Chan D, Wong E, Yin J, Nelson EA, Fok TF. The effects of obesity on pulmonary function. *Arch Dis Child*. 2003;88(4):361–363
37. Gibson N, Johnston K, Bear N, Stick S, Logie K, Hall GL. Expiratory flow limitation and breathing strategies in overweight adolescents during submaximal exercise. *Int J Obes*. 2014;38(1):22–26
38. Salome CM, King GG, Berend N. Physiology of obesity and effects on lung function. *J Appl Physiol (1985)*. 2010;108(1):206–211
39. Tantisira KG, Litonjua AA, Weiss ST, Fuhlbrigge AL; Childhood Asthma Management Program Research Group. Association of body mass with pulmonary function in the Childhood Asthma Management Program (CAMP). *Thorax*. 2003;58(12):1036–1041
40. Chu YT, Chen WY, Wang TN, Tseng HI, Wu JR, Ko YC. Extreme BMI predicts higher asthma prevalence and is associated with lung function impairment in school-aged children. *Pediatr Pulmonol*. 2009;44(5):472–479
41. Spathopoulos D, Paraskakis E, Trypsianis G, et al. The effect of obesity on pulmonary lung function of school aged children in Greece. *Pediatr Pulmonol*. 2009;44(3):273–280
42. Pérez-Padilla R, Rojas R, Torres V, Borja-Aburto V, Olaiz G, Grp EW; The Empece Working Group. Obesity among children residing in Mexico City and its impact on lung function: a comparison with Mexican-Americans. *Arch Med Res*. 2006;37(1):165–171
43. Chen Y, Rennie D, Cormier Y, Dosman JA. Waist circumference associated with pulmonary function in children. *Pediatr Pulmonol*. 2009;44(3):216–221
44. Consilvio NP, Di Pillo S, Verini M, et al. The reciprocal influences of asthma and obesity on lung function testing, AHR, and airway inflammation in prepubertal children. *Pediatr Pulmonol*. 2010;45(11):1103–1110
45. Sidoroff V, Hyvärinen M, Piippo-Savolainen E, Korppi M. Lung function and overweight in school aged children after early childhood wheezing. *Pediatr Pulmonol*. 2011;46(5):435–441
46. Huang F, del-Río-Navarro BE, Alcántara ST, et al. Plasminogen activator inhibitor-1, fibrinogen, and lung function in adolescents with asthma and obesity. *Endocr Res*. 2012;37(3):135–144
47. Jensen ME, Gibson PG, Collins CE, Wood LG. Airway and systemic inflammation in obese children with asthma. *Eur Respir J*. 2013;42(4):1012–1019
48. Jensen ME, Gibson PG, Collins CE, Hilton JM, Wood LG. Diet-induced weight loss in obese children with asthma: a randomized controlled trial. *Clin Exp Allergy*. 2013;43(7):775–784
49. Sánchez Jiménez J, Herrero Espinet FJ, Mengibar Garrido JM, et al. Asthma and insulin resistance in obese children and adolescents. *Pediatr Allergy Immunol*. 2014;25(7):699–705
50. Wang R, Custovic A, Simpson A, Belgrave DC, Lowe LA, Murray GS. Differing associations of BMI and body fat with asthma and lung function in children. *Pediatr Pulmonol*. 2014;49(11):1049–1057
51. Mancini MC. Metabolic syndrome in children and adolescents—criteria for diagnosis. *Diabetol Metab Syndr*. 2009;1(1):20
52. Messiah SE, Arheart KL, Lipshultz SE, Miller TL. Ethnic group differences in waist circumference percentiles among U.S. children and adolescents: estimates from the 1999–2008 National Health and Nutrition Examination Surveys. *Metab Syndr Relat Disord*. 2011;9(4):297–303
53. Huang SL, Shiao G, Chou P. Association between body mass index and allergy in teenage girls in Taiwan. *Clin Exp Allergy*. 1999;29(3):323–329
54. Schachter LM, Peat JK, Salome CM. Asthma and atopy in overweight children. *Thorax*. 2003;58(12):1031–1035
55. van Leeuwen JC, Hoogstrate M, Duiverman EJ, Thio BJ. Effects of dietary induced weight loss on exercise-induced bronchoconstriction in overweight and obese children. *Pediatr Pulmonol*. 2014;49(12):1155–1161
56. Luna-Pech JA, Torres-Mendoza BM, Luna-Pech JA, Garcia-Cobas CY, Navarrete-Navarro S, Elizalde-Lozano AM. Normocaloric diet improves asthma-related quality of life in obese pubertal adolescents. *Int Arch Allergy Immunol*. 2014;163(4):252–258
57. Khan UI, Rieder J, Cohen HW, Coupey SM, Wildman RP. Effect of modest changes in BMI on cardiovascular disease risk markers in severely obese, minority adolescents. *Obes Res Clin Pract*. 2010;4(3):e163–e246
58. Lee SC, Yang YH, Chuang SY, Liu SC, Yang HC, Pan WH. Risk of asthma associated with energy-dense but nutrient-poor dietary pattern in Taiwanese children. *Asia Pac J Clin Nutr*. 2012;21(1):73–81
59. Huang SL, Pan WH. Dietary fats and asthma in teenagers: analyses of the first Nutrition and Health Survey in Taiwan (NAHSIT). *Clin Exp Allergy*. 2001;31(12):1875–1880
60. Garcia-Marcos L, Canflanca IM, Garrido JB, et al. Relationship of asthma and rhinoconjunctivitis with obesity, exercise and Mediterranean diet in Spanish schoolchildren. *Thorax*. 2007;62(6):503–508
61. de Batlle J, Garcia-Aymerich J, Barraza-Villarreal A, Antó JM, Romieu I. Mediterranean diet is associated with reduced asthma and

- rinitis in Mexican children. *Allergy*. 2008;63(10):1310–1316
62. Nurmatov U, Devereux G, Sheikh A. Nutrients and foods for the primary prevention of asthma and allergy: systematic review and meta-analysis. *J Allergy Clin Immunol*. 2011;127(3):724–33.e1, 30
 63. Garcia-Marcos L, Castro-Rodríguez JA, Weinmayr G, Panagiotakos DB, Priftis KN, Nagel G. Influence of Mediterranean diet on asthma in children: a systematic review and meta-analysis. *Pediatr Allergy Immunol*. 2013;24(4):330–338
 64. Dandona P, Ghanim H, Chaudhuri A, Dhindsa S, Kim SS. Macronutrient intake induces oxidative and inflammatory stress: potential relevance to atherosclerosis and insulin resistance. *Exp Mol Med*. 2010;42(4):245–253
 65. Aljada A, Mohanty P, Ghanim H, et al. Increase in intranuclear nuclear factor kappaB and decrease in inhibitor kappaB in mononuclear cells after a mixed meal: evidence for a proinflammatory effect. *Am J Clin Nutr*. 2004;79(4):682–690
 66. Black PN, Sharpe S. Dietary fat and asthma: is there a connection? *Eur Respir J*. 1997;10(1):6–12
 67. Wendell SG, Baffi C, Holguin F. Fatty acids, inflammation, and asthma. *J Allergy Clin Immunol*. 2014;133(5):1255–1264
 68. Horrobin DF. Low prevalences of coronary heart disease (CHD), psoriasis, asthma and rheumatoid arthritis in Eskimos: are they caused by high dietary intake of eicosapentaenoic acid (EPA), a genetic variation of essential fatty acid (EFA) metabolism or a combination of both? *Med Hypotheses*. 1987;22(4):421–428
 69. Glowacka E, Jedynak-Wasowicz U, Sanak M, Lis G. Exhaled eicosanoid profiles in children with atopic asthma and healthy controls. *Pediatr Pulmonol*. 2013;48(4):324–335
 70. Lang JE, Mougey EB, Allayee H, et al; Nemours Network for Asthma Research. Nutri-genetic response to omega-3 fatty acids in obese asthmatics (N00A): rationale and methods. *Contemp Clin Trials*. 2013;34(2):326–335
 71. Arora P, Kumar V, Batra S. Vitamin A status in children with asthma. *Pediatr Allergy Immunol*. 2002;13(3):223–226
 72. Mizuno Y, Furusho T, Yoshida A, Nakamura H, Matsuura T, Eto Y. Serum vitamin A concentrations in asthmatic children in Japan. *Pediatr Int*. 2006;48(3):261–264
 73. Nakamura K, Wada K, Sahashi Y, et al. Associations of intake of antioxidant vitamins and fatty acids with asthma in pre-school children. *Public Health Nutr*. 2013;16(11):2040–2045
 74. Brehm JM, Celedón JC, Soto-Quiros ME, et al. Serum vitamin D levels and markers of severity of childhood asthma in Costa Rica. *Am J Respir Crit Care Med*. 2009;179(9):765–771
 75. Somashekar AR, Prithvi AB, Gowda MN. Vitamin D levels in children with bronchial asthma. *J Clin Diagn Res*. 2014;8(10):PC04–PC07
 76. Yao TC, Tu YL, Chang SW, et al; Prediction of Allergies in Taiwanese Children (PATCH) Study Group. Serum 25-hydroxyvitamin D levels in relation to lung function and exhaled nitric oxide in children. *J Pediatr*. 2014;165(6):1098–1103.e1
 77. Mann EH, Chambers ES, Pfeffer PE, Hawrylowicz CM. Immunoregulatory mechanisms of vitamin D relevant to respiratory health and asthma. *Ann N Y Acad Sci*. 2014;1317:57–69
 78. Ooi JH, Li Y, Rogers CJ, Cantorna MT. Vitamin D regulates the gut microbiome and protects mice from dextran sodium sulfate-induced colitis. *J Nutr*. 2013;143(10):1679–1686
 79. Verhulst SL, Vael C, Beunckens C, Nelen V, Goossens H, Desager K. A longitudinal analysis on the association between antibiotic use, intestinal microflora, and wheezing during the first year of life. *J Asthma*. 2008;45(9):828–832
 80. Cho SH, Stanciu LA, Holgate ST, Johnston SL. Increased interleukin-4, interleukin-5, and interferon- γ in airway CD4+ and CD8+ T cells in atopic asthma. *Am J Respir Crit Care Med*. 2005;171(3):224–230
 81. Litonjua AA, Gold DR. Asthma and obesity: common early-life influences in the inception of disease. *J Allergy Clin Immunol*. 2008;121(5):1075–1084; quiz 1085–1076
 82. Jones LR, Steer CD, Rogers IS, Emmett PM. Influences on child fruit and vegetable intake: sociodemographic, parental and child factors in a longitudinal cohort study. *Public Health Nutr*. 2010;13(7):1122–1130
 83. Carper JL, Orlet Fisher J, Birch LL. Young girls' emerging dietary restraint and disinhibition are related to parental control in child feeding. *Appetite*. 2000;35(2):121–129
 84. Santiago-Torres M, Adams AK, Carrel AL, LaRowe TL, Schoeller DA. Home food availability, parental dietary intake, and familial eating habits influence the diet quality of urban Hispanic children. *Child Obes*. 2014;10(5):408–415
 85. Arredondo EM, Elder JP, Ayala GX, Campbell N, Baquero B, Duerksen S. Is parenting style related to children's healthy eating and physical activity in Latino families? *Health Educ Res*. 2006;21(6):862–871
 86. Brogan K, Idalski Carcone A, Jen KL, Ellis D, Marshall S, Naar-King S. Factors associated with weight resilience in obesogenic environments in female African-American adolescents. *J Acad Nutr Diet*. 2012;112(5):718–724
 87. Scott HA, Gibson PG, Garg ML, et al. Dietary restriction and exercise improve airway inflammation and clinical outcomes in overweight and obese asthma: a randomized trial. *Clin Exp Allergy*. 2013;43(1):36–49
 88. Cox R, Skouteris H, Rutherford L, Fuller-Tyszkiewicz M, Dell'Aquila D, Hardy LL. Television viewing, television content, food intake, physical activity and body mass index: a cross-sectional study of preschool children aged 2–6 years. *Health Promot J Austr*. 2012;23(1):58–62
 89. Ghavamzadeh S, Khalkhali HR, Alizadeh M. TV viewing, independent of physical activity and obesogenic foods, increases overweight and obesity in adolescents. *J Health Popul Nutr*. 2013;31(3):334–342
 90. Sherriff A, Maitra A, Ness AR, et al. Association of duration of television viewing in early childhood with the

- subsequent development of asthma. *Thorax*. 2009;64(4):321–325
91. Rasmussen F, Lambrechtsen J, Siersted HC, Hansen HS, Hansen NC. Low physical fitness in childhood is associated with the development of asthma in young adulthood: the Odense schoolchild study. *Eur Respir J*. 2000;16(5):866–870
 92. Vlaski E, Stavric K, Seckova L, Kimovska M, Isjanovska R. Influence of physical activity and television-watching time on asthma and allergic rhinitis among young adolescents: preventive or aggravating? *Allergol Immunopathol (Madr)*. 2008;36(5):247–253
 93. Rastogi D, Khan UI, Isasi CR, Coupey SM. Associations of obesity and asthma with functional exercise capacity in urban minority adolescents. *Pediatr Pulmonol*. 2012;47(11):1061–1069
 94. O'Rourke RW, White AE, Metcalf MD, et al. Hypoxia-induced inflammatory cytokine secretion in human adipose tissue stromovascular cells. *Diabetologia*. 2011;54(6):1480–1490
 95. Ferrante AW Jr. The immune cells in adipose tissue. *Diabetes Obes Metab*. 2013;15(suppl 3):34–38
 96. Luczyński W, Wawrusiewicz-Kurylonek N, Ilendo E, et al. Generation of functional T-regulatory cells in children with metabolic syndrome. *Arch Immunol Ther Exp (Warsz)*. 2012;60(6):487–495
 97. Manigrasso MR, Ferroni P, Santilli F, et al. Association between circulating adiponectin and interleukin-10 levels in android obesity: effects of weight loss. *J Clin Endocrinol Metab*. 2005;90(10):5876–5879
 98. Catli G, Anik A, Abaci A, Kume T, Bober E. Low omentin-1 levels are related with clinical and metabolic parameters in obese children. *Exp Clin Endocrinol Diabetes*. 2013;121(10):595–600
 99. Hong EG, Ko HJ, Cho YR, et al. Interleukin-10 prevents diet-induced insulin resistance by attenuating macrophage and cytokine response in skeletal muscle. *Diabetes*. 2009;58(11):2525–2535
 100. Yuksel H, Sogut A, Yilmaz O, Onur E, Dinc G. Role of adipokines and hormones of obesity in childhood asthma. *Allergy Asthma Immunol Res*. 2012;4(2):98–103
 101. Nagel G, Koenig W, Rapp K, Wabitsch M, Zoellner I, Weiland SK. Associations of adipokines with asthma, rhinoconjunctivitis, and eczema in German schoolchildren. *Pediatr Allergy Immunol*. 2009;20(1):81–88
 102. Youssef DM, Elbehidy RM, Shokry DM, Elbehidy EM. The influence of leptin on Th1/Th2 balance in obese children with asthma. *J Bras Pneumol*. 2013;39(5):562–568
 103. Rastogi D, Fraser S, Oh J, et al. Inflammation, metabolic dysregulation, and pulmonary function among obese urban adolescents with asthma. *Am J Respir Crit Care Med*. 2015;191(2):149–160
 104. Baek HS, Kim YD, Shin JH, Kim JH, Oh JW, Lee HB. Serum leptin and adiponectin levels correlate with exercise-induced bronchoconstriction in children with asthma. *Ann Allergy Asthma Immunol*. 2011;107(1):14–21
 105. Dixon AE, Johnson SE, Griffes LV, et al. Relationship of adipokines with immune response and lung function in obese asthmatic and non-asthmatic women. *J Asthma*. 2011;48(8):811–817
 106. Visness CM, London SJ, Daniels JL, et al. Association of childhood obesity with atopic and nonatopic asthma: results from the National Health and Nutrition Examination Survey 1999–2006. *J Asthma*. 2010;47(7):822–829
 107. Visness CM, London SJ, Daniels JL, et al. Association of obesity with IgE levels and allergy symptoms in children and adolescents: results from the National Health and Nutrition Examination Survey 2005–2006. *J Allergy Clin Immunol*. 2009;123(5):1163–1169, 1169.e1–1169.e4
 108. Han Y-Y, Forno E, Celedón JC. Adiposity, fractional exhaled nitric oxide, and asthma in U.S. children. *Am J Respir Crit Care Med*. 2014;190(1):32–39
 109. Santamaria F, Montella S, De Stefano S, et al. Asthma, atopy, and airway inflammation in obese children. *J Allergy Clin Immunol*. 2007;120(4):965–967
 110. Erkoçoğlu M, Kaya A, Ozcan C, et al. The effect of obesity on the level of fractional exhaled nitric oxide in children with asthma. *Int Arch Allergy Immunol*. 2013;162(2):156–162
 111. Leung TF, Li CY, Lam CWK, et al. The relation between obesity and asthmatic airway inflammation. *Pediatr Allergy Immunol*. 2004;15(4):344–350
 112. Huang F, del-Río-Navarro BE, Monge JJ, et al. Endothelial activation and systemic inflammation in obese asthmatic children. *Allergy Asthma Proc*. 2008;29(5):453–460
 113. Michelson PH, Williams LW, Benjamin DK, Barnato AE. Obesity, inflammation, and asthma severity in childhood: data from the National Health and Nutrition Examination Survey 2001–2004. *Ann Allergy Asthma Immunol*. 2009;103(5):381–385
 114. Khan UI, Rastogi D, Isasi CR, Coupey SM. Independent and synergistic associations of asthma and obesity with systemic inflammation in adolescents. *J Asthma*. 2012;49(10):1044–1050
 115. Sah PK, Gerald Teague W, Demuth KA, Whitlock DR, Brown SD, Fitzpatrick AM. Poor asthma control in obese children may be overestimated because of enhanced perception of dyspnea. *J Allergy Clin Immunol Pract*. 2013;1(1):39–45
 116. Peters MC, Fahy JV. Type 2 immune responses in obese individuals with asthma. *Am J Respir Crit Care Med*. 2013;188(6):633–634
 117. Moore WC, Hastie AT, Li X, et al; National Heart, Lung, and Blood Institute's Severe Asthma Research Program. Sputum neutrophil counts are associated with more severe asthma phenotypes using cluster analysis. *J Allergy Clin Immunol*. 2014;133(6):1557–63.e5
 118. Periyalil HA, Gibson PG, Wood LG. Immunometabolism in obese asthmatics: are we there yet? *Nutrients*. 2013;5(9):3506–3530
 119. Sutherland ER, Goleva E, Strand M, Beuther DA, Leung DY. Body mass and glucocorticoid response in asthma. *Am J Respir Crit Care Med*. 2008;178(7):682–687
 120. Price D, Musgrave SD, Shepstone L, et al. Leukotriene antagonists as first-line

- or add-on asthma-controller therapy. *N Engl J Med.* 2011;364(18):1695–1707
121. Berry MA, Hargadon B, Shelley M, et al. Evidence of a role of tumor necrosis factor alpha in refractory asthma. *N Engl J Med.* 2006;354(7):697–708
 122. Levy-Marchal C, Arslanian S, Cutfield W, et al; ESPE-LWPES-ISPAD-APPES-APEG-SLEP-JSPE; Insulin Resistance in Children Consensus Conference Group. Insulin resistance in children: consensus, perspective, and future directions. *J Clin Endocrinol Metab.* 2010;95(12):5189–5198
 123. Goran MI, Ball GD, Cruz ML. Obesity and risk of type 2 diabetes and cardiovascular disease in children and adolescents. *J Clin Endocrinol Metab.* 2003;88(4):1417–1427
 124. Arshi M, Cardinal J, Hill RJ, Davies PSW, Wainwright C. Asthma and insulin resistance in children. *Respirology.* 2010;15(5):779–784
 125. Chen YC, Tung KY, Tsai CH, et al. Lipid profiles in children with and without asthma: interaction of asthma and obesity on hyperlipidemia. *Diabetes Metab Syndr.* 2013;7(1):20–25
 126. Hyun E, Ramachandran R, Hollenberg MD, Vergnolle N. Mechanisms behind the anti-inflammatory actions of insulin. *Crit Rev Immunol.* 2011;31(4):307–340
 127. Liu ML, Dong HY, Zhang B, et al. Insulin reduces LPS-induced lethality and lung injury in rats. *Pulm Pharmacol Ther.* 2012;25(6):472–477
 128. Lumeng CN, Saltiel AR. Inflammatory links between obesity and metabolic disease. *J Clin Invest.* 2011;121(6):2111–2117
 129. Dekkers BG, Schaafsma D, Tran T, Zaagsma J, Meurs H. Insulin-induced laminin expression promotes a hypercontractile airway smooth muscle phenotype. *Am J Respir Cell Mol Biol.* 2009;41(4):494–504
 130. Noveral JP, Bhala A, Hintz RL, Grunstein MM, Cohen P. Insulin-like growth factor axis in airway smooth muscle cells. *Am J Physiol.* 1994;267(6 pt 1):L761–L765
 131. Nie Z, Jacoby DB, Fryer AD. Hyperinsulinemia potentiates airway responsiveness to parasympathetic nerve stimulation in obese rats. *Am J Respir Cell Mol Biol.* 2014;51(2):251–261
 132. Bossé Y. Endocrine regulation of airway contractility is overlooked. *J Endocrinol.* 2014;222(2):R61–R73
 133. Fenger RV, Gonzalez-Quintela A, Linneberg A, et al. The relationship of serum triglycerides, serum HDL, and obesity to the risk of wheezing in 85,555 adults. *Respir Med.* 2013;107(6):816–824
 134. Lumia M, Luukkainen P, Tapanainen H, et al. Dietary fatty acid composition during pregnancy and the risk of asthma in the offspring. *Pediatr Allergy Immunol.* 2011;22(8):827–835
 135. Wood LG, Garg ML, Gibson PG. A high-fat challenge increases airway inflammation and impairs bronchodilator recovery in asthma. *J Allergy Clin Immunol.* 2011;127(5):1133–1140
 136. Rosenkranz SK, Townsend DK, Steffens SE, Harms CA. Effects of a high-fat meal on pulmonary function in healthy subjects. *Eur J Appl Physiol.* 2010;109(3):499–506
 137. von Mutius E. Gene-environment interactions in asthma. *J Allergy Clin Immunol.* 2009;123(1):3–11, quiz 12–13
 138. Szarc vel Szic K, Ndlovu MN, Haegeman G, Vanden Berghe W. Nature or nurture: let food be your epigenetic medicine in chronic inflammatory disorders. *Biochem Pharmacol.* 2010;80(12):1816–1832
 139. Almén MS, Jacobsson JA, Moschonis G, et al. Genome wide analysis reveals association of a FTO gene variant with epigenetic changes. *Genomics.* 2012;99(3):132–137
 140. Martínez JA, Cordero P, Campián J, Milagro FI. Interplay of early-life nutritional programming on obesity, inflammation and epigenetic outcomes. *Proc Nutr Soc.* 2012;71(2):276–283
 141. Stefanowicz D, Hackett TL, Garmaroudi FS, et al. DNA methylation profiles of airway epithelial cells and PBMCs from healthy, atopic and asthmatic children. *PLoS One.* 2012;7(9):e44213
 142. Kim YJ, Park SW, Kim TH, et al. Genome-wide methylation profiling of the bronchial mucosa of asthmatics: relationship to atopy. *BMC Med Genet.* 2013;14(39):39
 143. Kwon NH, Kim JS, Lee JY, Oh MJ, Choi DC. DNA methylation and the expression of IL-4 and IFN- γ promoter genes in patients with bronchial asthma. *J Clin Immunol.* 2008;28(2):139–146
 144. Rastogi D, Suzuki M, Grealley JM. Differential epigenome-wide DNA methylation patterns in childhood obesity-associated asthma. *Sci Rep.* 2013;3(2164):2164
 145. MacIver NJ, Michalek RD, Rathmell JC. Metabolic regulation of T lymphocytes. *Annu Rev Immunol.* 2013;31:259–283
 146. Melén E, Himes BE, Brehm JM, et al. Analyses of shared genetic factors between asthma and obesity in children. *J Allergy Clin Immunol.* 2010;126(3):631–7.e1, 8
 147. González JR, Cáceres A, Esko T, et al. A common 16p11.2 inversion underlies the joint susceptibility to asthma and obesity. *Am J Hum Genet.* 2014;94(3):361–372
 148. Oguri K, Tachi T, Matsuoka T. Visceral fat accumulation and metabolic syndrome in children: the impact of Trp64Arg polymorphism of the beta3-adrenergic receptor gene. *Acta Paediatr.* 2013;102(6):613–619
 149. Silverman EK, Kwiatkowski DJ, Sylvia JS, et al. Family-based association analysis of beta2-adrenergic receptor polymorphisms in the childhood asthma management program. *J Allergy Clin Immunol.* 2003;112(5):870–876
 150. Aoki T, Hirota T, Tamari M, et al. An association between asthma and TNF-308G/A polymorphism: meta-analysis. *J Hum Genet.* 2006;51(8):677–685
 151. Sookoian SC, González C, Pirola CJ. Meta-analysis on the G-308A tumor necrosis factor alpha gene variant and phenotypes associated with the metabolic syndrome. *Obes Res.* 2005;13(12):2122–2131
 152. Hamid YH, Urhammer SA, Glümer C, et al. The common T60N polymorphism of the lymphotoxin-alpha gene is associated with type 2 diabetes and other phenotypes of the metabolic syndrome. *Diabetologia.* 2005;48(3):445–451

153. Wang TN, Chen WY, Wang TH, Chen CJ, Huang LY, Ko YC. Gene-gene synergistic effect on atopic asthma: tumour necrosis factor-alpha-308 and lymphotoxin-alpha-Ncol in Taiwan's children. *Clin Exp Allergy*. 2004;34(2):184–188
154. Expert Panel on Integrated Guidelines for Cardiovascular Health and Risk Reduction in Children and Adolescents. Expert Panel on Integrated Guidelines for Cardiovascular Health and Risk Reduction in Children and Adolescents: summary report. *Pediatrics*. 2011; 128(suppl 5):S213–256

Pediatric Obesity-Related Asthma: The Role of Metabolic Dysregulation

Nandini Vijayakanthi, John M. Grealley and Deepa Rastogi

Pediatrics 2016;137;

DOI: 10.1542/peds.2015-0812 originally published online April 8, 2016;

Updated Information & Services

including high resolution figures, can be found at:
<http://pediatrics.aappublications.org/content/137/5/e20150812>

References

This article cites 151 articles, 18 of which you can access for free at:
<http://pediatrics.aappublications.org/content/137/5/e20150812#BIBL>

Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):

Obesity

http://www.aappublications.org/cgi/collection/obesity_new_sub

Allergy/Immunology

http://www.aappublications.org/cgi/collection/allergy:immunology_sub

Asthma

http://www.aappublications.org/cgi/collection/asthma_sub

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:

<http://www.aappublications.org/site/misc/Permissions.xhtml>

Reprints

Information about ordering reprints can be found online:
<http://www.aappublications.org/site/misc/reprints.xhtml>

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Pediatric Obesity-Related Asthma: The Role of Metabolic Dysregulation

Nandini Vijayakanthi, John M. Grealley and Deepa Rastogi

Pediatrics 2016;137;

DOI: 10.1542/peds.2015-0812 originally published online April 8, 2016;

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/137/5/e20150812>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2016 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

