

Diabetes in Pregnancy and Childhood Cognitive Development: A Systematic Review

Akilew Awoke Adane, MPH, Gita D. Mishra, PhD, Leigh R. Tooth, PhD

abstract

CONTEXT: The effect of diabetes during pregnancy on the cognitive development of offspring is unclear because of inconsistent findings from limited studies.

OBJECTIVE: This review was aimed to provide the best available scientific evidence on the associations between maternal pregnancy diabetes and the cognitive development of offspring.

DATA SOURCES: A search was conducted in the Embase, CINAHL, PubMed, PsycINFO, and Scopus databases.

STUDY SELECTION: Studies addressing the cognitive development of offspring (aged ≤ 12 years) as outcome and any diabetes in pregnancy as an exposure were included.

DATA EXTRACTION: Data were extracted and evaluated for quality by 2 independent reviewers.

RESULTS: Fourteen articles were eligible for the review. Ten studies investigated the associations between maternal pregestational diabetes or both pregestational and gestational diabetes and offspring's cognitive development; 6 studies found at least 1 negative association. Four studies exclusively examined the relationships between gestational diabetes and offspring's cognitive development; 2 studies found a negative association, 1 a positive association, and 1 a null association. The use of diverse cognitive and diabetes assessment tools/criteria, as well as statistical power, contributed to the inconsistent findings.

LIMITATIONS: The English-language restriction and publication bias in the included studies are potential limitations.

CONCLUSIONS: Although there are few data available regarding the associations between maternal pregnancy diabetes and offspring's cognitive development, this review found that maternal diabetes during pregnancy seems to be negatively associated with offspring's cognitive development. Large prospective studies that address potential confounders are needed to confirm the independent effect of maternal diabetes during pregnancy.

Centre for Longitudinal and Life Course Research, School of Public Health, the University of Queensland, Australia

Mr Adane designed the study, performed the systematic review, and drafted the manuscript; Prof Mishra and A/Prof Tooth contributed to the design of the study, interpretation of the results, and critical revision of the manuscript for important intellectual content; and all authors approved the final manuscript as submitted and agree to be accountable for all aspects of the work.

DOI: 10.1542/peds.2015-4234

Accepted for publication Feb 19, 2016

To cite: Adane AA, Mishra GD, Tooth LR. Diabetes in Pregnancy and Childhood Cognitive Development: A Systematic Review. *Pediatrics*. 2016;137(5):e20154234

Globally, the prevalence of both pregestational diabetes mellitus (PDM) and gestational diabetes mellitus (GDM) are increasing significantly due to the rising rates of obesity, aging, and population growth.¹⁻⁴ In women, the global prevalence of age-adjusted diabetes increased from 7.5% in 1980 to 9.2% in 2008.⁴ An increasing future trend in the prevalence of diabetes is also projected across all countries worldwide. For instance, in 2011, there were 366 million people with diabetes, and over the next 19 years, this figure is expected to rise to 552 million. Substantial increases are expected in developing countries related to lifestyle changes after rapid urbanization, globalization, and aging.⁵

Any type of diabetes during pregnancy is associated with poor maternal and perinatal outcomes. For instance, congenital malformation,⁶ preterm birth,^{7,8} large for gestational age at birth/macrosomia,^{7,9-11} and stillbirth^{6,10,11} are significantly more common in offspring of diabetic mothers than nondiabetic mothers. Even offspring of mothers with borderline GDM have been found to be born too early and to be macrosomic.¹²

Due to advances in medical care, offspring of mothers with diabetes are increasingly surviving the perinatal period. Consequently, the long-term effects of diabetes during pregnancy on later childhood health such as obesity/adiposity, mainly via large for gestational age at birth, has been clearly established.¹³⁻¹⁵

In contrast, data regarding the long-term effects of maternal diabetes during pregnancy on the offspring's cognitive development are relatively limited and, when data are available, the findings are inconclusive.¹⁶⁻²³

The mechanisms by which maternal diabetes during pregnancy is associated with the offspring's cognitive development are unclear. Excess glucose levels in patients with diabetes, a condition known as hyperglycemia, is hypothesized to cause cognitive impairment.²⁴ Animal

models have shown that maternal diabetes during pregnancy is usually associated with hyperglycemia and that more glucose will pass to the fetus, and this action may hinder the cognitive development of the offspring.²⁵ Although not always found,²⁶ metabolic complications related to diabetes during pregnancy such as acetonuria,^{27,28} ketoacidosis,²⁰ and glycosuria¹⁶ are also associated with offspring's cognitive impairment. It has been suggested by some investigators that offspring born to either mothers with PDM or GDM are more likely to have poorer cognitive and language development than those born to nondiabetic mothers.²⁸⁻³³ However, other investigators have either found no such associations^{17,18,20,22} or report contradictory findings.²³

To the best of our knowledge, systematically synthesized information on the associations between maternal diabetes during pregnancy and offspring's cognitive development, particularly in childhood, is lacking. The implications of such information are twofold: (1) it would provide decision-makers/clinicians with comprehensive information to deliver preconception counseling; and (2) it would enable the early identification of infants at future risk of cognitive development. It would also avail information for researchers to identify research gaps and guide future research development. Thus, the goal of the present systematic review was to provide the best available scientific evidence regarding the possible associations between maternal diabetes in pregnancy and the cognitive development of offspring.

METHODS

Data Sources and Search Strategies

The PubMed database was searched by using combinations of key words: (((((offspring) OR

child*)) AND (((((((((((("cognitive development") OR "cognitive functions") OR "school performance") OR "educational outcomes") OR "language development") OR "neuropsychological tests") OR "mental development") OR neurodevelopment) OR "child development") OR cognition) OR intelligence) OR "IQ") OR "intelligence tests")) AND (((mother*) OR pregnan*) OR women)) AND (((("type I diabetes") OR "type II diabetes") OR "gestational diabetes") OR "diabetes insipidus") OR "diabetes mellitus") OR diabetes). Similarly, systematic searches were conducted in the Embase, CINAHL, PsycINFO, and Scopus databases using the same key word combinations tailored to each database until June 2015. These searches were limited to studies in the English language with human domain restrictions. Moreover, the reference list of all identified relevant records were searched for additional studies. The authors were not contacted for additional studies or data.

Study Selection

Studies addressing the cognitive and/or language development of offspring (aged ≤ 12 years) as outcome and either PDM or GDM or both as a main exposure or confounder were included in this systematic review. Reviews, commentaries, and case or descriptive studies lacking relevant control groups were excluded.

Records deemed relevant from the title screening were further evaluated by using the information available in the abstract by 2 independent reviewers (A.A.A. and L.R.T.) using the eligibility criterion. The same authors further evaluated the full records of those eligible articles from the abstract review. Disagreements were resolved by face-to-face discussions and through joint review of the records. The third author helped resolve any disagreement between the first 2 reviewers (Fig 1).

Data Extraction

Data on the first author, publication year, country, study design, sample size, offspring's age at follow-up, diabetes type and assessment, cognitive type and measurement, confounders accounted/adjusted for, and main findings between maternal pregnancy diabetes and offspring's cognitive development were extracted by the reviewers by using a standardized data extraction format (Table 1).

Quality Assessment

The Newcastle-Ottawa quality assessment scale for case-control and cohort studies³⁴ was used to assess the quality of included studies. Two modifications were made: a "nonblind standard assessment" option was included in the outcome subsection of the cohort studies assessment scale and "the length of follow-up for the outcome to occur" criteria was omitted because the goal of this systematic review was to examine the effect of any diabetes during pregnancy on cognitive development of offspring to the age of 12 years. Consequently, a slightly modified score ranging from 0 (most likely biased) to 8 (highly unlikely to be biased) was calculated for each study.

Data Synthesis and Analysis

Meta-analysis was not performed because of heterogeneous cognitive outcomes and having too few studies that adjusted for potential confounders. Instead, a narrative review and qualitative summarizations were undertaken (Table 2).

RESULTS

A total of 1295 records were identified with English-language and human domain restrictions. After removal of duplicates, the titles of




FIGURE 1

Flow diagram of studies included in the systematic review of maternal diabetes during pregnancy and childhood cognitive development.

847 articles were screened; 679 were identified as not relevant or as further duplicates. The abstracts of 168 articles were then evaluated independently, and 111 records were excluded. Subsequent full-record evaluations of 68 articles resulted in a total of 19 eligible studies. However, 2 studies^{35,36} were further excluded because they were published with the same populations used in 2 included studies.^{18,20} In addition, 5 relatively different studies^{22,32,37-39} were conducted by using the same cohort study with varied sample sizes. All reported global cognitive development as a secondary outcome or confounder at the age of 1 year^{32,37-39} and 4 years.²² None of these studies primarily aimed to evaluate the global cognitive development of offspring of mothers with diabetes mellitus (ODM). Therefore, we included Nelson et al³² (ie, better sample size and relatively recent) and Townsend et al²² (ie, cognitive development was measured at the age of 4 years by using the Wechsler Intelligence Scale

for Children) in the review. Overall, 14 studies^{16-23,28-33} were eligible for the present review.

Quality Assessment

Using the adapted Newcastle-Ottawa quality assessment scale, the scores for each study ranged from 2 to 6 (from a total of 8). In addition to low overall scores, the following quality concerns were observed: one-half of the studies reported unadjusted results; in all studies (except 1 report³⁰) the population was obtained from unrepresentative populations, mainly from hospital facilities; and 7 of 14 studies^{16,17,22,23,29,31,33} reported the use of standard cognitive development assessment tools but failed to provide or report blind assessment.

Study Characteristics

The 14 studies were conducted between 1969 and 2015. The offspring's age at cognitive assessment ranged from 6 months

TABLE 1 Characteristics of Studies Reporting the Association Between Maternal Diabetes During Pregnancy and Cognitive Development of Offspring

First Author/ Year	Country	Study Design	Sample Size	Age at Follow-up	Diabetes Type	Diabetes Diagnosis	Cognitive Assessment	Cognitive Outcome	Main Findings on Maternal Pregnancy Diabetes and Offspring's Cognitive Development (95% CI)	Effect Size; Cohen's <i>d</i> (95% CI)
Bonilla et al, 2012 ²⁹	UK	Cohort study	6272 mother-child pairs	8 y	PDM + GDM	Record-based	WISC-III	IQ	DM associated with lower IQ (MD), -3.5 (-5.6, -1.5); <i>P</i> = .001	No sufficient data
Churchill et al, 1969 ³⁸	US	Retrospective cohort study	237 ODM and control subjects	8 mo and 4 y	PDM + GDM	Somogyi-Nelson glucose tolerance curves	BSID at 8 mo and SBIS at 4 y	IQ (<i>n</i> = 121) and MDI (<i>n</i> = 231)	DM associated with lower IQ (96 vs 103; <i>P</i> = .001) and MDI (78 vs 81; <i>P</i> = .001) DM without acetone has no association with MDI (81.0 vs 81.0), PDI (33.4 vs 34.2), and IQ (101.3 vs 100.8); all, <i>P</i> > .05	No sufficient data
Dionne et al, 2008 ³⁰	Canada	Retrospective cohort study	221 OGDM and 2612 control subjects	1.5-7 y	GDM	OGTT	MCDI at 18 and 30 mo, PPVT at 48 mo, expressive and receptive vocabulary at 60 mo, and EDI teacher-assessed communication at 72 and 84 mo	Language, NVIQ, and RM	QNTS GDM associated with lower (mean); expressive vocabulary at 18 mo (-0.25 ± 0.96 vs 0.02 ± 0.99) and at 30 mo (-0.33 ± 1.22 vs 0.04 ± 0.96), expressive grammar at 30 mo (-0.28 ± 1.18 vs 0.04 ± 0.96) and oral communication at 72/84 mo (-0.30 ± 1.21 vs 0.05 ± 0.96); all, <i>P</i> < .05	Expressive vocabulary Effect size varies from 0.27 to 0.41
			QNTS (<i>n</i> ranges from 721 to 861) and QLSCD (<i>n</i> varies from 955 to 1728)						No association with (mean) receptive vocabulary (-0.06 ± 1.00 vs 0.00 ± 0.99) and receptive grammar (-0.10 ± 1.16 vs 0.02 ± 0.98) at 18 mo, receptive grammar at 30 mo (0.04 ± 0.97 vs 0.01 ± 1.01), expressive vocabulary (-0.12 ± 0.90 vs 0.04 ± 0.99) and receptive vocabulary at 60 mo (-0.09 ± 0.86 vs 0.01 ± 1.01), math (-0.10 ± 1.02 vs 0.05 ± 0.93) and reading (-0.04 ± 0.96 vs 0.03 ± 0.99) scores at 72/84 mo; all, <i>P</i> > .05	Effect size varies from -0.37 to -0.30
									QLSCD GDM associated with lower (mean); expressive vocabulary (-0.28 ± 1.09 vs 0.02 ± 0.99) and grammar at 30 mo (-0.38 ± 1.21 vs 0.03 ± 0.97) but not with receptive vocabulary at 30 mo (-0.18 ± 1.10 vs 0.01 ± 0.98) and at 42 mo (-0.16 ± 0.81 vs 0.02 ± 1.00)	Expressive vocabulary Effect size varies from -0.34 to -0.17 Receptive vocabulary

TABLE 1 Continued

First Author/ Year	Country	Study Design	Sample Size	Age at Follow-up	Diabetes Type	Diabetes Diagnosis	Cognitive Assessment	Cognitive Outcome	Main Findings on Maternal Pregnancy Diabetes and Offspring's Cognitive Development (95% CI)	Effect Size; Cohen's <i>d</i> (95% CI)
Fraser et al, 2012 ¹⁶	UK	Cohort study	SEA	4 and 8 y	PDM + GDM	Record based	SEA at 4 y and WISC-III at 8 y	Educational attainment and IQ	GDM associated with language impairment (OR), 2.2 (1.4,3.5), RM (F = 5.31; P = .02), and NVIQ (F = 3.73; P = .05) but not with short-term memory (F = 2.86; P = .09)	Effect size varies from -0.18 to -0.06 Receptive grammar Effect size varies from -0.11 to 0.03 Oral communication Effect size (<i>d</i>) = -0.32 Math Effect size (<i>d</i>) = -0.15 Reading Effect size (<i>d</i>) = -0.07 For the others, no sufficient data PDM
Hod et al, 1999 ⁵¹	Israel	Cohort study	24 OGDM, 21 OPDM, and 5804 control subjects IQ	1 y	PDM	OGTT	BSID-II	MDI	PDM has no association with (MD) SEA, 0.04 (-1.68,1.75); FIQ, -0.54 (-9.61,8.52); VIQ, 1.5 (-7.63,10.74); and PIQ, -4.01 (-13.80,5.78)	SEA, -0.13 (-0.51,0.26)
Nelson et al, 2003 ³²	US	Cohort study	23 OGDM, 20 OPDM, and 5079 control subjects	1 y	PDM + GDM	OGTT	BSID-II	MDI	GDM associated with (MD) lower VIQ, -9.92 (-18.34, -1.50); not associated with FIQ, -5.93 (-14.24, 2.38); PIQ, -0.19 (-9.17, 8.78); or SEA, 0.30 (-1.12, 1.72)	FIQ, -0.13 (-0.51,0.26) VIQ, -0.01 (-0.40,0.37) PIQ, -0.27 (-0.66,0.11) GDM SEA, -0.03 (-0.37,0.31) FIQ, -0.40 (-0.75, -0.06) VIQ, -0.43 (-0.78, -0.09) PIQ, -0.19 (-0.53,0.15) MDI, -0.64 (-1.12,-0.17)
Nomura et al, 2012 ¹⁷	US	Cohort study	21 OGDM and 191 control subjects	3-4 y	GDM	Self-reported	WPPSI-III, developmental; neuropsychologic assessment	IQ, language	PDM associated with lower MDI (91 ± 9.0 vs 98 ± 12.1; P < .05) DM associated with lower MDI (100 ± 9 vs 104 ± 8; P < .03)	MDI, -0.49 (-0.94,-0.03)
									GDM associated with lower FIQ (109.2 ± 1.4 vs 113.6 ± 3.5), VIQ (110.5 ± 1.5 vs 113.6 ± 4.2), language (108.8 ± 1.4 vs 112.9 ± 3.9), and memory (96.6 ± 1.4 vs 101.1 ± 3.8); all, P < .05; No association with PIQ (109.8 ± 1.5 vs 111.9 ± 3.8); P = .14	FIQ, -1.30 (-2.01,-0.60) VIQ, -0.76 (-1.46,-0.07) Language, -1.09 (-1.79, -0.39) Memory, -1.23 (-1.93, -0.52) PIQ, -0.57 (-1.26,0.12)

TABLE 1 Continued

First Author/ Year	Country	Study Design	Sample Size	Age at Follow-up	Diabetes Type	Diabetes Diagnosis	Cognitive Assessment	Cognitive Outcome	Main Findings on Maternal Pregnancy Diabetes and Offspring's Cognitive Development (95% CI)	Effect Size; Cohen's <i>d</i> (95% CI)
Ornoy et al, 1999 ¹⁹	Israel	Retrospective cohort study	32 OGDM and 57 control subjects	5–12 y	GDM	Blood glucose concentration test	WISC-Revised	IQ	Young age (5–8 y) GDM associated with lower FIQ (111 ± 14 vs 121 ± 8) and VIQ (107 ± 11 vs 115 ± 11); both, <i>P</i> < .05; Not associated with PIQ (114 ± 17 vs 123 ± 11); <i>P</i> > .05 Older age (9–12 y) No association with FIQ (115 ± 13 vs 116 ± 12), VIQ (109 ± 12 vs 113 ± 13), or PIQ (119 ± 15 vs 117 ± 12); all, <i>P</i> > .05	Young group IQ, -0.95 (-1.40, -0.49)
Ornoy et al, 2001 ¹⁸	Israel	Retrospective cohort study	57 OPDM, 32 OGDM, and 57 control subjects	5–12 y	PDM + GDM	Blood glucose concentration test	WISC-Revised	IQ	PDM has no association with FIQ (117.7 ± 12 vs 118.5 ± 11), VIQ (112.4 ± 12 vs 114.4 ± 12), and PIQ (120.4 ± 19 vs 119.7 ± 11.5) GDM has no association with FIQ (113.5 ± 14.3 vs 118.5 ± 11), VIQ (108.0 ± 11.5 vs 114.4 ± 12), and PIQ (116.0 ± 16.0 vs 119.7 ± 11.5); all, <i>P</i> > .05	IQ, -0.08 (-0.44, 0.30) VIQ, -0.17 (-0.53, 0.20) PIQ, 0.05 (-0.32, 0.41) GDM IQ, -0.41 (-0.84, 0.03) VIQ, -0.54 (-0.98, -0.10) PIQ, -0.28 (-0.71, 0.16) PDM
Rizzo et al, 1991 ²⁰	USA	Cohort study	80 OPDM, 82 OGDM, and 29 control subjects	2–5 y	PDM + GDM	OGTT- O'Sullivan and Mahan criteria	BSID at 2 y SBIS at 3–5 y	MDI, IQ	PDM has no association with MDI (89 ± 18 vs 89 ± 13) and IQ (89 ± 14 vs 92 ± 10) GDM has no association with MDI (90 ± 14 vs 89 ± 13) and IQ (93 ± 13 vs 92 ± 10)	MDI, 0 (-0.43, 0.43) IQ, -0.23 (-0.67, 0.20) GDM MDI, 0.07 (-0.35, 0.50) IQ, 0.08 (-0.36, 0.52)

TABLE 1 Continued

First Author/ Year	Country	Study Design	Sample Size	Age at Follow-up	Diabetes Type	Diabetes Diagnosis	Cognitive Assessment	Cognitive Outcome	Main Findings on Maternal Pregnancy Diabetes and Offspring's Cognitive Development (95% CI)	Effect Size; Cohen's <i>d</i> (95% CI)
Sells et al, 1994 ²¹	US	Cohort study	109 OPDM (70 early entry and 39 late entry), and 90 control subjects	6, 12, 24, and 36 mo	PDM	Not stated	BSID at 6, 12, and 24 mo; SBIS; spontaneous language sample; and PPVT at 36 mo revised Vineland Adaptive	MDI, IQ, and language	PDM associated with lower PPVT scores (114 ± 11.0 vs 111 ± 10.1 vs 105 ± 11.4) and Vineland Communication subscale (111 ± 10.2 vs 110 ± 9.2 vs 103 ± 11.5)	PPVT Early entry, -0.28 (-0.65, 0.09) Later entry, -0.81 (-1.30, -0.32)
							Behavior scales at 12, 24, and 36 mo		PDM has no association with mean length of utterance (4.1 ± 1.0 vs 4.0 ± 0.86 vs 3.7 ± 0.91) MDI at 6 mo (107 ± 12.1 vs 104 ± 14.4 vs 106 ± 13.7), at 12 mo (117 ± 12.5 vs 113 ± 15.3 vs 112 ± 3.5) and at 24 mo (118 ± 18.4 vs 118 ± 19.4 vs 112 ± 16.8)	Communication Early entry, -0.10 (-0.48, 0.27) Later entry, -0.76 (-1.25, -0.27)
									SBIS Verbal reasoning (114 ± 12.4 vs 110 ± 10.2 vs 105 ± 14.2), abstract reasoning (104 ± 11.8 vs 104 ± 8.7 vs 103 ± 9.7), quantitative reasoning (105 ± 9.5 vs 104 ± 9.1 vs 96.5 ± 8.7), short-term memory (110 ± 12.1 vs 110 ± 10.5 vs 108 ± 9.6) and composite score (110 ± 9.6 vs 109 ± 7.9 vs 103 ± 11.0). All comparisons are control versus early entry versus late entry	Length of Utterance Early entry, -0.11 (-0.48, 0.27) Later entry, -0.41 (-0.89, 0.07) MDI Early entry, ranges -0.29 to 0 Later entry, ranges -0.33 to -0.08 SBIS Early entry, ranges -0.35 to 0 Later entry, ranges -0.70 to -0.09 FIQ, -0.16 (-0.88,0.56) VIQ, 0.06 (-0.66,0.77) PIQ, -0.31 (-1.03,0.41)
Townsend et al, 2005 ²²	US	Cohort study	15 ODM and 15 control subjects	4 y	PDM + GDM	Not stated	WPPSI-Revised at 4 y	IQ	DM has no association with FIQ (118 ± 16 vs 121 ± 21), VIQ (116 ± 16 vs 115 ± 19), or PIQ (116 ± 14 vs 121 ± 18)	

TABLE 1 Continued

First Author/ Year	Country	Study Design	Sample Size	Age at Follow-up	Diabetes Type	Diabetes Diagnosis	Cognitive Assessment	Cognitive Outcome	Main Findings on Maternal Pregnancy Diabetes and Offspring's Cognitive Development (95% CI)	Effect Size; Cohen's <i>d</i> (95% CI)
Veena et al, 2010 ²³	India	Cohort study	32 OGDM and 483 control subjects	9–10 y	GDM	OGTT	KABC-II (learning, long-term retrieval/ storage) Word order Pattern reasoning	Learning long-term retrieval/ storage Short-term memory Reasoning ability	GDM associated with higher mean cognitive scores of long-term retrieval/ storage (β), 0.38 (0.01, 0.75), verbal ability—names, 0.46 (0.09, 0.83) GDM has no association with: attention and concentration, 0.32 (–0.04, 0.67); visuospatial ability, 0.01 (–0.36, 0.39); verbal ability—animals, 0.22 (–0.16, 0.61); reasoning ability, 0.14 (–0.23, 0.51); and short-term memory, 0.03 (–0.35, 0.40)	Long-term retrieval/ storage, 0.49 (0.13, 0.85) Verbal ability— names, 0.45 (0.09, 0.81) Attention and concentration, 0.54 (0.19, 0.90) Verbal ability—animals, 0.34 (–0.02, 0.69) Short-term memory, 0.23 (–0.13, 0.59) All effect sizes are unadjusted
Yamashita et al, 1996 ³³	Japan	Cohort study	33 OPDM, 3 OGDM, and 34 control subjects	3–4 y	PDM + GDM	OGTT	Tanaka-Binet intelligence scale	IQ Attention and concentration	DM associated with lower IQ (98 ± 17 vs 113 ± 15)	IQ, –0.93 (–1.43, –0.44)

BSID, Bayley Scales of Infant Development; CI, confidence interval; DM, diabetes mellitus; EDI, Early Development Instrument; FIQ, Full IQ; KABC, Kaufman Assessment Battery for Children; MCDI, MacArthur Communicative Development Inventory; MD, mean difference; NVIQ, nonverbal IQ; OGTT, oral glucose tolerance test; OR, odds ratio; PIQ, performance IQ; PPVT, Peabody Picture Vocabulary Test; QLSOD, Quebec Longitudinal Study of Child Development; QNTS, Quebec Newborn Twin Study; RM, recognition memory; SBIS, Stanford-Binet Intelligence Scale; SEA, school entry assessment; VIQ, verbal IQ; WISC, Wechsler Intelligence Scale for Children; WPPSI-III, Wechsler Preschool and Primary Scale of Intelligence, Third Edition.

to 12 years. Most of the studies were conducted in the United States, Israel, and the United Kingdom; 3 were conducted in India, Japan, and Canada. The majority of the studies ($n = 10$) were prospective cohorts. Although the measurement tools were diverse, offspring's IQ according to several different IQ scales was the most commonly assessed cognitive outcome, followed by the mental development index (MDI) of the Bayley Scales of Infant Development. Overall, 10 of 14 studies^{16,18,20–22,28,29,31–33} investigated the associations between maternal PDM or both PDM and GDM with cognitive development of offspring. The remaining 4 studies^{17,19,23,30} exclusively examined the associations between maternal GDM and offspring's cognitive development (Table 1).

Association Between PDM and Offspring's Cognitive Development

In 1969, Churchill et al²⁸ reported the negative associations between maternal diabetes in pregnancy (ie, both PDM and GDM) and offspring's cognitive development. They administered the MDI at age 8 months ($n = 121$) and the Stanford-Binet Intelligence Scale at age 4 years ($n = 231$). In this study, maternal diabetes during pregnancy with acetonuria was associated with lower MDI and IQ scores. ODM, complicated by acetonuria, scored ~ 2.5 and 7 points lower in MDI and IQ scores, respectively, compared with offspring of mothers without diabetes. However, in the absence of acetonuria, ODM had MDI and IQ scores similar to control subjects.

In 1991, Rizzo et al²⁰ conducted a cohort study of 89 offspring of mothers with pregestational diabetes mellitus (OPDM), 99 offspring of mothers with gestational diabetes mellitus

(OGDM), and 35 control subjects aged 2 to 5 years. The study examined the correlation between maternal metabolism during pregnancy and offspring's cognitive and behavioral functioning. Accordingly, the MDI scores of the offspring correlated inversely with the mother's third-trimester plasma β -hydroxybutyrate levels, and the Binet's IQ scores correlated inversely with the mother's third-trimester plasma β -hydroxybutyrate and free fatty acid levels. However, the investigators found similar MDI and Binet's IQ scores among groups. In support of this outcome, Ornoy et al,¹⁸ using 32 OGDM, 57 OPDM, and 57 control subjects matched on age, socioeconomic status (SES), gestational age, birth order, and family size, reported similar Wechsler's IQ scores across these groups. Despite lack of statistical significance, OGDM scored 5 points below control subjects.

Nelson et al³² aimed to evaluate cross-modal recognition memory of infants, and they provided a simple MDI score comparison (unadjusted) between 52 ODM and 75 control subjects at the age of 1 year. They found a significant difference between these groups (mean \pm SD, 100 ± 9 vs 104 ± 8 ; $P < .03$). However, using a small subset (15 ODM and 15 control subjects) of the same population but at the age of 4 years, Townsend et al²² supported the null association between maternal diabetes during pregnancy and offspring's cognitive development as measured by using Wechsler's IQ scale. The investigators failed to provide explanations for this discrepancy except for the cognitive measurement variations.

In another study, Sells et al²¹ examined the neurodevelopmental outcomes of offspring born to mothers with insulin-dependent PDM. The study included 109 OPDM (70 early entry and 39 late entry) and 90 control subjects, and the

investigators administered various cognitive and language development measures at 6 to 36 months of age. In this study, PDM was significantly associated with lower scores of language development in late-entry groups, although this association was not observed between maternal PDM and offspring's MDI and the Binet IQ scores.

Hod et al³¹ confirmed the negative association in 31 OPDM and 41 control subjects at 1 year of age between maternal PDM and offspring MDI; OPDM scored 7 points below control subjects (91 ± 9 vs 98 ± 12 ; $P < .05$). Similarly, Yamashita et al³³ found significantly lower Tanaka-Binet IQ scores in 15 OPDM (98 ± 17) compared with 15 control subjects (113 ± 15 ; $P < .0001$). However, both studies failed to account for any potential confounder.

Bonilla et al²⁹ ($n = 6272$) and Fraser et al¹⁶ examined the cognitive development of offspring born to women with PDM and GDM at the age of 4 and 8 years in 2 different large cohort studies. Bonilla et al reported that maternal diabetes during pregnancy was negatively associated with Wechsler's IQ scores; ODM scored 3.5 (95% confidence interval, -5.6 to -1.5 ; $P = .001$) points lower compared with offspring of women without diabetes. The investigators did not report the outcomes according to diabetes type, failed to account for potential confounders such as SES and prepregnancy BMI, and did not provide information about the nature and size of the control group. Fraser et al also reported that both PDM and GDM were associated with lower offspring school entry assessment (age 4 years, ~ 5849 children) and Wechsler's IQ (age 8 years, ~ 5124 children) scores. However, after full adjustments were made for various confounders, the negative associations persisted only between maternal GDM and verbal IQ (mean

TABLE 2 Summary of the Associations Between Diabetes in Pregnancy and Offspring's Cognitive Development, and Confounders and Covariates

Study/Year	Association	Potential Confounders/Covariates												
		Maternal					Offspring							
		Age	BMI	SES	Education	HDP	Depression	Lifestyle Behaviors	Parity	Duration of Breastfeeding	Birth Weight	GA	Age	Gender
Bonilla et al, 2012 ²⁹	↓	✓								✓	✓	✓	✓	✓
Churchill et al, 1969 ²⁸	↓	✓		✓						✓	✓	✓	✓	✓
Dionne et al, 2008 ³⁰	↓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓	✓
Fraser et al, 2012 ¹⁶	↓	✓						✓		✓	✓	✓	✓	✓
Hod et al, 1999 ³¹	↓	✓						✓		✓	✓	✓	✓	✓
Nelson et al, 2003 ³²	↓	✓						✓		✓	✓	✓	✓	✓
Nomura et al, 2012 ¹⁷	↔	✓						✓		✓	✓	✓	✓	✓
Ornoy et al, 1999 ^{18a}	↔			✓				✓		✓	✓	✓	✓	✓
Ornoy et al, 2001 ¹⁸	↔			✓				✓		✓	✓	✓	✓	✓
Rizzo et al, 1991 ²⁰	↔			✓				✓		✓	✓	✓	✓	✓
Sells et al, 1994 ²¹	↓				✓									
Townsend et al, 2005 ²²	↔													
Veena et al, 2010 ²³	↑	✓	✓	✓	✓						✓	✓	✓	✓
Yamashita et al, 1996 ³³	↓										✓	✓	✓	✓

↓, significant negative association; ↑, significant positive association; ↔, no significant association; GA, gestational age; HDP, hypertensive disorder of pregnancy.

^a Found significant association between GDM and IQ scores in young age (5–8 years) groups only.

difference, 9.92) scores. Despite full model adjustment, Fraser et al acknowledged the presence of the small number of mothers with diabetes (~44) and a significant loss to follow-up (ie, IQ data were available for ~49% of the cohort).

Association Between GDM and Offspring's Cognitive Development

Although the associations between maternal GDM and offspring's cognitive development was separately reported in 7 studies,^{16–20,23,30} only 4 studies^{17,19,23,30} exclusively addressed GDM. Hence, to avoid repetition, only the last 4 studies are presented here.

In a retrospective cohort study, Ornoy et al¹⁹ compared the neuropsychological function of 32 school-aged children born to mothers with well-controlled GDM and 57 control subjects matched according to age, birth order, and parental SES. Although the study was underpowered and lacked any adjustment for confounders, the younger aged (ie, 5–8 years) OGDM scored 8 points (107 ± 11 vs 115 ± 11) and 10 points (111 ± 14 vs 121 ± 8) lower in Wechsler's verbal IQ and full-scale IQ scores, respectively, than control subjects; such associations were not observed in older children (aged 9–12 years) or on the performance IQ scale.

Dionne et al³⁰ compared OGDM and control subjects (aged 1.5–7 years) in various language development measures. In this study, OGDM scored between 0.27 and 0.41 SD below control subjects on expressive vocabulary and grammar at 18 and 30 months. At 72 and 84 months, maternal GDM was also associated with lower mean scores in oral communication. In both cases, the model was adjusted for gender, gestational age, birth weight, Apgar score, gestational

hypertension, alcohol use, and smoking in pregnancy. The same authors also reported an odds ratio of 2.2 (95% confidence interval, 1.4 to 3.5) for the risk of language impairment. At 42 and 60 months of age, no differences between OGDM and control subjects on expressive and receptive vocabulary were observed. In addition, the groups did not differ in mean reading and math scores.

In contrary, Veena et al²³ reported that maternal GDM was associated with higher mean cognitive scores of long-term retrieval/storage, verbal ability, attention, and concentration; these scores were measured by using the Kaufman Assessment Battery for Children, the Wechsler Intelligence Scale for Children, and other cognitive tests in 32 school-aged OGDM and 483 control children after adjustments were made for various confounders. The investigators acknowledged the relatively small number of OGDM and requested larger studies.

More recently, Nomura et al¹⁷ examined the independent and synergistic effect of maternal GDM and low SES on the cognitive and language development of 21 OGDM and 191 control subjects. They compared preschool-aged children (aged 3–4 years) exposed to 1 of 4 conditions: neither mother's GDM nor low SES ($n = 97$); mother's GDM but not low SES ($n = 12$); no mother's GDM but low SES ($n = 94$); and both mother's GDM and low SES ($n = 9$). From these pairwise comparisons, they found that children born to both diabetic and low SES mothers had lower verbal and full-scale IQ and language composite scores on the Wechsler Preschool and Primary Scale of Intelligence. Compared with the first group, maternal GDM alone was also significantly associated with lower cognitive (full and verbal IQ)

and language development but with attenuated effect sizes.

DISCUSSION

To our knowledge, this article is the first systematic review to examine the associations between maternal diabetes during pregnancy and childhood cognitive development. We found a few geographically limited studies, the majority of which were small and did not adjust for important confounders. Although not conclusive, 8 of the 14 studies seemed to support the negative association between maternal diabetes during pregnancy and offspring's cognitive and language development. Generally, the effect sizes were heterogeneous, varying from -1.30 to 0.54 . Effect sizes were consistently larger for language development compared with performance IQ, suggesting that the latter is less likely to be affected by maternal diabetes during pregnancy. The finding has relevant clinical implications because the prevalence of GDM and type 2 diabetes mellitus are increasing.^{3,40} It is worth noting, however, that the evidence is from limited observational studies (ie, whether the observed association was exclusively due to diabetes during pregnancy, its complications, or confounders was unclear with the available evidence).

Most of the studies^{16,29,30} conducted with relatively larger sample sizes, totaling ≥ 2833 offspring, produced consistent negative associations. Others that failed to detect such associations were conducted with smaller sample sizes^{18,20,22} and therefore had less power to detect true differences between groups.

Only 2 of the 14 eligible studies^{16,23} fully accounted for potential confounders such as maternal prepregnancy BMI, SES, maternal age, alcohol use, and smoking during pregnancy and offspring-related

covariates. Maternal prepregnancy overweight/obesity is a well-known strong predictor of both maternal PDM (particularly type 2 diabetes mellitus) and GDM.⁴¹ For instance, in the United States, $>80\%$ and 49% of patients with diabetes were found to be overweight and obese, respectively.⁴² Together with the increasing trend of overweight and obesity and the subsequent effect of these conditions on maternal diabetes,⁵ the relationship between these factors and later cognitive development of the offspring must be better understood. To date, there have been only a few studies to show the independent associations of maternal prepregnancy obesity and offspring's cognitive development.^{43–45} Hence, studies in the present review that perceived a negative association between maternal diabetes during pregnancy and offspring's cognitive development could have been confounded by maternal prepregnancy obesity.

Studies in young offspring consistently found that both maternal PDM and GDM reduced their cognitive and language development, whereas the majority of the studies in older children showed no effect of maternal diabetes on their cognitive development. This outcome suggests that either the intrauterine effect of diabetes may diminish as children get older or postnatal factors such as SES³⁰ accounted for the association. Alternatively, the effect of diabetes in pregnancy may be reversible as the cognitive abilities in young children are prone to changes, mainly to their home environment.⁴⁶

Maternal SES, measured according to maternal education, occupation, and income, is a powerful determinant of health. In a recent systematic review,⁴⁷ SES was a significant confounder in the association between preterm birth and cognitive deficit, and these investigators recommended the

need to adjust the role of SES in studies reporting child cognitive development. Similarly, Nomura et al,¹⁷ included in this systematic review, revealed a negative synergistic effect of maternal low SES and GDM on offspring's IQ and language development. Moreover, a large cohort study⁴⁸ using data from Swedish population registers found lower IQ scores in nonsibling men born to diabetic mothers, but no such association was seen within sibships discordant in their exposure to maternal diabetes in pregnancy; these findings reflect the role of the shared environment (specifically SES). However, few of the included studies matched control subjects on the basis of SES¹⁹ or adjusted for SES¹⁶ in the relationships between maternal diabetes during pregnancy and offspring's cognitive development. Likewise, other potential confounders, including maternal age,^{49,50} alcohol use,^{51,52} and smoking during pregnancy,⁵³ and offspring-related covariates, such as gestational age⁵⁴ and birth weight,^{27,55} were rarely adjusted for in these studies.

A review by Ornoy⁵⁶ concluded that the cognitive ability of offspring born to mothers with well-controlled diabetes is usually normal. Despite substantial challenges, maternal metabolism control has multiple benefits, including but not limited to reduction of perinatal adverse outcomes and long-term effects in offspring exposed to maternal diabetes in pregnancy.⁵⁷ However, the conclusion was offered from limited, mainly small^{18,20,21} and descriptive^{26,27,58} studies. In addition, mixed results have been reported, with null or positive associations between different measures of maternal metabolic control and diverse measures of cognitive ability.^{26,29,33,39} Even with well-controlled maternal diabetes, offspring's cognitive development has been found to be significantly

impaired.^{19,31} Regardless of maternal metabolic control, in our systematic review, 1 study¹⁶ fully adjusted for potential confounders with relatively good power, as well as other studies^{19,21,29-33} that partially adjusted for potential confounders, reported significantly impaired cognitive development in offspring born to mothers with either PDM or GDM.

In contrast, in school-aged Indian children born to a population with higher rates of diabetes in pregnancy (6.9%), Veena et al²³ reported higher cognitive scores on a variety of cognitive measurements in OGDM compared with offspring of women without GDM. The investigators reported that GDM has been positively associated with higher SES, unlike in developed nations, and hence the observed association could be attributable to residual SES confounders. Although there are no data regarding the maximum threshold of maternal glucose level that is deleterious to the growing fetus, Veena et al and others^{29,48} reported positive associations between maternal glucose levels during pregnancy and offspring's cognitive scores. An optimal maternal glucose level during pregnancy is vital for the growing fetus. However, tight glycemic control is usually associated with hypoglycemia, a condition which, if it occurs repeatedly, could lead to impaired cognitive development.⁵⁹

The observed discrepancies between some of the studies that found negative,^{30,33} null,^{18,20,22} and positive associations²³ could also be partly due to cognitive test variations and the subsequent different cognitive outcomes. The effect of maternal diabetes in pregnancy may vary according to specific types of cognitive domains, as reflected by heterogeneous effect sizes. However, we are unaware

of any evidence supporting this suggestion. Various cognitive development measurements of offspring, obtained from self-report to highly standardized and blindly administered tools, were used. In addition, despite the utilization of standard cognitive measurement tools, one-half of the studies^{16,17,22,23, 29,31,33} failed to provide or report blind outcome assessment. Assessors' awareness of exposure status in randomized controlled trials has been found to be a source of information bias⁶⁰; it may also be a problem in observational cohort studies that did not commence as case-control designs. Likewise, maternal diabetes status was obtained either from hospital records, interview, or from objectively measured but varied glucose tolerance test results. These variations in measurement may also have contributed to the observed inconsistencies (eg, between Nomura et al¹⁷ and Fraser et al¹⁶). Moreover, universal GDM screening was not available in these studies,^{16,17} and it is possible that control subjects could have GDM. If so, this finding may have diluted the associations.

Some of the strengths of this systematic review are that it included studies without time restriction, employed independent reviewers, and used a standardized quality assessment tool. However, publication bias in the included studies is an inevitable limitation. A recent systematic review revealed a strong tendency for publication of positive or significant results.⁶¹ In addition, we restricted our systematic review to published articles and did not contact experts for additional data. Although we excluded limited studies through language restriction, relevant studies could have been missed from this restriction. More importantly, meta-analysis was not performed because of the limited number of studies

with diverse cognitive outcomes and model adjustment.

CONCLUSIONS

Few data are available regarding the influence of maternal diabetes during pregnancy on the subsequent cognitive development of offspring. The present review found that maternal diabetes during pregnancy seems negatively associated with offspring's childhood cognitive development. The effect was substantial in the offspring's language development during a young age, particularly verbal IQ. However, the extent to which

the observed association is due to potential confounders such as prepregnancy obesity, maternal SES, or other confounders is unclear. The use of diverse cognitive and diabetes assessment tools and criteria, sample size, and population differences contributed to the inconsistent findings. Larger prospective studies that address potential confounders are needed to confirm the independent effect of maternal diabetes on cognitive development of offspring. Future research must also determine whether the negative association is due to maternal diabetes itself or to metabolic complications.

ABBREVIATIONS

DM:	diabetes mellitus
GDM:	gestational diabetes mellitus
MDI:	mental development index
ODM:	offspring of mothers with diabetes mellitus
OGDM:	offspring of mothers with gestational diabetes mellitus
OPDM:	offspring of mothers with pregestational diabetes mellitus
PDM:	pregestational diabetes mellitus
SES:	socioeconomic status

Address correspondence to Akilew Awoke Adane, MPH, School of Public Health, Faculty of Medicine and Biomedical Sciences, the University of Queensland, Herston Rd, Herston QLD 4006, Australia. E-mail: a.adane@uq.edu.au

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2016 by the American Academy of Pediatrics

FINANCIAL DISCLOSURE: The authors have indicated they have no financial relationships relevant to this article to disclose.

FUNDING: Mr Adane is supported by an International Postgraduate Research Scholarship and a UQ Centennial scholarship. Prof Mishra is funded by an Australian Research Council Future Fellowship (FT120100812). A/Prof Tooth's salary is paid for by the Australian Longitudinal Study on Women's Health, funded by the Australian Government Department of Health.

POTENTIAL CONFLICT OF INTEREST: The authors have indicated they have no potential conflicts of interest to disclose.

REFERENCES

1. Bhattarai MD. Three patterns of rising type 2 diabetes prevalence in the world: need to widen the concept of prevention in individuals into control in the community. *JNMA J Nepal Med Assoc.* 2009;48(174):173–179
2. Dunstan DW, Zimmet PZ, Welborn TA, et al. The rising prevalence of diabetes and impaired glucose tolerance: the Australian Diabetes, Obesity and Lifestyle Study. *Diabetes Care.* 2002;25(5):829–834
3. Ferrara A. Increasing prevalence of gestational diabetes mellitus: a public health perspective. *Diabetes Care.* 2007;30(suppl 2):S141–S146
4. Danaei G, Finucane MM, Lu Y, et al; Global Burden of Metabolic Risk Factors of Chronic Diseases Collaborating Group (Blood Glucose). National, regional, and global trends in fasting plasma glucose and diabetes prevalence since 1980: systematic analysis of health examination surveys and epidemiological studies with 370 country-years and 2.7 million participants. *Lancet.* 2011;378(9785):31–40
5. Whiting DR, Guariguata L, Weil C, Shaw J. IDF diabetes atlas: global estimates of the prevalence of diabetes for 2011 and 2030. *Diabetes Res Clin Pract.* 2011;94(3):311–321
6. Casson IF, Clarke CA, Howard CV, et al. Outcomes of pregnancy in insulin dependent diabetic women: results of a five year population cohort study. *BMJ.* 1997;315(7103):275–278
7. von Katterfeld B, Li J, McNamara B, Langridge AT. Maternal and neonatal outcomes associated with gestational diabetes in women from culturally and linguistically diverse backgrounds in western Australia. *Diabet Med.* 2012;29(3):372–377
8. Rosenberg TJ, Garbers S, Lipkind H, Chiasson MA. Maternal obesity and diabetes as risk factors for adverse pregnancy outcomes: differences among 4 racial/ethnic groups. *Am J Public Health.* 2005;95(9):1545–1551
9. Catalano PM, McIntyre HD, Cruickshank JK, et al; HAPO Study Cooperative Research Group. The hyperglycemia and adverse pregnancy outcome study: associations of GDM and obesity with pregnancy outcomes. *Diabetes Care.* 2012;35(4):780–786
10. Shand AW, Bell JC, McElduff A, Morris J, Roberts CL. Outcomes of pregnancies in women with pregestational diabetes mellitus and gestational diabetes mellitus; a

- population-based study in New South Wales, Australia, 1998-2002. *Diabet Med.* 2008;25(6):708–715
11. Silva IS, Higgins C, Swerdlow AJ, et al. Birthweight and other pregnancy outcomes in a cohort of women with pre-gestational insulin-treated diabetes mellitus, Scotland, 1979-95. *Diabet Med.* 2005;22(4):440–447
 12. Ju H, Rumbold AR, Willson KJ, Crowther CA. Borderline gestational diabetes mellitus and pregnancy outcomes. *BMC Pregnancy Childbirth.* 2008;8:31
 13. Gillman MW, Rifas-Shiman S, Berkey CS, Field AE, Colditz GA. Maternal gestational diabetes, birth weight, and adolescent obesity. *Pediatrics.* 2003;111(3). Available at: www.pediatrics.org/cgi/content/full/111/3/e221
 14. Morgan K, Rahman M, Atkinson M, et al. Association of diabetes in pregnancy with child weight at birth, age 12 months and 5 years—a population-based electronic cohort study. *PLoS One.* 2013;8(11):e79803
 15. Wright GS, Rifas-Shiman SL, Rich-Edwards JW, Taveras EM, Gillman MW, Oken E. Intrauterine exposure to gestational diabetes, child adiposity, and blood pressure. *Am J Hypertens.* 2009;22(2):215–220
 16. Fraser A, Nelson SM, Macdonald-Wallis C, Lawlor DA. Associations of existing diabetes, gestational diabetes, and glycosuria with offspring IQ and educational attainment: the Avon Longitudinal Study of Parents and Children. *Exp Diabetes Res.* 2012;2012:963735
 17. Nomura Y, Marks DJ, Grossman B, et al. Exposure to gestational diabetes mellitus and low socioeconomic status: effects on neurocognitive development and risk of attention-deficit/hyperactivity disorder in offspring. *Arch Pediatr Adolesc Med.* 2012;166(4):337–343
 18. Ornoy A, Ratzon N, Greenbaum C, Wolf A, Dulitzky M. School-age children born to diabetic mothers and to mothers with gestational diabetes exhibit a high rate of inattention and fine and gross motor impairment. *J Pediatr Endocrinol Metab.* 2001;14(suppl 1):681–689
 19. Ornoy A, Wolf A, Ratzon N, Greenbaum C, Dulitzky M. Neurodevelopmental outcome at early school age of children born to mothers with gestational diabetes. *Arch Dis Child Fetal Neonatal Ed.* 1999;81(1):F10–F14
 20. Rizzo T, Metzger BE, Burns WJ, Burns K. Correlations between antepartum maternal metabolism and child intelligence. *N Engl J Med.* 1991;325(13):911–916
 21. Sells CJ, Robinson NM, Brown Z, Knopp RH. Long-term developmental follow-up of infants of diabetic mothers. *J Pediatr.* 1994;125(1):S9–S17
 22. Townsend EL, Georgieff MK, Nelson CA. Neurobehavioral functioning in five-year-old children born to diabetic mothers. *Cogniție Creier Comportament.* 2005;9(2):363–381
 23. Veena SR, Krishnaveni GV, Srinivasan K, et al. Childhood cognitive ability: relationship to gestational diabetes mellitus in India. *Diabetologia.* 2010;53(10):2134–2138
 24. Cukierman T, Gerstein HC, Williamson JD. Cognitive decline and dementia in diabetes—systematic overview of prospective observational studies. *Diabetologia.* 2005;48(12):2460–2469
 25. Biessels GJ, Gispen WH. The impact of diabetes on cognition: what can be learned from rodent models? *Neurobiol Aging.* 2005;26(1 suppl 1):36–41
 26. Persson B, Gentz J. Follow-up of children of insulin-dependent and gestational diabetic mothers. Neuropsychological outcome. *Acta Paediatr Scand.* 1984;73(3):349–358
 27. Stehbens JA, Baker GL, Kitchell M. Outcome at ages 1, 3, and 5 years of children born to diabetic women. *Am J Obstet Gynecol.* 1977;127(4):408–413
 28. Churchill JA, Berendes HW, Nemore J. Neuropsychological deficits in children of diabetic mothers. A report from the Collaborative Study of Cerebral Palsy. *Am J Obstet Gynecol.* 1969;105(2):257–268
 29. Bonilla C, Lawlor DA, Ben-Shlomo Y, et al. Maternal and offspring fasting glucose and type 2 diabetes-associated genetic variants and cognitive function at age 8: a Mendelian randomization study in the Avon Longitudinal Study of Parents and Children. *BMC Med Genet.* 2012;13:90
 30. Dionne G, Boivin M, Séguin JR, Pérusse D, Tremblay RE. Gestational diabetes hinders language development in offspring. *Pediatrics.* 2008;122(5). Available at: www.pediatrics.org/cgi/content/full/122/5/e1073
 31. Hod M, Levy-Shiff R, Lerman M, Schindel B, Ben-Rafael Z, Bar J. Developmental outcome of offspring of pregestational diabetic mothers. *J Pediatr Endocrinol Metab.* 1999;12(6):867–872
 32. Nelson CA, Wewerka SS, Borscheid AJ, Deregnier RA, Georgieff MK. Electrophysiologic evidence of impaired cross-modal recognition memory in 8-month-old infants of diabetic mothers. *J Pediatr.* 2003;142(5):575–582
 33. Yamashita Y, Kawano Y, Kuriya N, et al. Intellectual development of offspring of diabetic mothers. *Acta Paediatr.* 1996;85(10):1192–1196
 34. Wells G, Shea B, O'Connell D, et al. The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomised studies in meta-analyses. Available at: www.ohri.ca/programs/clinical_epidemiology/oxford.asp. Accessed September 9, 2015
 35. Rizzo TA, Ogata ES, Dooley SL, Metzger BE, Cho NH. Perinatal complications and cognitive development in 2- to 5-year-old children of diabetic mothers. *Am J Obstet Gynecol.* 1994;171(3):706–713
 36. Ornoy A, Ratzon N, Greenbaum C, Peretz E, Soriano D, Dulitzky M. Neurobehaviour of school age children born to diabetic mothers. *Arch Dis Child Fetal Neonatal Ed.* 1998;79(2):F94–F99
 37. DeBoer T, Wewerka S, Bauer PJ, Georgieff MK, Nelson CA. Explicit memory performance in infants of diabetic mothers at 1 year of age. *Dev Med Child Neurol.* 2005;47(8):525–531
 38. Deregnier RA, Nelson CA, Thomas KM, Wewerka S, Georgieff MK. Neurophysiologic evaluation of auditory recognition memory in

- healthy newborn infants and infants of diabetic mothers. *J Pediatr*. 2000;137(6):777–784
39. Nelson CA, Wewerka S, Thomas KM, Tribby-Walbridge S, deRegnier R, Georgieff M. Neurocognitive sequelae of infants of diabetic mothers. *Behav Neurosci*. 2000;114(5):950–956
 40. Olokoba AB, Obateru OA, Olokoba LB. Type 2 diabetes mellitus: a review of current trends. *Oman Med J*. 2012;27(4):269–273
 41. Chu SY, Callaghan WM, Kim SY, et al. Maternal obesity and risk of gestational diabetes mellitus. *Diabetes Care*. 2007;30(8):2070–2076
 42. Nguyen NT, Nguyen XM, Lane J, Wang P. Relationship between obesity and diabetes in a US adult population: findings from the National Health and Nutrition Examination Survey, 1999–2006. *Obes Surg*. 2011;21(3):351–355
 43. Basatemur E, Gardiner J, Williams C, Melhuish E, Barnes J, Sutcliffe A. Maternal prepregnancy BMI and child cognition: a longitudinal cohort study. *Pediatrics*. 2013;131(1):56–63
 44. Bliddal M, Olsen J, Støvring H, et al. Maternal pre-pregnancy BMI and intelligence quotient (IQ) in 5-year-old children: a cohort based study. *PLoS One*. 2014;9(4):e94498
 45. Huang L, Yu X, Keim S, Li L, Zhang L, Zhang J. Maternal prepregnancy obesity and child neurodevelopment in the Collaborative Perinatal Project. *Int J Epidemiol*. 2014;43(3):783–792
 46. Ronfani L, Vecchi Brumatti L, Mariuz M, et al. The complex interaction between home environment, socioeconomic status, maternal IQ and early child neurocognitive development: a multivariate analysis of data collected in a newborn cohort study. *PLoS One*. 2015;10(5):e0127052
 47. Wong HS, Edwards P. Nature or nurture: a systematic review of the effect of socio-economic status on the developmental and cognitive outcomes of children born preterm. *Matern Child Health J*. 2013;17(9):1689–1700
 48. Fraser A, Almqvist C, Larsson H, Långström N, Lawlor DA. Maternal diabetes in pregnancy and offspring cognitive ability: sibling study with 723,775 men from 579,857 families. *Diabetologia*. 2014;57(1):102–109
 49. Morinis J, Carson C, Quigley MA. Effect of teenage motherhood on cognitive outcomes in children: a population-based cohort study. *Arch Dis Child*. 2013;98(12):959–964
 50. Sutcliffe AG, Barnes J, Belsky J, Gardiner J, Melhuish E. The health and development of children born to older mothers in the United Kingdom: observational study using longitudinal cohort data. *BMJ*. 2012;345:e5116
 51. Willford J, Leech S, Day N. Moderate prenatal alcohol exposure and cognitive status of children at age 10. *Alcohol Clin Exp Res*. 2006;30(6):1051–1059
 52. Coles CD, Brown RT, Smith IE, Platzman KA, Erickson S, Falek A. Effects of prenatal alcohol exposure at school age. I. Physical and cognitive development. *Neurotoxicol Teratol*. 1991;13(4):357–367
 53. Clifford A, Lang L, Chen R. Effects of maternal cigarette smoking during pregnancy on cognitive parameters of children and young adults: a literature review. *Neurotoxicol Teratol*. 2012;34(6):560–570
 54. Woythaler MA, McCormick MC, Smith VC. Late preterm infants have worse 24-month neurodevelopmental outcomes than term infants. *Pediatrics*. 2011;127(3). Available at: www.pediatrics.org/cgi/content/full/127/3/e622
 55. Shenkin SD, Starr JM, Deary IJ. Birth weight and cognitive ability in childhood: a systematic review. *Psychol Bull*. 2004;130(6):989–1013
 56. Ornoy A. Growth and neurodevelopmental outcome of children born to mothers with pregestational and gestational diabetes. *Pediatr Endocrinol Rev*. 2005;3(2):104–113
 57. Persaud OD. Maternal diabetes and the consequences for her offspring. *J Dev Disabil*. 2007;13(1):101–134
 58. Cummins M, Norrish M. Follow-up of children of diabetic mothers. *Arch Dis Child*. 1980;55(4):259–264
 59. ter Braak EW, Evers IM, Willem Erkelens D, Visser GH. Maternal hypoglycemia during pregnancy in type 1 diabetes: maternal and fetal consequences. *Diabetes Metab Res Rev*. 2002;18(2):96–105
 60. Hróbjartsson A, Boutron I. Blinding in randomized clinical trials: imposed impartiality. *Clin Pharmacol Ther*. 2011;90(5):732–736
 61. Dwan K, Gamble C, Williamson PR, Kirkham JJ; Reporting Bias Group. Systematic review of the empirical evidence of study publication bias and outcome reporting bias—an updated review. *PLoS One*. 2013;8(7):e66844

Diabetes in Pregnancy and Childhood Cognitive Development: A Systematic Review

Akilew Awoke Adane, Gita D. Mishra and Leigh R. Tooth

Pediatrics 2016;137;

DOI: 10.1542/peds.2015-4234 originally published online April 21, 2016;

Updated Information & Services

including high resolution figures, can be found at:
<http://pediatrics.aappublications.org/content/137/5/e20154234>

References

This article cites 58 articles, 14 of which you can access for free at:
<http://pediatrics.aappublications.org/content/137/5/e20154234#BIBL>

Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):
Developmental/Behavioral Pediatrics
http://www.aappublications.org/cgi/collection/development:behavioral_issues_sub
Endocrinology
http://www.aappublications.org/cgi/collection/endocrinology_sub
Diabetes Mellitus
http://www.aappublications.org/cgi/collection/diabetes_mellitus_sub

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
<http://www.aappublications.org/site/misc/Permissions.xhtml>

Reprints

Information about ordering reprints can be found online:
<http://www.aappublications.org/site/misc/reprints.xhtml>

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Diabetes in Pregnancy and Childhood Cognitive Development: A Systematic Review

Akilew Awoke Adane, Gita D. Mishra and Leigh R. Tooth
Pediatrics 2016;137;

DOI: 10.1542/peds.2015-4234 originally published online April 21, 2016;

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/137/5/e20154234>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2016 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

