

Vaccines and Febrile Seizures: Quantifying the Risk

Mark H. Sawyer, MD, FAAP,^a Geoff Simon, MD, FAAP,^b Carrie Byington, MD, FAAP^c



Vaccines can cause fever, and fever in young children can lead to febrile seizures; these facts are not new. Febrile seizures are the most common seizure disorder of childhood and occur in ~5% of all children, usually those younger than 24 months. We know that vaccines, often administered to children in this age group, can trigger febrile seizures. The article by Duffy et al¹ in this edition of *Pediatrics* provides pediatric practitioners with a fairly precise estimate of how often influenza vaccine, conjugated pneumococcal vaccine (PCV), and diphtheria-tetanus-acellular pertussis (DTaP) vaccines given alone or in various combinations lead to febrile seizures. This study, conducted by the Vaccine Safety Datalink² (VSD), and others like it, are important as we engage in dialogue with parents about the risks and benefits of vaccines.

The VSD, started in 1990, is a collaborative project between the Centers for Disease Control and Prevention's Immunization Safety Office and 9 health care organizations and was created to address the upsurge of concerns about vaccine safety that have marked the past 2 decades. By leveraging the collective patient populations enrolled in these large organizations and the ability to evaluate the electronic medical records of those patients, the VSD can tell us scientifically, rapidly, and with good precision what happens to people after they receive vaccines. It is perhaps the best tool we have to assess vaccine safety.

Duffy et al¹ used the VSD to follow up on observations made between

2010 and 2012 linking specific influenza vaccine products, sometimes given with other vaccines, to febrile seizures.^{3,4} In the current study, the VSD used its patient power to carefully evaluate the frequency of febrile seizures after influenza vaccine given during 5 seasons (2006–2007 through 2010–2011) either alone or in combinations with PCV and DTaP because these vaccines are recommended to be given together routinely. Each of these vaccines alone was already known to cause fever, so it was reasonable to think that the potential additive fevers caused when these vaccines are given at the same time could lead to febrile seizures. The authors found that, when given alone, only PCV vaccine, but none of the influenza vaccines or DTaP, were associated with an increased rate of febrile seizures. What about the combination of all 3? The answer is that febrile seizures happen, but not often. As reported by Duffy et al,¹ influenza, DTaP, and PCV vaccines given together can lead to febrile seizures at a rate of up to 30 in 100 000 children immunized. This means for the average pediatrician, who may care for 1000 children younger than 5 including 3 to 500 between 6 and 24 months of age annually, one could expect to see at most 1 child who experiences a febrile seizure every 5 to 10 years due to administration of these vaccines together in the first 2 years of life.⁵ This would be in addition to the 30 to 75 patients in each birth year cohort in a practice that would experience a febrile seizure from other causes given the background rate of 2% to 5%.

^aUniversity of California, San Diego Department of Pediatrics and Rady Children's Hospital San Diego, San Diego, California; ^bNemours duPont Pediatrics, Wilmington, Delaware; and ^cDepartment of Pediatrics, University of Utah, Salt Lake City, Utah

Drs Sawyer and Simon are members and Dr Byington is the Chair of the American Academy of Pediatrics Committee on Infectious Disease. Dr Simon is the Chair of the American Academy of Pediatrics Committee on Practice and Ambulatory Medicine.

Opinions expressed in these commentaries are those of the author and not necessarily those of the American Academy of Pediatrics or its Committees.

DOI: 10.1542/peds.2016-0976

Accepted for publication Apr 6, 2016

Address correspondence to Mark H. Sawyer, MD, 3020 Children's Way, #5124, San Diego, CA 92123. E-mail: mhsawyer@ucsd.edu

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2016 by the American Academy of Pediatrics

FINANCIAL DISCLOSURE: Dr Byington has intellectual property in and receives royalties from BioFire Diagnostics, Inc (Salt Lake City, UT); Drs Sawyer and Simon have indicated they have no financial relationships relevant to this article to disclose.

FUNDING: Dr Byington is supported by the HA and Edna Benning Presidential Endowment; National Center for Advancing Translational Sciences of the National Institutes of Health under award 1ULTR001067. The content is solely the

To cite: Sawyer MH, Simon G, Byington C. Vaccines and Febrile Seizures: Quantifying the Risk. *Pediatrics*. 2016;138(1):e20160976

Does this mean we should stop giving these vaccines together or stop giving them at all? We say, emphatically, no. With the results of this study, we can accurately calculate the risks and benefits of this practice. The risk is 1 febrile seizure per pediatric practice every 5 to 10 years. Febrile seizures, although frightening to parents, rarely have any long-term sequelae.^{6,7} The benefits of giving these vaccines simultaneously include decreased office visits associated with painful vaccines, decreased episodes of vaccine-associated fussiness, and, most important, the assurance that children will be fully immunized and protected from infections that carry real morbidity and mortality. It is

well established that the vaccines we miss when we fail to give all the vaccines we can (simultaneously at each health care visit) may never be administered to some children, thus leaving them at risk for the diseases the vaccines prevent.⁸ It goes without saying that influenza, diphtheria, tetanus, pertussis, and pneumococcal infections may result in serious illness. These infections also have the potential to cause fevers and febrile seizures. Without vaccines to prevent these illnesses, pediatricians would see many more than 1 case of most of these infections each decade. In fact, they would see children in their practices with both febrile seizures and life-threatening

infections. The risk from these diseases far outweighs the risk from the vaccines. Fortunately, because of the surveillance and research of the VSD, we no longer need to wonder how often adverse events happen after vaccinations; instead, we can measure them scientifically, and studies like that by Duffy et al¹ increase our confidence in vaccines.

ABBREVIATIONS

DTaP: diphtheria-tetanus-acellular pertussis
PCV: conjugated pneumococcal vaccine
VSD: Vaccine Safety Datalink

responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. Funded by the National Institutes of Health (NIH).

POTENTIAL CONFLICT OF INTEREST: The authors have indicated they have no potential conflicts of interest to disclose.

COMPANION PAPER: A companion to this article can be found online at www.pediatrics.org/cgi/doi/10.1542/peds.2016-0320.

REFERENCES

1. Duffy J, Weintraub E, Hambidge SJ, et al. Risk of febrile seizure following vaccination among children age 6 through 23 months. *Pediatrics*. 2016;138(1):e20160320
2. Centers for Disease Control and Prevention. Vaccine Safety Datalink. Available at: www.cdc.gov/vaccinesafety/ensuringsafety/monitoring/vsd/. Accessed March 10, 2016
3. Broder KR, Martin DB, Vellozzi C. In the heat of a signal: responding to a vaccine safety signal for febrile seizures after 2010–11 influenza vaccine in young children, United States. *Vaccine*. 2012;30(11):2032–2034
4. Armstrong PK, Dowse GK, Effler PV, et al. Epidemiological study of severe febrile reactions in young children in Western Australia caused by a 2010 trivalent inactivated influenza vaccine. *BMJ Open*. 2011;1(1):e000016
5. American Academy of Pediatrics. Profile of Pediatric Visits. 2010. Available at: <https://www.aap.org/en-us/professional-resources/practice-support/financing-and-payment/Billing-and-Payment/pages/Profile-of-Pediatric-Office-Visits.aspx>
6. Verity CM, Greenwood R, Golding J. Long-term intellectual and behavioral outcomes of children with febrile convulsions. *N Engl J Med*. 1983;308(24):1723–1728
7. American Academy of Pediatrics; Subcommittee on Febrile Seizures. Neurodiagnostic evaluation of the child with a simple febrile seizure. *Pediatrics*. 2011;127(2):389–394
8. Hammer LD, Curry ES, Harlor AD, et al; Committee on Practice and Ambulatory Medicine; Council on Community Pediatrics. Increasing immunization coverage. *Pediatrics*. 2010;125(6):1295–1304

Vaccines and Febrile Seizures: Quantifying the Risk

Mark H. Sawyer, Geoff Simon and Carrie Byington

Pediatrics 2016;138;

DOI: 10.1542/peds.2016-0976 originally published online June 6, 2016;

Updated Information & Services

including high resolution figures, can be found at:
<http://pediatrics.aappublications.org/content/138/1/e20160976>

References

This article cites 4 articles, 3 of which you can access for free at:
<http://pediatrics.aappublications.org/content/138/1/e20160976#BIBL>

Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):
Infectious Disease
http://www.aappublications.org/cgi/collection/infectious_diseases_sub
Vaccine/Immunization
http://www.aappublications.org/cgi/collection/vaccine:immunization_sub
Preventive Medicine
http://www.aappublications.org/cgi/collection/preventative_medicine_sub

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
<http://www.aappublications.org/site/misc/Permissions.xhtml>

Reprints

Information about ordering reprints can be found online:
<http://www.aappublications.org/site/misc/reprints.xhtml>

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®



PEDIATRICS[®]

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Vaccines and Febrile Seizures: Quantifying the Risk

Mark H. Sawyer, Geoff Simon and Carrie Byington

Pediatrics 2016;138;

DOI: 10.1542/peds.2016-0976 originally published online June 6, 2016;

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/138/1/e20160976>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 345 Park Avenue, Itasca, Illinois, 60143. Copyright © 2016 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN[®]

