

Screening in Primary Care: What Is the Best Way to Identify At-Risk Youth for Substance Use?

Elizabeth J. D'Amico, PhD,^a Layla Parast, PhD,^a Lisa S. Meredith, PhD,^a Brett A. Ewing, MS,^a William G. Shadel, PhD,^b Bradley D. Stein, PhD^b

abstract

BACKGROUND: It is important to improve primary care providers' capability to identify youth at risk for alcohol and other drug use. To our knowledge, this is the first study to use *Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition* criteria to compare screeners for youth for both alcohol and marijuana, given that these are the most frequently used substances by this age group.

METHODS: We compared the psychometric performance of 4 screeners: the National Institute on Alcohol Abuse and Alcoholism Screening Guide (NIAAA SG), the Alcohol Use Disorders Identification Test, the Car-Relax-Alone-Forget-Family and Friends-Trouble (CRAFFT) screener, and the Personal Experience Screening Questionnaire Problem Severity Scale (PESQ-PS) in identifying alcohol and marijuana use outcomes. Youth age 12 through 18 ($N = 1573$; 27% black, 51% Hispanic) were screened with the NIAAA SG, followed by a Web survey that included the other screeners and outcomes.

RESULTS: Sensitivity for alcohol outcomes indicated that the NIAAA SG (0.87) did not perform as well as the CRAFFT (0.97) or PESQ-PS (0.97) screeners but performed better than the Alcohol Use Disorders Identification Test (0.70). The pattern for sensitivity across screeners for marijuana outcomes was similar.

CONCLUSIONS: An important tradeoff in primary care settings is precision versus practicality. Because of brevity and focus on frequency of drinking, the NIAAA SG offers ease of administration and is good at identifying youth with probably problematic drinking levels. The PESQ-PS and the CRAFFT correctly identify more at-risk youth for alcohol and marijuana than the NIAAA SG. Future work is needed to elucidate how to efficiently and accurately identify at-risk youth in the primary care setting, including determining the best cutoff points to use to increase sensitivity.



^aRAND Corporation, Santa Monica, California; and ^bRAND Corporation, Pittsburgh, Pennsylvania

Dr D'Amico conceptualized and designed the study and drafted the initial manuscript; Dr Parast and Ms Ewing carried out initial analyses and reviewed and revised the manuscript; Drs Meredith, Shadel, and Stein reviewed and revised the manuscript; and all authors approved the final manuscript as submitted.

This trial has been registered at www.clinicaltrials.gov (identifier NCT01797835).

DOI: 10.1542/peds.2016-1717

Accepted for publication Sep 27, 2016

Address correspondence to Elizabeth J. D'Amico, PhD, RAND Corporation, 1776 Main St, Santa Monica, CA 90401. E-mail: damico@rand.org

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

WHAT'S KNOWN ON THIS SUBJECT: Most adolescents are not screened for substance use in primary care because of time constraints or insufficient training. Providers need a screener that can be easily incorporated into an appointment; however, there is a tradeoff between precision and practicality.

WHAT THIS STUDY ADDS: We examined sensitivity, specificity, and positive and negative predictive value for the new National Institute on Alcohol Abuse and Alcoholism screener and 3 widely used adolescent screeners for various levels of alcohol and marijuana use and impairment in a large, racially and ethnically diverse sample of adolescents.

To cite: D'Amico EJ, Parast L, Meredith LS, et al. Screening in Primary Care: What Is the Best Way to Identify At-Risk Youth for Substance Use? *Pediatrics*. 2016;138(6):e20161717

It is important to improve the capability of primary care providers and associated health care staff to identify youth at risk for alcohol and other drug (AOD) use because use during adolescence is associated with academic problems, poorer mental health, future use of other illicit drugs (including heroin and cocaine), and a higher likelihood of abuse or dependence in adulthood.¹⁻³ The primary care setting provides a unique opportunity to screen significant numbers of adolescents⁴ and identify those at risk for problematic use. Guidelines propose that providers screen for AOD use and provide brief counseling and referrals where appropriate⁵ and that doctors screen all patients for alcohol use starting in middle school.⁶

Unfortunately, most adolescents are not screened for AOD use in primary care settings,⁷⁻¹¹ and significant numbers of at-risk youth remain unidentified and never receive appropriate preventive or treatment services.¹²⁻¹⁴ Lack of screening and preventive services is even more profound among younger adolescents age 11 to 14 and socioeconomically disadvantaged youth.^{15,16} Lack of primary care screening typically is caused by provider time constraints, discomfort discussing AOD use, insufficient training, or lack of referral options.^{9,10,17,18}

Given these concerns, providers need an easy-to-administer screener that takes little time or training, can be incorporated into a primary care appointment to determine an adolescent's risk level, and will facilitate appropriate referral or treatment. There is a difficult tradeoff between precision and practicality in the primary care setting. To increase screening in pediatric settings, the National Institute on Alcohol Abuse and Alcoholism (NIAAA) developed a brief 2-item screening guide (SG) with 2 age-specific

screening questions about friend and self-drinking.¹⁹ Age-sensitive cutoff points can assist providers in determining whether brief advice, counseling, or referral is appropriate. For example, any report of drinking if the youth is ≤ 15 years old warrants brief advice and counseling. For youth ≥ 16 years old, the threshold is a bit higher, and drinking ≥ 6 days in the past year is considered moderate or high risk.

In 2014, Kelly et al¹¹ evaluated the NIAAA SG in a sample of 525 youth age 12 to 17 years (54% female) who were mainly black (92.8%). Their focus was on expanding the NIAAA SG to include drug and tobacco questions for a total of 10 questions and determining cutoff points for their Brief Screener for Tobacco, Alcohol and Other Drugs. They established cutoff points for this new screener for each substance and compared the Brief Screener for Tobacco, Alcohol and Other Drugs cutoff points with the *Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition* (DSM-5) criteria.¹¹ In 2016, Clark et al²⁰ examined cutoff points for the NIAAA SG and found that ≤ 3 days of alcohol use in the past year yielded optimal psychometric performance.

Several studies²¹⁻²⁵ have reviewed adolescent screeners to determine performance against DSM-4 and DSM-5 criteria. Some screeners focus only on alcohol, such as the NIAAA SG, whereas others address AOD more broadly. A comparison of the Alcohol Use Disorders Identification Test (AUDIT; 10 items; alcohol only), Car-Relax-Along-Forget-Family and Friends-Trouble (CRAFFT; 6 items; AOD), Cut Down, Annoyed, Guilty, Eye-Opener (CAGE; 4 items; alcohol only), and Tolerance, Worry, Eye-Opener, Cutoff (TWEAK; 5 items; alcohol only) in a sample of youth recruited in an emergency department found that Cut Down, Annoyed, Guilty, Eye-Opener items were reported infrequently by

adolescents and that the AUDIT performed the best across the range of cutoff scores.²¹ The Problem Severity Scale on the Personal Experience Screening Questionnaire (PESQ-PS; 18 items; AOD)^{23,26,27} had the highest reliability estimates when compared with several other screening measures that address behavior and consequences for youth, such as the Rutgers Alcohol Problem Index.²⁸ Furthermore, the 2011 American Academy of Pediatrics guidelines recommended routine use of the CRAFFT, which measures AOD use,²⁹ has been validated in diverse populations,^{30,31} and has good sensitivity and specificity with new DSM-5 criteria.³²

Thus, across diverse studies, the AUDIT, CRAFFT, and PESQ-PS appear to be the most sensitive and reliable screeners with adolescents. They are short, taking ~5 to 10 minutes to complete and score. The NIAAA SG¹⁹ could also easily be integrated into primary care practice given its brevity,¹¹ and studies have shown that asking youth about alcohol use frequency is often the best predictor of their alcohol use over time.^{33,34} We therefore compared these 4 screeners in identifying both alcohol and marijuana use among adolescents because these substances are the most frequently reported substances for this age group.³⁵

This study moves the field forward by comparing 4 screeners in a large racially and ethnically diverse sample of adolescents, ages 12 through 18, recruited across 4 primary care clinics in Los Angeles, California and Pittsburgh, Pennsylvania. The study's purpose was to inform primary care providers seeking to screen adolescents for alcohol and marijuana use about the 4 screeners' strengths and weaknesses; we compared screeners with respect to sensitivity, specificity, positive

predictive value (PPV), and negative predictive value (NPV) for various levels of alcohol and marijuana use and impairment.

METHODS

Procedures

This study involved 4 clinics (1 in Los Angeles, 3 in Pittsburgh). We obtained a certificate of confidentiality; procedures were approved by both the RAND institutional review board and the 4 clinics. Every youth age 12 through 18 who came for an appointment during the 2.5-year study period (April 2013–November 2015) was asked to be in the project. We obtained parental consent and youth assent (<18 years old) or consent (18 years old). Youth were screened with the NIAAA SG, completed a survey via the Web, and paid \$25.

Setting, Participants, and Surveys

The Pittsburgh and Los Angeles area clinics are family-based community health clinics that provide care for ethnically and racially diverse and underserved populations of youth. Sites offer both longitudinal, continuity-based care and episode-based urgent care to their patients. Clinics in both cities have a large percentage of minority patients and serve a high proportion of low-income patients. Approximately 3309 youth were approached to be in the project. Of these youth, 27% ($n = 892$) were ineligible because of age, lack of English proficiency, being present for an appointment other than their own, or disability status; 18.5% ($n = 614$) declined to participate, mostly because of time constraints or youth being at the clinic for family planning and not wanting their parents to know they were there. This process yielded a total sample of 1803 youth who enrolled or provided consent to

contact. Of the 1803 youth, 230 did not complete the baseline within the field period or had unreliable contact information. The final enrolled sample included 1573 youth. Screening and surveys were completed in a private clinic space without a parent present. RAND staff first screened youth in person by using the NIAAA SG; youth then completed a Web-based survey comprising other screeners and outcome variables on a laptop immediately after completion of the NIAAA SG (see Supplemental Information for screener questions).

Screeners

NIAAA SG

Two screening questions were asked in a different order depending on age¹⁹: “In the past year, on how many days have you had more than a few sips of beer, wine, or any drink containing alcohol?” and “Do any of your friends drink alcohol?” Youth age 12 to 14 years were first asked about friend drinking and then self-drinking as a less threatening way to gauge use, whereas youth age ≥ 15 years (and 14-year-olds in high school) were first asked about self-drinking and then friend drinking. Adolescents were categorized based solely on the days of use question according to the published NIAAA risk assessment guide, which results in 4 risk categories: no risk, lower risk, moderate risk, and highest risk. For example, youth age 12 to 15 years were categorized as moderate risk if they reported 1 to 5 days of use; adolescents 16 years old were categorized as moderate risk if they reported 6 to 11 days of use.

CRAFFT

The 6-item CRAFFT²⁹ addresses both alcohol and other drugs (eg, “Do you ever use AOD to *Relax*, feel better about yourself, or fit in?”). Response options are “yes” or “no,” and a “yes”

response to ≥ 2 questions indicates risk.

PESQ-PS

The 18-question PESQ-PS²⁶ assesses AOD use rated on a 4-point response scale (never, once or twice, sometimes, often). The summed score categorizes adolescents into 3 groups, no AOD problem (“Green Flag”), mild or moderate AOD problem (“Yellow Flag”), and severe AOD problem (“Red Flag”), by using established thresholds. Thresholds vary depending on sex and age.

AUDIT

The AUDIT³⁶ focuses on frequency and consequences of drinking. It was modified slightly to be developmentally appropriate for youth.²¹ Youth are categorized into Zone I (alcohol education), Zone II (simple advice), Zone III (simple advice plus brief intervention and follow-up), or Zone IV (referral to specialist).

Outcomes

Youth with an alcohol use disorder (AUD) or cannabis use disorder (CUD) were identified via the Diagnostic Interview Schedule for Children Version IV (DISC-IV)³⁷ computerized version, valid and reliable in adolescent populations.^{38–40} We used DSM-5 criteria available in this version (eg, the craving item was included; see Supplemental Figure) to identify subjects with AUD or CUD. Adolescents were classified as having an AUD or CUD if they reported ≥ 2 of the 11 criteria for AUD or CUD. Past-year alcohol and marijuana use was assessed via well-established measures with adolescents.⁴¹ We asked, “During the past year, how many times did you [drink at least one full drink of alcohol] [use marijuana]?” Responses ranged from 1 = “0 times” to 6 = “11–20 times” and were dichotomized (1 = “any use” versus 0 = “no use”).

TABLE 1 Sample Characteristics by Age Group

	Overall, <i>N</i> = 1573, Mean (SD) or <i>N</i> (%)	Age 12–14, <i>N</i> = 498, Mean (SD) or <i>N</i> (%)	Age 15–18, <i>N</i> = 1075, Mean (SD) or <i>N</i> (%)
Age	15.5 (1.9)	13.2 (0.8)	16.6 (1.1)
Sex			
Male	662 (42.5%)	235 (48%)	427 (40.1%)
Female	894 (57.5%)	255 (52%)	639 (59.9%)
Race or ethnicity			
White	232 (14.7%)	78 (15.7%)	154 (14.3%)
Black	420 (26.7%)	166 (33.3%)	254 (23.6%)
Hispanic	808 (51.4%)	209 (42%)	599 (55.7%)
Other or multiracial	113 (7.2%)	45 (9%)	68 (6.3%)
Prevalence: past-year use			
Alcohol use	655 (41.7%)	66 (13.3%)	589 (54.9%)
Heavy alcohol use	347 (22.1%)	24 (4.8%)	323 (30.1%)
Marijuana use	575 (36.6%)	77 (15.5%)	498 (46.4%)
Heavy marijuana use	302 (19.3%)	38 (7.7%)	264 (24.7%)
Prevalence: DSM-5 diagnosis			
AUD	61 (3.9%)	4 (0.8%)	57 (5.4%)
CUD	211 (13.6%)	23 (4.7%)	188 (17.8%)

Percentages are among nonmissing values; 17 missing sex, 10 missing past-year alcohol or marijuana use responses, 35 missing responses needed to determine DSM-5 Diagnostic Interview Schedule for Children diagnosis category.

For past-year heavy alcohol use, respondents were asked, “During the past year, how many times have you tried five or more drinks of alcohol in a row, that is, within a couple of hours?” with the same response options and dichotomization. For past-year heavy marijuana use, respondents were asked, “On days you use marijuana, how often do you use it?” Responses were “Once,” “Twice,” and “3 or more times,” and responses of “twice” and “3 or more” were considered heavy marijuana use.

Statistical Analysis

We examined the proportion of youth identified as at risk for each screener. We estimated sensitivity, specificity, PPV, and NPV against DSM-5 diagnoses of AUD and CUD, past-year any use, and past-year heavy alcohol and marijuana use. Sensitivity is the probability that the screener correctly identifies at-risk youth (as at risk), whereas specificity is the probability that the screener correctly identifies no-risk youth (as low or no risk). PPV is the probability of a case screened as positive actually being positive; this depends on specificity of the test and

prevalence of the condition. NPV is the probability of a negative diagnosis indicating a true negative. For screeners with multiple risk categories, we used the lowest established risk threshold to dichotomize adolescents into low- and high-risk categories. Specifically, for the AUDIT, youth categorized in Zone II or above were considered at risk, for the PESQ-PS, youth categorized in the yellow or red flag categories were considered at risk, and for the NIAAA SG screener, youth categorized as moderate or high risk were considered at risk. We had <2.5% of data missing for any 1 variable; pairwise deletion was used to handle missingness. We performed analyses via R version 3.2.4 (R Foundation for Statistical Computing, Vienna, Austria).

RESULTS

More than half the sample was female; youth were racially and ethnically diverse, with a mean age of 15.5 years (Table 1). More than 40% of adolescents reported drinking >1 time in the past year, with almost a quarter reporting

heavy use. About 37% reported using marijuana >1 time in the past year, with 1 in 5 reporting heavy marijuana use. According to DSM-5 criteria, 3.6% of youth were identified with an AUD; 13.6% were identified with a CUD.

Table 2 shows number and proportion of adolescents in each risk category by screener, stratified by age. Older youth were more likely to be identified as at risk. Overall, the NIAAA SG, CRAFFT, PESQ-PS, and AUDIT identified 19%, 30%, 31%, and 8% of adolescents as at risk, respectively.

Table 3 presents sensitivity, specificity, PPV, and NPV and associated 95% confidence intervals for screeners by AUD, past-year alcohol use, and past-year heavy alcohol use. For sensitivity, for all alcohol outcomes, the NIAAA SG did not perform as well as the CRAFFT or PESQ-PS screeners but performed better than the AUDIT. For AUD, the sensitivity was 0.87 for the NIAAA SG, 0.98 for the CRAFFT, 0.97 for the PESQ-PS, and 0.70 for the AUDIT. Specificity tended to be high, but it was lowest for the CRAFFT and PESQ-PS when AUD was used as the outcome. For PPV, among adolescents identified as at risk on the NIAAA SG,

TABLE 2 Number and Percentage of Youth Identified at Risk for Alcohol or Drug Use by Screener

	Overall, N = 1573, N (%)	Age 12–14, N = 498, N (%)	Age 15–18, N = 1075, N (%)
NIAAA			
No risk	992 (63.1%)	445 (89.4%)	547 (50.9%)
Lower risk	287 (18.2%)	0 (0%)	287 (26.7%)
Moderate risk	199 (12.7%)	46 (9.2%)	153 (14.2%)
Highest risk	95 (6%)	7 (1.4%)	88 (8.2%)
CRAFFT			
<2	1100 (70.2%)	445 (89.7%)	655 (61.2%)
≥2	467 (29.8%)	51 (10.3%)	416 (38.8%)
PESQ-PS			
Green Flag	1064 (69%)	432 (89.8%)	632 (59.6%)
Yellow Flag	227 (14.7%)	28 (5.8%)	199 (18.8%)
Red Flag	250 (16.2%)	21 (4.4%)	229 (21.6%)
AUDIT			
Zone I: Education	1440 (91.8%)	490 (98.6%)	950 (88.6%)
Zone II: Simple Advice	106 (6.8%)	4 (0.8%)	102 (9.5%)
Zone III: Advice and Counseling	13 (0.8%)	2 (0.4%)	11 (1%)
Zone IV: Referral	10 (0.6%)	1 (0.2%)	9 (0.8%)

Percentages are among nonmissing values; 6 youth missing ≥1 response needed to calculate CRAFFT score, 32 youth missing ≥1 response needed to calculate PESQ-PS score, 4 youth missing ≥1 response needed to calculate AUDIT score.

TABLE 3 Sensitivity, Specificity, PPV, and NPV for Each Screener, With 95% Confidence Intervals, for 3 Different Outcomes: DSM-5 Diagnosis of AUD, Past-Year Alcohol Use, and Past-Year Heavy Alcohol Use

	Sensitivity	Specificity	PPV	NPV
AUD				
NIAAA	0.87 (0.76–0.94)	0.84 (0.82–0.86)	0.19 (0.14–0.24)	0.99 (0.99–1.00)
CRAFFT	0.98 (0.91–1.00)	0.73 (0.71–0.76)	0.13 (0.10–0.17)	1.00 (0.99–1.00)
PESQ-PS	0.97 (0.88–1.00)	0.72 (0.70–0.74)	0.12 (0.09–0.15)	1.00 (0.99–1.00)
AUDIT	0.70 (0.57–0.81)	0.94 (0.93–0.96)	0.34 (0.26–0.43)	0.99 (0.98–0.99)
Past-year alcohol use				
NIAAA	0.40 (0.37–0.44)	0.97 (0.95–0.98)	0.90 (0.86–0.93)	0.69 (0.67–0.72)
CRAFFT	0.61 (0.57–0.65)	0.93 (0.91–0.94)	0.85 (0.82–0.88)	0.77 (0.74–0.79)
PESQ-PS	0.64 (0.60–0.67)	0.93 (0.91–0.94)	0.86 (0.83–0.89)	0.78 (0.75–0.80)
AUDIT	0.19 (0.16–0.22)	0.99 (0.98–1.00)	0.94 (0.88–0.97)	0.63 (0.61–0.66)
Past-year heavy alcohol use				
NIAAA	0.56 (0.51–0.61)	0.92 (0.90–0.93)	0.66 (0.60–0.71)	0.88 (0.86–0.90)
CRAFFT	0.81 (0.76–0.85)	0.84 (0.82–0.86)	0.59 (0.55–0.64)	0.94 (0.92–0.95)
PESQ-PS	0.85 (0.80–0.88)	0.84 (0.82–0.86)	0.61 (0.56–0.65)	0.95 (0.94–0.96)
AUDIT	0.33 (0.28–0.39)	0.99 (0.98–0.99)	0.89 (0.82–0.94)	0.84 (0.82–0.86)

95% confidence intervals obtained by using exact binomial confidence limits.⁴²

19% had an AUD, 90% reported past-year alcohol use, and 66% reported heavy past-year alcohol use.

Table 4 parallels Table 3 for marijuana outcomes. Similar to alcohol, the NIAAA SG performed better than the AUDIT but did not perform as well as the CRAFFT or PESQ-PS in terms of sensitivity. For example, among adolescents with CUD, the CRAFFT, PESQ-PS and NIAAA SG correctly identified 88%, 91%, and 54% as at risk, whereas the AUDIT correctly identified 32%. For PPV, among

adolescents identified as at risk on the NIAAA SG, 40% had a CUD, 77% reported past-year marijuana use, and 50% reported past-year heavy marijuana use. For both PPV and NPV, the NIAAA SG performed similarly to the CRAFFT and PESQ-PS when CUD and heavy marijuana use were examined.

DISCUSSION

To our knowledge, this is the first study to use the DSM-5 AUD and CUD criteria to compare several different screeners for youth. We

compared 4 adolescent screeners for various levels of alcohol and marijuana use and impairment with a large, racially and ethnically diverse multisite primary care population. The CRAFFT and PESQ-PS, which address AOD, identified about one-third of youth as at risk, the NIAAA SG identified ~19%, and the AUDIT identified ~8%. The CRAFFT and PESQ-PS had excellent sensitivity for detecting an AUD and also did well for CUD. The NIAAA SG, briefer and focused exclusively on alcohol, was better at identifying youth with an AUD

TABLE 4 Sensitivity, Specificity, PPV, and NPV for Each Screener, With 95% Confidence Intervals, for 3 Different Outcomes: DSM-5 Diagnosis of CUD, Past-Year Marijuana Use, and Past-Year Heavy Marijuana Use

	Sensitivity	Specificity	PPV	NPV
CUD				
NIAAA	0.54 (0.47–0.60)	0.87 (0.85–0.89)	0.40 (0.34–0.46)	0.92 (0.91–0.94)
CRAFFT	0.88 (0.83–0.92)	0.80 (0.78–0.82)	0.41 (0.36–0.45)	0.98 (0.97–0.99)
PESQ-PS	0.91 (0.86–0.94)	0.79 (0.77–0.81)	0.41 (0.37–0.46)	0.98 (0.97–0.99)
AUDIT	0.32 (0.26–0.39)	0.96 (0.94–0.97)	0.54 (0.44–0.62)	0.90 (0.88–0.92)
Past-year marijuana use				
NIAAA	0.39 (0.35–0.44)	0.93 (0.92–0.95)	0.77 (0.72–0.82)	0.73 (0.70–0.75)
CRAFFT	0.68 (0.64–0.72)	0.92 (0.90–0.94)	0.84 (0.80–0.87)	0.83 (0.81–0.86)
PESQ-PS	0.72 (0.68–0.75)	0.93 (0.91–0.94)	0.85 (0.81–0.88)	0.85 (0.83–0.87)
AUDIT	0.20 (0.17–0.24)	0.99 (0.98–0.99)	0.91 (0.84–0.95)	0.68 (0.66–0.71)
Past-year heavy marijuana use				
NIAAA	0.48 (0.42–0.54)	0.88 (0.87–0.90)	0.50 (0.44–0.56)	0.88 (0.86–0.89)
CRAFFT	0.84 (0.79–0.88)	0.83 (0.81–0.85)	0.54 (0.49–0.59)	0.96 (0.94–0.97)
PESQ-PS	0.86 (0.81–0.90)	0.82 (0.80–0.84)	0.54 (0.49–0.59)	0.96 (0.95–0.97)
AUDIT	0.29 (0.24–0.34)	0.97 (0.96–0.98)	0.67 (0.59–0.75)	0.85 (0.83–0.87)

95% confidence intervals obtained by using exact binomial confidence limits.⁴²

versus a CUD, and the AUDIT had lower sensitivity for both disorders. Screeners had higher sensitivity for identifying youth who reported past-year heavy alcohol use compared with any past-year drinking. This result is not surprising given that one would generally expect youth with heavy drinking to be easier to identify as high risk compared with youth who have had any past drinking in the last year (which may include very light drinkers). In addition, screeners that addressed AOD versus only alcohol did better at identifying youth who reported heavy past-year marijuana use and any past-year marijuana use.

Overall, specificity was lower than sensitivity for all screeners for AUD and CUD, although it was still good. Given that the potential harm of a false positive is low in this setting, whereas risks associated with a false negative are high, providers might be more willing to accept lower specificity for high sensitivity to ensure that at-risk youth are identified. As expected, the PPV was better for all screeners in identifying past-year use for alcohol and marijuana use versus AUD and CUD because the PPV depends on both specificity of the test and prevalence of the condition.

The NPV was also high for all screeners.

An important tradeoff in primary care settings is precision versus practicality. Incorporating screening into everyday practice can be difficult if the screener is long or not intuitive. Because of its brevity and focus on frequency of drinking, the NIAAA SG offers ease of administration, and results show that it is good at identifying youth with problematic drinking levels. However, the PESQ-PS and the CRAFFT correctly identify more at-risk youth for alcohol than the NIAAA SG. One recent study found that computer self-entry for the CRAFFT was valid and time-efficient.⁴³ Other work has also shown that brief screens can be completed electronically as part of routine care.⁴⁴ Future work is needed to elucidate how to most efficiently and accurately identify more at-risk youth in the primary care setting, including determining the best cutoff points to use to increase sensitivity.

Although the NIAAA SG performed well for alcohol outcomes, it did not do as well identifying youth who report marijuana use. This result is not surprising because the NIAAA SG questions focus on alcohol. However, identifying youth

at risk for marijuana use is also important, particularly because it affects more domains of functioning in adolescence than alcohol.⁴⁵ Furthermore, in this sample, 72% of youth who reported past-year alcohol use also reported past-year marijuana use, and CUD in this sample was >3 times as common as AUD, emphasizing the importance of asking about marijuana use in primary care settings. If a positive screen occurs, providers need to discuss potential harms of marijuana so youth better understand how the drug can affect functioning in both adolescence and adulthood. This discussion is particularly important because many youth view marijuana use as less harmful than alcohol use,⁴⁶ perhaps because of continuing changes in state laws regarding medical and recreational use of marijuana. Interventions as brief as 15 minutes can lead to subsequent reductions in youth AOD use,^{47,48} but time constraints, issues of confidentiality, and knowledge of what to do after a positive a screen remain common barriers to screening in the primary care setting.^{49–51} Training designed to increase provider confidence in screening and discussing AOD use with at-risk youth^{52,53} and addressing confidentiality concerns⁵¹ is 1 approach to

decreasing barriers, as is incorporating technology and providing computerized brief interventions in primary care settings for at-risk youth.⁵⁴

Limitations of the current study include self-report, although self-report limitations are often exaggerated,⁵⁵ and rates of AOD use in our sample were similar to national rates.⁴¹ The sample may also not be generalizable to all adolescents in primary care clinics; however, clinics did cross 2 states, and the sample was racially and ethnically diverse, with a wide range of ages. We also always administered the NIAAA SG to youth first, which could be a reason for lower sensitivity.⁴³ Finally, we cannot speak to performance of these screeners as clinical tools. For example, knowing their doctor will see their answers might change an adolescent's likelihood of disclosure. Future

studies should examine screener performance under more realistic clinical conditions.

CONCLUSIONS

In sum, many youth were identified as at risk by these 4 screeners; however, some screeners performed better than others, depending on the substance and severity (eg, frequency of use versus a disorder). Future work could focus on quantifying tradeoffs in precision and practicality of these screeners in primary care.

ACKNOWLEDGMENTS

The authors thank the clinics that participated and supported this project. We also thank Kirsten Becker and Jennifer Parker for overseeing the survey administrations and Michael Woodward for help with manuscript formatting.

ABBREVIATIONS

AOD: alcohol and other drug
AUD: alcohol use disorder
AUDIT: Alcohol Use Disorders Identification Test
CRAFT: Car-Relax-Alone-Forget-Family and Friends-Trouble
CUD: cannabis use disorder
DISC-IV: Diagnostic Interview Schedule for Children Version IV
DSM-5: *Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition*
NIAAA SG: National Institute on Alcohol Abuse and Alcoholism Screening Guide
NPV: negative predictive value
PESQ-PS: Personal Experience Screening Questionnaire Problem Severity Scale
PPV: positive predictive value

Copyright © 2016 by the American Academy of Pediatrics

FINANCIAL DISCLOSURE: The authors have indicated they have no financial relationships relevant to this article to disclose.

FUNDING: Supported by a grant from the National Institute of Alcohol Abuse and Alcoholism (R01AA021786) to Elizabeth D'Amico. Funded by the National Institutes of Health (NIH).

POTENTIAL CONFLICT OF INTEREST: The authors have indicated they have no potential conflicts of interest to disclose.

REFERENCES

1. Berg N, Kiviruusu O, Karvonen S, et al. A 26-year follow-up study of heavy drinking trajectories from adolescence to mid-adulthood and adult disadvantage. *Alcohol Alcohol*. 2013;48(4):452–457
2. Brook JS, Lee JY, Brown EN, Finch SJ. Comorbid trajectories of tobacco and marijuana use as related to psychological outcomes. *Subst Abuse*. 2012;33(2):156–167
3. van Gastel WA, Tempelaar W, Bun C, et al. Cannabis use as an indicator of risk for mental health problems in adolescents: a population-based study at secondary schools. *Psychol Med*. 2013;43(9):1849–1856
4. Bloom B, Cohen RA, Freeman G. Summary health statistics for US children: National Health Interview Survey, 2007. *Vital Health Stat* 10. 2009;10(239):1–80
5. Elster AB, Kuznets NK. *Guidelines for Adolescent Preventive Services (GAPS): Recommendations and Rationale*. Baltimore, MD: Williams & Wilkins; 1993
6. Kokotailo PK; Committee on Substance Abuse. Alcohol use by youth and adolescents: a pediatric concern. *Pediatrics*. 2010;125(5):1078–1087
7. Johnson SB, Millstein SG. Prevention opportunities in health care settings. *Am Psychol*. 2003;58(6–7):475–481
8. Carballo JJ, Oquendo MA, Giner L, et al. Prevalence of alcohol misuse among adolescents and young adults evaluated in a primary care setting. *Int J Adolesc Med Health*. 2006;18(1):197–202
9. Van Hook S, Harris SK, Brooks T, et al; New England Partnership for Substance Abuse Research. The “Six T’s”: barriers to screening teens for substance abuse in primary care. *J Adolesc Health*. 2007;40(5):456–461
10. Gordon AJ, Ettaro L, Rodriguez KL, Mocik J, Clark DB. Provider, patient, and family perspectives of adolescent alcohol use and treatment in rural settings. *J Rural Health*. 2011;27(1):81–90
11. Kelly SM, Gryczynski J, Mitchell SG, Kirk A, O’Grady KE, Schwartz RP. Validity

- of brief screening instrument for adolescent tobacco, alcohol, and drug use. *Pediatrics*. 2014;133(5):819–826
12. Ozer EM, Adams SH, Lustig JL, et al. Can it be done? Implementing adolescent clinical preventive services. *Health Serv Res*. 2001; 36(6 pt 2):150–165
 13. Klein JD, Allan MJ, Elster AB, et al. Improving adolescent preventive care in community health centers. *Pediatrics*. 2001;107(2):318–327
 14. Chung PJ, Lee TC, Morrison JL, Schuster MA. Preventive care for children in the United States: quality and barriers. *Annu Rev Public Health*. 2006;27:491–515
 15. Stevens GD, Shi L. Racial and ethnic disparities in the primary care experiences of children: a review of the literature. *Med Care Res Rev*. 2003;60(1):3–30
 16. Millstein SG, Marcell AV. Screening and counseling for adolescent alcohol use among primary care physicians in the United States. *Pediatrics*. 2003;111(1):114–122
 17. Halpern-Felsher BL, Ozer EM, Millstein SG, et al. Preventive services in a health maintenance organization: how well do pediatricians screen and educate adolescent patients? *Arch Pediatr Adolesc Med*. 2000;154(2):173–179
 18. Ozer EM, Adams SH, Gardner LR, Mailloux DE, Wibbelsman CJ, Irwin CE Jr. Provider self-efficacy and the screening of adolescents for risky health behaviors. *J Adolesc Health*. 2004;35(2):101–107
 19. NIAAA. *Alcohol Screening and Brief Intervention for Youth: A Practitioner's Guide*. Rockville, MD: NIAAA, DHHS and AAP; 2011. NIH publication 11-7805
 20. Clark DB, Martin CS, Chung T, et al. Screening for underage drinking and *Diagnostic and Statistical Manual of Mental Disorders, 5th Edition* alcohol use disorder in rural primary care practice. *J Pediatr*. 2016;173:214–220
 21. Chung T, Colby SM, Barnett NP, Rohsenow DJ, Spirito A, Monti PM. Screening adolescents for problem drinking: performance of brief screens against DSM-IV alcohol diagnoses. *J Stud Alcohol*. 2000;61(4):579–587
 22. Knight JR, Sherritt L, Harris SK, Gates EC, Chang G. Validity of brief alcohol screening tests among adolescents: a comparison of the AUDIT, POSIT, CAGE, and CRAFFT. *Alcohol Clin Exp Res*. 2003;27(1):67–73
 23. Shields AL, Campfield DC, Miller CS, Howell RT, Wallace K, Weiss RD. Score reliability of adolescent alcohol screening measures: a meta-analytic inquiry. *J Child Adolesc Subst Abuse*. 2008;17(4):75–97
 24. Reinert DF, Allen JP. The Alcohol Use Disorders Identification Test: an update of research findings. *Alcohol Clin Exp Res*. 2007;31(2):185–199
 25. Cook RL, Chung T, Kelly TM, Clark DB. Alcohol screening in young persons attending a sexually transmitted disease clinic. Comparison of AUDIT, CRAFFT, and CAGE instruments. *J Gen Intern Med*. 2005;20(1):1–6
 26. Winters KC. Development of an adolescent alcohol and other drug abuse screening scale: Personal Experience Screening Questionnaire. *Addict Behav*. 1992;17(5):479–490
 27. Winters KC, Kaminer Y. Screening and assessing adolescent substance use disorders in clinical populations. *J Am Acad Child Adolesc Psychiatry*. 2008;47(7):740–744
 28. White HR, Labouvie EW. Towards the assessment of adolescent problem drinking. *J Stud Alcohol*. 1989;50(1):30–37
 29. Knight JR, Sherritt L, Shrier LA, Harris SK, Chang G. Validity of the CRAFFT substance abuse screening test among adolescent clinic patients. *Arch Pediatr Adolesc Med*. 2002;156(6):607–614
 30. Cummins LH, Chan KK, Burns KM, Blume AW, Larimer M, Marlatt GA. Validity of the CRAFFT in American-Indian and Alaska-Native adolescents: screening for drug and alcohol risk. *J Stud Alcohol*. 2003;64(5):727–732
 31. Subramaniam M, Cheok C, Verma S, Wong J, Chong SA. Validity of a brief screening instrument: CRAFFT in a multiethnic Asian population. *Addict Behav*. 2010;35(12):1102–1104
 32. Mitchell SG, Kelly SM, Gryczynski J, et al. The CRAFFT cut-points and DSM-5 criteria for alcohol and other drugs: a reevaluation and reexamination. *Subst Abus*. 2014;35(4):376–380
 33. Orenstein A, Davis RB, Wolfe H; CASPAR Alcohol and Drug Education Program. Comparing screening instruments. *J Alcohol Drug Educ*. 2001;40(3):119–131
 34. Clark DB, Chung T, Martin C. Alcohol use frequency as a screen for alcohol use disorders in adolescents. *Int J Adolesc Med Health*. 2006;18(1):181–187
 35. Miech RA, Johnston LD, O'Malley PM, Bachman JG, Schulenberg JE. *Monitoring the Future National Survey Results on Drug Use, 1975–2015. Vol. I: Secondary School Students*. Ann Arbor, MI: Institute for Social Research, The University of Michigan; 2016
 36. Babor TF, de la Fuente JR, Saunders J, Grant MA. *AUDIT: The Alcohol Use Disorders Identification Test: Guidelines for Use in Primary Health Care*. Geneva, Switzerland: World Health Organization; 1992
 37. Shaffer D, Fisher P, Lucas C, Comer J. *Diagnostic Interview Schedule for Children (DISC-IV) Scoring Manual*. New York, NY: NIMH DISC Editorial Board and Mark Davies; 2007
 38. Wasserman GA, McReynolds LS, Lucas CP, Fisher P, Santos L. The voice DISC-IV with incarcerated male youths: prevalence of disorder. *J Am Acad Child Adolesc Psychiatry*. 2002;41(3):314–321
 39. NIMH. NHANES data documentation National Institute of Mental Health Diagnostic Interview Schedule for Children (DISC) (YDQ): years of coverage: 1999–2004. Available at: www.cdc.gov/nchs/data/nhanes/limited_access/ydq.pdf. Accessed October 30, 2011
 40. Shaffer D, Lucas CP, Richters JE. *Diagnostic Assessment in Child and Adolescent Psychopathology*. New York, NY: Guilford Press; 1999
 41. Johnston LD, O'Malley PM, Bachman JG, Schulenberg JE. *Monitoring the Future National Survey Results on Drug Use, 1975–2012. Vol. I: Secondary School Students*. Ann Arbor, MI: Institute for Social Research, The University of Michigan; 2013
 42. Collett D. *Modelling Binary Data*. 2nd ed. Boca Raton, FL: CRC Press; 2003

43. Harris SK, Knight JRJ Jr, Van Hook S, et al. Adolescent substance use screening in primary care: validity of computer self-administered versus clinician-administered screening. *Subst Abuse*. 2016;37(1):197–203
44. Levy S, Weiss R, Sherritt L, et al. An electronic screen for triaging adolescent substance use by risk levels. *JAMA Pediatr*. 2014;168(9):822–828
45. D'Amico EJ, Tucker JS, Miles JNV, Ewing BA, Shih RA, Pedersen ER. Alcohol and marijuana use trajectories in a diverse longitudinal sample of adolescents: examining use patterns from age 11 to 17 years. *Addiction*. 2016;111(10):1825–1835
46. D'Amico EJ, Houck JM, Hunter SB, Miles JNV, Osilla KC, Ewing BA. Group motivational interviewing for adolescents: change talk and alcohol and marijuana outcomes. *J Consult Clin Psychol*. 2015;83(1):68–80
47. D'Amico EJ, Miles JNV, Stern SA, Meredith LS. Brief motivational interviewing for teens at risk of substance use consequences: a randomized pilot study in a primary care clinic. *J Subst Abuse Treat*. 2008;35(1):53–61
48. Jensen CD, Cushing CC, Aylward BS, Craig JT, Sorell DM, Steele RG. Effectiveness of motivational interviewing interventions for adolescent substance use behavior change: a meta-analytic review. *J Consult Clin Psychol*. 2011;79(4):433–440
49. Ozechowski TJ, Becker SJ, Hogue A. SBIRT-A: adapting SBIRT to maximize developmental fit for adolescents in primary care. *J Subst Abuse Treat*. 2016;62:28–37
50. Helitzer DL, Sussman AL, de Hernandez BU, Kong AS. The “ins” and “outs” of provider–parent communication: perspectives from adolescent primary care providers on challenges to forging alliances to reduce adolescent risk. *J Adolesc Health*. 2011;48(4):404–409
51. Committee on Adolescence American Academy of Pediatrics. Achieving quality health services for adolescents. *Pediatrics*. 2008;121(6):1263–1270
52. Lustig JL, Ozer EM, Adams SH, et al. Improving the delivery of adolescent clinical preventive services through skills-based training. *Pediatrics*. 2001;107(5):1100–1107
53. Buckelew SM, Adams SH, Irwin CE Jr, Gee S, Ozer EM. Increasing clinician self-efficacy for screening and counseling adolescents for risky health behaviors: results of an intervention. *J Adolesc Health*. 2008;43(2):198–200
54. Harris SK, Knight JR. Putting the screen in screening: technology-based alcohol screening and brief interventions in medical settings. *Alcohol Res*. 2014;36(1):63–79
55. Chan D. So why ask me? Are self-report data really that bad? In: Lance CE, Vandenberg RJ, eds. *Statistical and Methodological Myths and Urban Legends: Doctrine, Verity and Fable in the Organizational and Social Sciences*. New York, NY: Psychology Press; 2008:309–336

Screening in Primary Care: What Is the Best Way to Identify At-Risk Youth for Substance Use?

Elizabeth J. D'Amico, Layla Parast, Lisa S. Meredith, Brett A. Ewing, William G. Shadel and Bradley D. Stein

Pediatrics 2016;138;

DOI: 10.1542/peds.2016-1717 originally published online November 18, 2016;

Updated Information & Services	including high resolution figures, can be found at: http://pediatrics.aappublications.org/content/138/6/e20161717
References	This article cites 45 articles, 6 of which you can access for free at: http://pediatrics.aappublications.org/content/138/6/e20161717#BIBL
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Adolescent Health/Medicine http://www.aappublications.org/cgi/collection/adolescent_health_medicine_sub Substance Use http://www.aappublications.org/cgi/collection/substance_abuse_sub
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.aappublications.org/site/misc/Permissions.xhtml
Reprints	Information about ordering reprints can be found online: http://www.aappublications.org/site/misc/reprints.xhtml

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Screening in Primary Care: What Is the Best Way to Identify At-Risk Youth for Substance Use?

Elizabeth J. D'Amico, Layla Parast, Lisa S. Meredith, Brett A. Ewing, William G. Shadel and Bradley D. Stein

Pediatrics 2016;138;

DOI: 10.1542/peds.2016-1717 originally published online November 18, 2016;

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/138/6/e20161717>

Data Supplement at:

<http://pediatrics.aappublications.org/content/suppl/2016/11/16/peds.2016-1717.DCSupplemental>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2016 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

