

The Well-Appearing Newborn at Risk for Early-Onset Sepsis: We Can Do Better

James J. Cummings, MD, MS

A common question in the newborn nursery is when to do a sepsis evaluation. When one considers pathogens such as group B *Streptococcus*, the risk of invasive infection is higher in the newborn than at any other time of life.¹ Overall, the incidence of early-onset sepsis (EOS, defined as a positive blood or cerebrospinal fluid culture within the first 72 hours after birth) in infants born at term is estimated to be ~1 per 2000.² Although case fatality rates are <2%, the incidence of long-term neurologic sequelae can be as high as 50%.^{1,2}

A baseline risk of <1 per 1000 may be too low to justify routine evaluation for EOS, but perinatal factors can significantly increase that risk. Questions then arise: Whom to test, when to treat with antibiotics pending test results, when to perform a lumbar puncture, and how long to treat with antibiotics if cultures remain negative? In an effort to aid the clinician, guidelines^{3,4} and risk-based tools^{5,6} have been developed.

How does the newborn nursery provider use these available resources when managing the well-appearing term infant? The study by Mukhopadhyay et al⁷ in this issue of *Pediatrics* examines this question. In a Web-based survey within a newborn outcome research network of 81 nurseries across the United States, they found wide variation in how well-appearing newborns are assessed and managed for EOS. Although the majority of respondents used 1 of

the published guidelines or sepsis risk calculators, >25% relied on local protocols or individual provider discretion. Frequency of intervention varied from doing less to doing more than directed by published resources. Notably, 2 of the 81 responders reported that their site practice was to provide observation and routine newborn care to all well-appearing newborns, without consideration of perinatal sepsis risk factors.

In general, clinical practice guidelines are often not followed^{8,9}; for the healthy-appearing newborn with a low risk of EOS there may also be concern that our current guidelines and risk-based calculators result in too many unnecessary evaluations and antibiotic exposures in infants who do not have EOS.^{10,11} But what is more concerning in the study by Mukhopadhyay et al are the extremes of practice variation, from observation without regard to perinatal risk factors to intervention based solely on fetal tachycardia, neither of which is supported by any of the published guidelines or risk-based models.

The bugbear of chorioamnionitis also looms large in the findings of Mukhopadhyay et al. It was 1 of the 2 most common clinical risk factors that prompted evaluation and treatment; although this finding is consistent with currently published guidelines,^{3,4} neither the guidelines nor the survey provides rigorous criteria for the diagnosis of chorioamnionitis, leaving it to individual interpretation. For respondents who primarily

Pediatrics, The Bernard & Millie Duker Children's Hospital at Albany Medical Center, Albany, New York

Opinions expressed in these commentaries are those of the author and not necessarily those of the American Academy of Pediatrics or its Committees.

DOI: 10.1542/peds.2016-4211

Accepted for publication Dec 19, 2016

Address correspondence to James J. Cummings, MD, MS, Vice Chair and Professor of Pediatrics, The Bernard & Millie Duker Children's Hospital at Albany Medical Center, 43 New Scotland Ave, Mailstop 88, Albany, NY 12208. E-mail: cumminj1@mail.amc.edu

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2017 by the American Academy of Pediatrics

FINANCIAL DISCLOSURE: The author has indicated he has no financial relationships relevant to this article to disclose.

FUNDING: No external funding.

POTENTIAL CONFLICT OF INTEREST: The author has indicated he has no potential conflicts of interest to disclose.

COMPANION PAPER: A companion to this article can be found online at www.pediatrics.org/cgi/doi/10.1542/peds.2016-2845.

To cite: Cummings JJ. The Well-Appearing Newborn at Risk for Early-Onset Sepsis: We Can Do Better. *Pediatrics*. 2017;139(3):e20164211

used risk-based calculators, most considered an obstetrical diagnosis of chorioamnionitis an additional risk factor, despite the fact that those calculators were specifically designed to avoid such subjective considerations.⁶

Earlier this year, a panel of experts in maternal, fetal, and neonatal care grappled with the vagaries of the term "maternal chorioamnionitis". In their summary report they recommended a more refined definition of intrauterine inflammation or infection (so-called Triple I), gave criteria for suspected and confirmed Triple I, and emphasized that isolated maternal fever was insufficient to make a diagnosis of Triple I.¹² However, this report is unlikely to change the findings of Mukhopadhyay et al, because almost all cases the newborn provider will have to struggle with will fall into the suspected category, a situation for which the panel recommends that "care should be individualized." Confirmation of Triple I requires either abnormal amniotic fluid laboratories or placental pathology; amniocentesis is rarely done, and pathology results typically are not available for several days after birth, leaving the newborn provider with no clear direction when evaluating the newly born infant.

The study by Mukhopadhyay et al has the usual shortcomings of survey research. In particular, it is limited by its sample size and depth of inquiry. Still, if such variation was evident among a limited sample, even wider variation is possible from a larger study. In a limited survey

we also gain little insight into the barriers to guideline adherence or why providers choose an alternative approach.⁹ What the survey by Mukhopadhyay et al does tell us is that assessment and management of the well-appearing newborn at risk for EOS have significant room for improvement, perhaps through a better understanding of the relative risks of harm and benefit for each of our treatment options (observation alone, evaluation and observation, or empirical treatment), the value and limitations of laboratory tests, the barriers that prevent providers from appropriately using available guidelines and risk-based tools, and how our current guidelines can provide better guidance.

ABBREVIATION

EOS: early-onset sepsis

REFERENCES

1. Simonsen KA, Anderson-Berry AL, Delair SF, Davies HD. Early-onset neonatal sepsis. *Clin Microbiol Rev.* 2014;27(1):21–47
2. Weston EJ, Pondo T, Lewis MM, et al. The burden of invasive early-onset neonatal sepsis in the United States, 2005–2008. *Pediatr Infect Dis J.* 2011;30(11):937–941
3. Polin RA; Committee on Fetus and Newborn. Management of neonates with suspected or proven early-onset bacterial sepsis. *Pediatrics.* 2012;129(5):1006–1015
4. Verani J, McGee L, Schrag S, Division of Bacterial Diseases, National Center for Immunization and Respiratory Diseases, Centers for Disease and Prevention (CDC). Prevention of perinatal group B streptococcal

disease: revised guidelines from CDC, 2010. *MMWR Morb Mortal Wkly Rep.* 2010;59(RR10):1–32

5. Escobar GJ, Puopolo KM, Wi S, et al. Stratification of risk of early-onset sepsis in newborns ≥ 34 weeks' gestation. *Pediatrics.* 2014;133(1):30–36
6. Puopolo KM, Draper D, Wi S, et al. Estimating the probability of neonatal early-onset infection on the basis of maternal risk factors. *Pediatrics.* 2011;128(5). Available at: www.pediatrics.org/cgi/content/full/113/2/e1155
7. Mukhopadhyay S, Taylor JA, Von Kohorn I, et al. Variation in sepsis evaluation across a national network of nurseries. *Pediatrics.* 2017;139(3):e20162845
8. Cabana MD, Rand CS, Powe NR, et al. Why don't physicians follow clinical practice guidelines? A framework for improvement. *JAMA.* 1999;282(15):1458–1465
9. Fischer F, Lange K, Klose K, Greiner W, Kraemer A. Barriers and strategies in guideline implementation: a scoping review. *Healthcare (Basel).* 2016;4(3):36
10. Kiser C, Nawab U, McKenna K, Aghai ZH. Role of guidelines on length of therapy in chorioamnionitis and neonatal sepsis. *Pediatrics.* 2014;133(6):992–998
11. Mukhopadhyay S, Dukhovny D, Mao W, Eichenwald EC, Puopolo KM. 2010 perinatal GBS prevention guideline and resource utilization. *Pediatrics.* 2014;133(2):196–203
12. Higgins RD, Saade G, Polin RA, et al; Chorioamnionitis Workshop Participants. Evaluation and management of women and newborns with a maternal diagnosis of chorioamnionitis: summary of a workshop. *Obstet Gynecol.* 2016;127(3):426–436

The Well-Appearing Newborn at Risk for Early-Onset Sepsis: We Can Do Better

James J. Cummings

Pediatrics 2017;139;

DOI: 10.1542/peds.2016-4211 originally published online February 8, 2017;

| | |
|---|--|
| Updated Information & Services | including high resolution figures, can be found at: http://pediatrics.aappublications.org/content/139/3/e20164211 |
| References | This article cites 11 articles, 6 of which you can access for free at: http://pediatrics.aappublications.org/content/139/3/e20164211#BIBL |
| Subspecialty Collections | This article, along with others on similar topics, appears in the following collection(s): Fetus/Newborn Infant http://www.aappublications.org/cgi/collection/fetus:newborn_infant_sub |
| Permissions & Licensing | Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.aappublications.org/site/misc/Permissions.xhtml |
| Reprints | Information about ordering reprints can be found online: http://www.aappublications.org/site/misc/reprints.xhtml |

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®



PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

The Well-Appearing Newborn at Risk for Early-Onset Sepsis: We Can Do Better

James J. Cummings

Pediatrics 2017;139;

DOI: 10.1542/peds.2016-4211 originally published online February 8, 2017;

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/139/3/e20164211>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 345 Park Avenue, Itasca, Illinois, 60143. Copyright © 2017 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®

