

# Effectiveness of Tympanostomy Tubes for Otitis Media: A Meta-analysis

Dale W. Steele, MD, MS,<sup>a,b,c,d</sup> Gaelen P. Adam, MLIS,<sup>a</sup> Mengyang Di, MD, PhD,<sup>a</sup> Christopher H. Halladay, ScM,<sup>a</sup> Ethan M. Balk, MD, MPH,<sup>a,b</sup> Thomas A. Trikalinos, MD, PhD<sup>a,b</sup>

abstract

**CONTEXT:** Tympanostomy tube placement is the most common ambulatory surgery performed on children in the United States.

**OBJECTIVES:** The goal of this study was to synthesize evidence for the effectiveness of tympanostomy tubes in children with chronic otitis media with effusion and recurrent acute otitis media.

**DATA SOURCES:** Searches were conducted in Medline, the Cochrane Central Trials Registry and Cochrane Database of Systematic Reviews, Embase, and the Cumulative Index to Nursing and Allied Health Literature.

**STUDY SELECTION:** Abstracts and full-text articles were independently screened by 2 investigators.

**DATA EXTRACTION:** A total of 147 articles were included. When feasible, random effects network meta-analyses were performed.

**RESULTS:** Children with chronic otitis media with effusion treated with tympanostomy tubes compared with watchful waiting had a net decrease in mean hearing threshold of 9.1 dB (95% credible interval: -14.0 to -3.4) at 1 to 3 months and 0.0 (95% credible interval: -4.0 to 3.4) by 12 to 24 months. Children with recurrent acute otitis media may have fewer episodes after placement of tympanostomy tubes. Associated adverse events are poorly defined and reported.

**LIMITATIONS:** Sparse evidence is available, applicable only to otherwise healthy children.

**CONCLUSIONS:** Tympanostomy tubes improve hearing at 1 to 3 months compared with watchful waiting, with no evidence of benefit by 12 to 24 months. Children with recurrent acute otitis media may have fewer episodes after tympanostomy tube placement, but the evidence base is severely limited. The benefits of tympanostomy tubes must be weighed against a variety of associated adverse events.



<sup>a</sup>Evidence-based Practice Center, Center for Evidence Synthesis in Health, <sup>b</sup>Department of Health Services, Policy and Practice, School of Public Health, <sup>c</sup>Department of Emergency Medicine, Section of Pediatrics—Hasbro Children's Hospital, and <sup>d</sup>Department of Pediatrics, Alpert Medical School, Brown University, Providence, Rhode Island

Dr Steele conceptualized and designed the review, designed data collection instruments and supervised data collection, performed data analysis and interpretation of data, and drafted and revised the manuscript; Ms Adam performed the literature search, participated in data collection, performed analysis and interpretation of data, and critically reviewed the manuscript; Dr Di and Mr Halliday participated in data collection, analysis, and interpretation of data, and reviewed and revised the manuscript; Dr Balk assisted in conceptualization and design, reviewed data collection, collected data, and critically reviewed the manuscript; and Dr Trikalinos oversaw protocol conception and design and analysis and interpretation of data, performed supplemental analysis, and critically reviewed and revised the manuscript. All authors approved the final manuscript as submitted.

The authors of this report are responsible for its content. Statements in the report should not be construed as endorsement by the Agency for Healthcare Research and Quality or the US Department of Health and Human Services.

**To cite:** Steele DW, Adam GP, Di M, et al. Effectiveness of Tympanostomy Tubes for Otitis Media: A Meta-analysis. *Pediatrics*. 2017;139(6):e20170125

Myringotomy with tympanostomy tube placement is the most common ambulatory surgery performed on children in the United States,<sup>1</sup> with 667 000 children aged <15 years undergoing tympanostomy tube placement in 2006.<sup>2</sup> The effectiveness of tympanostomy tubes for chronic otitis media with effusion (OME) and recurrent acute otitis media (AOM) is likely influenced by the many factors that affect the prognosis for middle ear disease in children. These factors include current age, age at first diagnosis, frequency of respiratory tract infections, and day care exposure.<sup>3</sup>

The American Academy of Otolaryngology–Head and Neck Surgery clinical practice guideline recommends that clinicians should offer bilateral tympanostomy tube insertion to children with bilateral OME for  $\geq 3$  months and who have documented hearing difficulties. They may offer tympanostomy tubes to children with unilateral or bilateral OME with symptoms that are likely attributable to OME which include, but are not limited to, vestibular problems, poor school performance, behavioral problems, ear discomfort, or reduced quality of life.<sup>4</sup>

The American Academy of Pediatrics' clinical practice guideline for diagnosis and management of AOM states that clinicians may offer tympanostomy tubes for recurrent AOM (3 episodes in 6 months or 4 episodes in 1 year with 1 episode in the preceding 6 months).<sup>5</sup> The American Academy of Otolaryngology–Head and Neck Surgery clinical practice guideline further recommends that clinicians not perform tympanostomy tube insertion when middle ear effusion is not present at the time of assessment for tube placement; they argue that the presence of effusion serves as both a marker for the accuracy of the diagnosis of AOM and an indicator of underlying Eustachian tube dysfunction with decreased ability

to clear middle ear fluid after an episode of AOM.<sup>4</sup>

In the present systematic review, we synthesized the available evidence regarding the effectiveness of tympanostomy tubes (with or without adenoidectomy) compared with watchful waiting in children with chronic OME and children with recurrent AOM. We also summarized the frequency of adverse events associated with tympanostomy tubes.

This review is derived from an Agency for Healthcare Research and Quality–commissioned comparative effectiveness review (Tympanostomy Tubes in Children With Otitis Media) conducted by the Brown Evidence-based Practice Center. The full review and review protocol (PROSPERO registry number: CRD42015029623) are available at <http://www.effectivehealthcare.ahrq.gov>.

## METHODS

The approaches outlined in the Agency for Healthcare Research and Quality's Methods Guide for Comparative Effectiveness Reviews were followed.<sup>6</sup>

### Search Strategy and Study Selection


This study evaluated published, peer-reviewed studies in which at least 1 arm included children receiving tympanostomy tubes; conference abstracts were excluded. We included randomized controlled trials (RCTs) and nonrandomized comparative studies (NRCS), prospective and retrospective, in which treatment with tympanostomy tubes was assigned on a per-patient basis. Studies with per-ear assignment were excluded (eg, tympanostomy tubes placed by design in 1 ear only). For adverse events, prospective surgical studies enrolling at least 50 subjects (including arms of RCTs or NRCS with  $\geq 50$  patients) and population-based retrospective single-group studies (registry

studies) with at least 1000 subjects were included.

Literature searches without language restrictions were conducted in Medline, the Cochrane Central Trials Registry and Cochrane Database of Systematic Reviews, Embase, and the Cumulative Index to Nursing and Allied Health Literature (from inception) to identify primary research studies meeting the study criteria. Citations found by literature searches were independently screened by 2 researchers, using the open-source, online software *abstrackr* (<http://abstrackr.cebml.brown.edu/>).<sup>7</sup> Conflicts were resolved by discussion until a group consensus was reached.

### Data Extraction and Analysis

Each study was extracted by 1 methodologist; the extractions were reviewed and confirmed by at least 1 other methodologist. We conducted quantitative analysis for outcomes with at least 5 studies which report results that could be combined in a meta-analysis. Bayesian network meta-analysis was performed by using the R *gemtc* package.<sup>8</sup> Estimation was performed with Markov chain Monte Carlo methods via the JAGS software,<sup>9</sup> using initial values drawn randomly from the marginal distributions of the priors of respective parameters. We fit 4 Markov chain Monte Carlo chains. After a burn-in of 5000 iterations, we monitored the convergence of random effects means and variances automatically by checking every 10 000 iterations whether the Gelman-Rubin diagnostic was  $< 1.05$  with 95% probability for all monitored parameters. After convergence was reached, an additional 10 000 iterations were run. All models converged within 10 000 iterations. Model fit was assessed by comparing the posterior mean of the residual deviance versus the number of data points. The ratio of residual deviance to number of



**FIGURE 1**  
Literature flow diagram. CINAHL, Cumulative Index to Nursing and Allied Health Literature.

data points ranged from 0.97 to 1.06, suggesting an adequate model fit.

Network meta-analysis is an extension of pairwise meta-analyses that simultaneously combines direct (when interventions are compared head-to-head) and indirect (when interventions are compared through other reference interventions) evidence. Combining the direct and indirect evidence not only improves the precision of estimates but also provides estimates for all pairwise comparisons, including those missing from the direct evidence.<sup>10</sup> Statistical heterogeneity was explored qualitatively. Because of the relatively small number of studies, and the little variability in characteristics, meta-regression analyses were not performed.

### Assessment of Study Risk of Bias and Strength of Evidence

The methodologic quality of each study was assessed on the basis of predefined criteria. For RCTs, the Cochrane risk of bias tool was

used.<sup>11</sup> The strength of evidence was graded according to the Agency for Healthcare Research and Quality's methods guide on assessing the strength of evidence.<sup>12</sup>

## RESULTS

Figure 1 displays the results of the literature search and selection process.

### Effectiveness of Tympanostomy Tubes in Children With OME

A total of 54 publications were identified. Of these, 29 articles reported the results of 16 RCTs.<sup>13–41</sup> Twenty-four publications reported the results of 24 NRCS that assessed the effectiveness of tympanostomy tubes in pediatric patients with chronic middle ear effusion.<sup>42–65</sup> These studies evaluated multiple interventions (tympanostomy tubes, tympanostomy tubes with adenoideotomy, myringotomy with adenoideotomy, myringotomy alone,

adenoideotomy alone, oral antibiotic prophylaxis, and watchful waiting).

Hearing thresholds were measured in 16 RCTs. In 10 of these RCTs, mean hearing thresholds were reported according to study arm at various time points. For the network meta-analysis of these RCTs, we classified hearing thresholds obtained at 1 to 3 months as “early”; hearing thresholds obtained between 12 and 24 months were classified as “late.” Not all studies had interventions at both early and late time points. Thus, the network of comparators differs for early and late comparisons.

Figure 2 illustrates the effectiveness of various interventions at 1 to 3 months compared with watchful waiting. Mean hearing thresholds improved (ie, decreased) by an average of 9.1 dB after insertion of the tympanostomy tubes and by 10 dB after tympanostomy tube insertion with adenoideotomy. As shown in Table 1, the strategies with the highest probability of being among the 3 most effective interventions with respect to early improvements in hearing thresholds were tympanostomy tubes, tympanostomy tubes with adenoideotomy, and myringotomy with adenoideotomy.

Five RCTs reported hearing thresholds at 12 to 24 months. As shown in Fig 3, by 12 to 24 months, the mean difference in hearing thresholds for tympanostomy tubes alone, compared with watchful waiting, was 0 dB (95% credible interval: –4 to 3). As can be seen in Table 2, tympanostomy tube insertion with adenoideotomy and myringotomy with adenoideotomy were the 2 most effective strategies with respect to late hearing thresholds. Tympanostomy tubes alone, antibiotic prophylaxis, and watchful waiting were among the 3 least effective strategies.

Eight studies (5 RCTs, 3 NRCS, and 1 that combined both designs) in 12


articles reported on 119 quality of life and patient-centered outcomes (cognitive, language, and behavioral) in 1665 children over multiple time points and arms. In general, the results were not significant (only 14 of 119 had significant results), and they varied in magnitude and direction, even across subscales of the same test.

### Effectiveness of Tympanostomy Tubes in Children With Recurrent AOM

We identified 8 publications, reporting results from 7 RCTs<sup>66–72</sup> and 2 NRCS<sup>69,73</sup> that reported outcomes for children with recurrent AOM.

Three RCTs compared tympanostomy tubes with placebo or no treatment. The first trial reported that 3 of 20 children in the placebo group had no further episodes of AOM, compared with 12 of 22 in the tympanostomy tube group ( $P = .01$ ), with an attack rate of 2.0 infections per child in the placebo group, compared with 0.86 in the tympanostomy tube group ( $P = .006$ ). The authors reported a post hoc subgroup comparison of treatment failure ( $\geq 2$  episodes in 3 months) rates. Children without middle ear effusion at study entry had significantly fewer bouts of AOM ( $P < .05$ ) and lower attack rates than children with middle ear effusion. However, in a logistic model of treatment failure, adjusted for presence of middle ear effusion, the treatment according to subgroup interaction term is nonsignificant ( $P = .69$ ). This analysis relies on a small sample ( $n = 42$ ) and is therefore underpowered, but it provides no evidence that the presence or absence of middle ear effusion at study entry influenced the efficacy of tympanostomy tubes.<sup>66</sup>

A second trial reported that the rate of new episodes per arm was 1.08 in the placebo group versus 1.02 in the tympanostomy tube group ( $P = .25$ ).




**FIGURE 2**

Early (1–3 months) decrease (improvement) in mean hearing thresholds compared with watchful waiting. CrI, credible interval; TT, tympanostomy tubes.

**TABLE 1** Probabilities That an Intervention Is Among the 3 Most Effective With Respect to Early Hearing Thresholds

Intervention	Probability (%) of Being Among the 3 Most Effective Interventions	Probability (%) of Being Among the 3 Least Effective Interventions
TT	97	3
TT + adenoidectomy	96	4
Myringotomy	8	92
Myringotomy + adenoidectomy	91	9
Antibiotic prophylaxis	6	94
Watchful waiting	1	99

TT, tympanostomy tubes.



**FIGURE 3**

Late (12–24 months) decrease (improvement) in mean hearing thresholds compared with watchful waiting. CrI, credible interval; TT, tympanostomy tubes.

In the placebo group, 40% had no further episodes of AOM, compared with 35% in the tympanostomy tube group. However, tympanostomy tube placement significantly decreased the percentage of time with AOM (6.6%) compared with placebo (15.0%;  $P < .001$ ).<sup>67</sup>

The third, most recent trial reported failure rates (defined as at least 2 episodes of AOM in 2 months, 3 in 6 months, or persistent effusion lasting at least 2 months), percentage of children with no recurrent AOM, cumulative number of AOM episodes, and 1-year incidence rates. There

was an absolute difference in the proportion of failures of  $-13\%$  (95% confidence interval:  $-25$  to  $-1$ ) between the tympanostomy tube and control groups, favoring tympanostomy tubes. The 1-year incidence rate (infections per child per year) was 0.55 (95% confidence interval: 0.93 to 0.17) lower in the tympanostomy tube group compared with the control group.<sup>71</sup>

We were unable to provide pooled results due to the small number of studies, multiple interventions, and heterogeneity in reported outcomes. The limited available evidence suggests that tympanostomy tube placement decreases the risk of recurrent AOM. Aside from the first study,<sup>66</sup> we found no direct evidence to evaluate whether the presence of middle ear effusion at the time of surgical evaluation modifies the effectiveness of tympanostomy tube placement for recurrent AOM because the other 2 studies specifically excluded patients with current middle ear effusion.<sup>67,71</sup>

Three RCTs evaluated tympanostomy tubes alone versus adenoidectomy and tympanostomy tubes. Of these trials, none reported a difference in recurrent episodes of AOM.<sup>69-71</sup>

Risk of bias across outcomes ranged from moderate to high.

### Adverse Events Associated With Tympanostomy Tubes

We extracted data on the occurrence of 11 adverse events from 85 cohorts and from the tympanostomy tube arms of included RCTs and NRCS. The number of publications reporting each event and the median (with 25th and 75th percentiles) percentage of patients and ears are summarized in Supplemental Table 4; references for these adverse events are also supplied.<sup>13,19,21,30,39,52,67,71,74-158</sup>

In general, the study-specific definitions of adverse events were poorly reported and/or highly

**TABLE 2** Probabilities That an Intervention Is Among the 2 Most Effective With Respect to Late Hearing Thresholds

Intervention	Probability (%) of Being Among the 2 Most Effective Interventions	Probability (%) of Being Among the 3 Least Effective Interventions
TT	5	95
TT + adenoidectomy	92	8
Myringotomy + adenoidectomy	88	12
Antibiotic prophylaxis	10	90
Watchful waiting	4	96

TT, tympanostomy tubes.

variable between studies. Not all cohorts followed up all patients until extrusion of the tube, nor was follow-up complete in all studies. Several adverse event categories (eg, otorrhea, premature extrusion, myringosclerosis) have very wide interquartile ranges, likely due to highly variable definitions. Other adverse events, such as hearing loss and cholesteatoma, are likely confounded by the severity of preexisting and ongoing middle ear disease.

The main conclusions and interpretations, including strength of evidence assessments based on our meta-analysis, are summarized in Supplemental Table 3.

## DISCUSSION

Tympanostomy tube placement (compared with watchful waiting) in children with chronic middle ear effusion, results in improved average hearing thresholds at 1 to 3 months after surgery (a period when the majority of tubes are functioning). Mean hearing thresholds after tube placement with or without adenoidectomy improved by  $\sim 10$  dB when assessed at 1 to 3 months.

By 1 to 2 years after surgery, when most tubes have extruded, hearing thresholds are no longer different, likely reflecting the usually favorable natural history of spontaneous resolution of middle ear effusion in most children in both groups. There is limited evidence regarding quality of life outcomes, but neither of the 2

studies that evaluated parental stress and health-related quality of life found significant improvements in surgically treated children compared with watchful waiting.

Tympanostomy tubes did not consistently improve cognition, behavior, or quality of life, but low statistical power prevents any definitive conclusions, and the results apply to otherwise healthy children with no baseline disorders or delays in language, cognition, or behavior. With the exception of a few NRCS, comparative trials systematically exclude patients with cleft palate and Down syndrome, thus limiting the applicability of the evidence for these and other similar subgroups who experience a higher burden of middle ear disease. Similarly, patients at increased risk of developmental or behavioral sequelae from middle ear disease have not been included (or at least identified) in trials to date.

Given the sparse data and limitations inherent to the synthesis of aggregate data, we were unable to perform an analysis of factors which would predict those children more likely to benefit from tympanostomy tubes for chronic middle ear effusion. Additional insight was provided by an individual patient data meta-analysis that focused on interactions between treatment and baseline characteristics.<sup>159</sup> The meta-analysis found significant interactions between day care attendance in children aged  $\leq 3$  years, and in children  $>4$  years of age with a hearing level of  $\geq 25$  dB

in both ears, and concluded that tympanostomy tubes might be most effective in young children attending day care, or in older children with persistent hearing impairments at least 12 weeks. However, average hearing level at baseline did not obviously modify effectiveness. Our meta-analysis of hearing levels used average pure tone hearing levels (typically reported as an average over frequencies of 500, 1000, 2000, and 4000 Hz). This single outcome measure is likely insufficient to fully elucidate the complex relationships between hearing, speech perception, and development in children.

Our network meta-analysis suggests a trend toward improved hearing thresholds in children undergoing adenoidectomy, but credible intervals are wide and include the null effect. An individual patient data meta-analysis of children with persistent OME concluded that adenoidectomy is most beneficial in children aged  $\geq 4$  years, with no significant benefit of adenoidectomy in children  $< 4$  years old.<sup>160</sup> This evidence is reflected in a recently updated clinical practice guideline for OME, which promotes tympanostomy tubes as the primary surgical intervention for younger children and reserves adenoidectomy for children  $\geq 4$  years or those with a distinct indication for the procedure other than OME.<sup>161</sup>

In children with recurrent AOM, the limited available evidence suggests that tympanostomy tubes decrease the number of additional episodes and the overall number of episodes of recurrent AOM. The degree to which the presence of middle ear fluid at the time of evaluation for surgery will affect the effectiveness of the tympanostomy tubes is unclear.

Three RCTs consistently found no difference in recurrent episodes of AOM in children who underwent tympanostomy tube placement with adenoidectomy compared with tympanostomy tube placement alone.<sup>69–71</sup>

Our systematic review of adverse events associated with tympanostomy tube placement provides a descriptive summary of the observed frequency in published cohorts. However, the study-specific definitions of adverse events are highly variable and poorly reported. Some adverse events, such as hearing loss and cholesteatoma, are likely confounded by the severity of preexisting and ongoing middle ear disease.

### Limitations

The available evidence base is composed of studies that evaluate multiple interventions. Several of these (eg, myringotomy alone and oral antibiotic prophylaxis) are rarely used or not recommended<sup>5</sup> in current practice. We used indirect evidence from a network meta-analysis to augment the direct evidence relating to the comparisons of current interest. The key assumption of the network meta-analysis is that of consistency of direct and indirect effects. Consistency is likely to hold when the distribution of effect modifiers is similar across trials. If this assumption is violated, there may be inconsistency between the direct evidence and indirect evidence of treatment comparisons.

Reporting of possible sociodemographic risk factors is sparse and inconsistent, which limits our ability to draw conclusions about which of these factors might influence the relative effectiveness of tympanostomy tubes.

Current recommended indications for tympanostomy tube placement largely reflect the inclusion criteria used in existing clinical trials. Given the usually favorable natural history of middle ear effusion, well-validated prognostic models are urgently needed to stratify the risk of individual children in terms of their risk for persistence of middle ear effusion and/or recurrent AOM.

Assessing the effectiveness of tympanostomy tubes in children with recurrent AOM is particularly challenging because an episode of AOM in control children (with an intact tympanic membrane) results in otalgia and inflammatory changes, whereas children with a functioning tympanostomy tube may present with varying degrees of otorrhea. Outcomes that rely on simple counts or rates of otorrhea fail to account for the variable character of otorrhea with respect to duration, character, and patient impact. For example, otorrhea may be transient (of little to no concern), recurrent (of more concern but usually readily managed), or chronic (of considerable concern and difficult to manage).

### Future Research Needs

Pragmatic trials are needed, particularly in children with recurrent AOM, but also in children with chronic OME or some combination of both. Because tympanostomy tubes are no longer effective after extrusion, future trials should record per-ear and per-patient outcomes conditional on whether the tympanostomy tube has been extruded. Future studies should also conduct appropriate analyses to estimate the causal effects of tympanostomy tubes among children who still have the tubes in place. Future trials would benefit from standardization of disease definitions and consistent definition of core outcomes and adverse events.

Exploring treatment effect heterogeneity (ie, differential effects of interventions in populations at different risk levels for outcomes of interest) should be a priority. There is particular need for randomized studies evaluating tympanostomy tubes in higher risk groups, such as patients with cleft palate, Down syndrome, and children with neurodevelopmental disorders.

## CONCLUSIONS

Overall, the evidence suggests that tympanostomy tubes in children with persistent middle-ear effusion result in short-term improvements in hearing compared with watchful waiting. However, there is no evidence of a sustained benefit.

Our network meta-analysis of hearing thresholds suggests the possibility of a more sustained improvement in hearing thresholds in at least some children who undergo adenoidectomy and tympanostomy tube placement. A nuanced understanding of which children may benefit from adenoidectomy is limited by the small evidence base and our use of aggregate data.

The evidence suggests that treatment with tympanostomy tubes did not

improve cognition, behavior, or quality of life. However, the evidence is sparse and prevents any definitive conclusions. The results apply to otherwise healthy children without baseline disorders or delays in language or cognition. They provide little guidance for the treatment of children who may be at increased risk for speech, language, or learning problems because of baseline sensory, physical, cognitive, or behavioral factors.

Children with recurrent AOM may have fewer episodes after tympanostomy tube placement, but the evidence base is severely limited. It is unclear whether quality of life outcomes are improved. The benefits of tympanostomy tubes must be weighed against a variety of adverse

events associated with—although not necessarily caused by—this treatment.

## ACKNOWLEDGMENTS

Special thanks to Joseph Lau for his advice and guidance; Ian Pan and Nathan Coopersmith for assistance with screening and extraction; and Jenni Quiroz and Jenna Legault and for their assistance throughout the systematic review process.

## ABBREVIATIONS

AOM: acute otitis media  
NRCS: nonrandomized comparative studies  
OME: otitis media with effusion  
RCT: randomized controlled trial

**DOI:** <https://doi.org/10.1542/peds.2017-0125>

Accepted for publication Mar 15, 2017

Address correspondence to Dale W. Steele, MD, MS, Center for Evidence Synthesis in Health, Box G-S121-8, Brown University, Providence, RI 02912. E-mail: dale\_steele@brown.edu

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2017 by the American Academy of Pediatrics

**FINANCIAL DISCLOSURE:** All authors received funding for this project under Contract HHS290-2015-00002-I from the Agency for Healthcare Research and Quality, US Department of Health and Human Services.

**FUNDING:** Supported by the Agency for Healthcare Research and Quality (contract: HHS290-2015-00002-I).

**POTENTIAL CONFLICT OF INTEREST:** The authors have indicated they have no potential conflicts of interest to disclose.

## REFERENCES

1. Rettig E, Tunkel DE. Contemporary concepts in management of acute otitis media in children. *Otolaryngol Clin North Am*. 2014;47(5):651–672
2. Cullen KA, Hall MJ, Golosinskiy A. Ambulatory surgery in the United States, 2006. *Natl Health Stat Rep*. 2009;(11):1–25
3. Hoffman HJ, Daly KA, Bainbridge KE, et al. Panel 1: epidemiology, natural history, and risk factors. *Otolaryngol Head Neck Surg*. 2013;148(suppl 4):E1–E25
4. Rosenfeld RM, Schwartz SR, Pynnonen MA, et al. Clinical practice guideline: tympanostomy tubes in children. *Otolaryngol Head Neck Surg*. 2013;149(suppl 1):S1–S35
5. Lieberthal AS, Carroll AE, Chonmaitree T, et al. The diagnosis and management of acute otitis media. *Pediatrics*. 2013;131(3). Available at: [www.pediatrics.org/cgi/content/full/131/3/e964](http://www.pediatrics.org/cgi/content/full/131/3/e964)
6. Agency for Healthcare Research and Quality. *Methods Guide for Effectiveness and Comparative Effectiveness Reviews*. Rockville, MD: Agency for Healthcare Research and Quality; 2014
7. Wallace BC, Small K, Brodley CE, Lau J, Trikalinos TA. Deploying an interactive machine learning system in an evidence-based practice center: abstract. In: *Proceedings of the 2nd ACM SIGHIT International Health Informatics Symposium*. Miami, FL: ACM; 2012:819–824
8. van Valkenhoef G, Lu G, de Brock B, Hillege H, Ades AE, Welton NJ. Automating network meta-analysis. *Res Synth Methods*. 2012;3(4):285–299
9. Plummer M. JAGS: a program for analysis of Bayesian graphical models using Gibbs sampling. In: *Proceedings of the 3rd International Workshop on Distributed Statistical Computing (DSC)*; March 20–22, 2003; Vienna, Austria
10. Dias S, Sutton AJ, Ades AE, Welton NJ. Evidence synthesis for decision making 2: a generalized linear modeling framework for pairwise and network meta-analysis of randomized

- controlled trials. *Med Decis Making*. 2013;33(5):607–617
11. Higgins JP, Altman DG, Gøtzsche PC, et al; Cochrane Bias Methods Group; Cochrane Statistical Methods Group. The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *BMJ*. 2011;343:d5928
  12. Berkman ND, Lohr KN, Ansari M, et al; AHRQ Methods for Effective Health Care. *Grading the Strength of a Body of Evidence When Assessing Health Care Interventions for the Effective Health Care Program of the Agency for Healthcare Research and Quality: An Update. Methods Guide for Effectiveness and Comparative Effectiveness Reviews*. Rockville, MD: Agency for Healthcare Research and Quality; 2008
  13. Bernard PA, Stenstrom RJ, Feldman W, Durieux-Smith A. Randomized, controlled trial comparing long-term sulfonamide therapy to ventilation tubes for otitis media with effusion. *Pediatrics*. 1991;88(2):215–222
  14. Casselbrant ML, Mandel EM, Rockette HE, Kurs-Lasky M, Fall PA, Bluestone CD. Adenoidectomy for otitis media with effusion in 2-3-year-old children. *Int J Pediatr Otorhinolaryngol*. 2009;73(12):1718–1724
  15. Chaudhuri GR, Bandyopadhyay SN, Basu SK. Role of grommet in otitis media with effusion: a necessity or nuisance? A comparative study. *Indian J Otolaryngol Head Neck Surg*. 2006;58(3):271–273
  16. D'Eredità R, Shah UK. Contact diode laser myringotomy for medium-duration middle ear ventilation in children. *Int J Pediatr Otorhinolaryngol*. 2006;70(6):1077–1080
  17. Hall AJ, Maw AR, Steer CD. Developmental outcomes in early compared with delayed surgery for glue ear up to age 7 years: a randomised controlled trial. *Clin Otolaryngol*. 2009;34(1):12–20
  18. Hartman M, Rovers MM, Ingels K, Zielhuis GA, Severens JL, van der Wilt GJ. Economic evaluation of ventilation tubes in otitis media with effusion. *Arch Otolaryngol Head Neck Surg*. 2001;127(12):1471–1476
  19. Ingels K, Rovers MM, van der Wilt GJ, Zielhuis GA. Ventilation tubes in infants increase the risk of otorrhoea and antibiotic usage. *B-ENT*. 2005;1(4):173–176
  20. Johnston LC, Feldman HM, Paradise JL, et al. Tympanic membrane abnormalities and hearing levels at the ages of 5 and 6 years in relation to persistent otitis media and tympanostomy tube insertion in the first 3 years of life: a prospective study incorporating a randomized clinical trial. *Pediatrics*. 2004;114(1). Available at: [www.pediatrics.org/cgi/content/full/114/1/e58](http://www.pediatrics.org/cgi/content/full/114/1/e58)
  21. Mandel EM, Rockette HE, Bluestone CD, Paradise JL, Nozza RJ. Myringotomy with and without tympanostomy tubes for chronic otitis media with effusion. *Arch Otolaryngol Head Neck Surg*. 1989;115(10):1217–1224
  22. Mandel EM, Rockette HE, Bluestone CD, Paradise JL, Nozza RJ. Efficacy of myringotomy with and without tympanostomy tubes for chronic otitis media with effusion. *Pediatr Infect Dis J*. 1992;11(4):270–277
  23. Maw R, Wilks J, Harvey I, Peters TJ, Golding J. Early surgery compared with watchful waiting for glue ear and effect on language development in preschool children: a randomised trial. *Lancet*. 1999;353(9157):960–963
  24. MRC Multicentre Otitis Media Study Group. The role of ventilation tube status in the hearing levels in children managed for bilateral persistent otitis media with effusion. *Clin Otolaryngol Allied Sci*. 2003;28(2):146–153
  25. MRC Multicentre Otitis Media Study Group. Speech reception in noise: an indicator of benefit from otitis media with effusion surgery. *Clin Otolaryngol Allied Sci*. 2004;29(5):497–504
  26. MRC Multicentre Otitis Media Study Group. Adjuvant adenoidectomy in persistent bilateral otitis media with effusion: hearing and revision surgery outcomes through 2 years in the TARGET randomised trial. *Clin Otolaryngol*. 2012;37(2):107–116
  27. Nguyen LH, Manoukian JJ, Yoskovitch A, Al-Sebeih KH. Adenoidectomy: selection criteria for surgical cases of otitis media. *Laryngoscope*. 2004;114(5):863–866
  28. Paradise JL, Campbell TF, Dollaghan CA, et al. Developmental outcomes after early or delayed insertion of tympanostomy tubes. *N Engl J Med*. 2005;353(6):576–586
  29. Paradise JL, Dollaghan CA, Campbell TF, et al. Otitis media and tympanostomy tube insertion during the first three years of life: developmental outcomes at the age of four years. *Pediatrics*. 2003;112(2):265–277
  30. Paradise JL, Feldman HM, Campbell TF, et al. Effect of early or delayed insertion of tympanostomy tubes for persistent otitis media on developmental outcomes at the age of three years. *N Engl J Med*. 2001;344(16):1179–1187
  31. Paradise JL, Feldman HM, Campbell TF, et al. Early versus delayed insertion of tympanostomy tubes for persistent otitis media: developmental outcomes at the age of three years in relation to prerandomization illness patterns and hearing levels. *Pediatr Infect Dis J*. 2003;22(4):309–314
  32. Paradise JL, Feldman HM, Campbell TF, et al. Tympanostomy tubes and developmental outcomes at 9 to 11 years of age. *N Engl J Med*. 2007;356(3):248–261
  33. Popova D, Varbanova S, Popov TM. Comparison between myringotomy and tympanostomy tubes in combination with adenoidectomy in 3-7-year-old children with otitis media with effusion. *Int J Pediatr Otorhinolaryngol*. 2010;74(7):777–780
  34. Rach GH, Zielhuis GA, van Baarle PW, van den Broek P. The effect of treatment with ventilating tubes on language development in preschool children with otitis media with effusion. *Clin Otolaryngol Allied Sci*. 1991;16(2):128–132
  35. Rovers MM, Krabbe PF, Straatman H, Ingels K, Zielhuis GA, van der Wilt GJ. Randomised controlled trial of the effect of ventilation tubes (grommets) on quality of life at age 1-2 years. *Arch Dis Child*. 2001;84(1):45–49
  36. Rovers MM, Straatman H, Ingels K, van der Wilt GJ, van den Broek P, Zielhuis GA. The effect of ventilation tubes on language development in infants with otitis media with effusion: a randomized trial. *Pediatrics*.

- 2000;106(3). Available at: [www.pediatrics.org/cgi/content/full/106/3/e42](http://www.pediatrics.org/cgi/content/full/106/3/e42)
37. Rovers MM, Straatman H, Ingels K, van der Wilt GJ, van den Broek P, Zielhuis GA. Generalizability of trial results based on randomized versus nonrandomized allocation of OME infants to ventilation tubes or watchful waiting. *J Clin Epidemiol.* 2001;54(8):789–794
  38. Rovers MM, Straatman H, Ingels K, van der Wilt GJ, van den Broek P, Zielhuis GA. The effect of short-term ventilation tubes versus watchful waiting on hearing in young children with persistent otitis media with effusion: a randomized trial. *Ear Hear.* 2001;22(3):191–199
  39. Velepich M, Starcevic R, Bonifacic M, et al. The clinical status of the eardrum: an inclusion criterion for the treatment of chronic secretory otitis media in children. *Int J Pediatr Otorhinolaryngol.* 2011;75(5):686–690
  40. Vlastos IM, Houlakis M, Kandiloros D, Manolopoulos L, Ferekidis E, Yiotakis I. Adenoidectomy plus tympanostomy tube insertion versus adenoidectomy plus myringotomy in children with obstructive sleep apnoea syndrome. *J Laryngol Otol.* 2011;125(3):274–278
  41. Wilks J, Maw R, Peters TJ, Harvey I, Golding J. Randomised controlled trial of early surgery versus watchful waiting for glue ear: the effect on behavioural problems in pre-school children. *Clin Otolaryngol Allied Sci.* 2000;25(3):209–214
  42. Yagi HI. The surgical treatment of secretory otitis media in children. *J Laryngol Otol.* 1977;91(3):267–270
  43. Robson AK, Blanshard JD, Jones K, Albery EH, Smith IM, Maw AR. A conservative approach to the management of otitis media with effusion in cleft palate children. *J Laryngol Otol.* 1992;106(9):788–792
  44. Hubbard TW, Paradise JL, McWilliams BJ, Elster BA, Taylor FH. Consequences of unremitting middle-ear disease in early life. Otologic, audiologic, and developmental findings in children with cleft palate. *N Engl J Med.* 1985;312(24):1529–1534
  45. Forquer BD, Linthicum FH Jr. Middle ear effusion in children: a report of treatment in 500 cases. *West J Med.* 1982;137(5):370–374
  46. Marshak G, Neriah ZB. Adenoidectomy versus tympanostomy in chronic secretory otitis media. *Ann Otol Rhinol Laryngol Suppl.* 1980;89(3 pt 2):316–318
  47. Peters SA, Grievink EH, van Bon WH, Schilder AG. The effects of early bilateral otitis media with effusion on educational attainment: a prospective cohort study. *J Learn Disabil.* 1994;27(2):111–121
  48. Grievink EH, Peters SA, van Bon WH, Schilder AG. The effects of early bilateral otitis media with effusion on language ability: a prospective cohort study. *J Speech Hear Res.* 1993;36(5):1004–1012
  49. Schilder AG, Hak E, Straatman H, Zielhuis GA, van Bon WH, van den Broek P. Long-term effects of ventilation tubes for persistent otitis media with effusion in children. *Clin Otolaryngol Allied Sci.* 1997;22(5):423–429
  50. Coyte PC, Croxford R, McIsaac W, Feldman W, Friedberg J. The role of adjuvant adenoidectomy and tonsillectomy in the outcome of the insertion of tympanostomy tubes. *N Engl J Med.* 2001;344(16):1188–1195
  51. Xu HF, Xu LR, He Y, Zheng Q, Zheng Y, Liao XY. [Treatment of cleft palate with secretory otitis media]. *Zhonghua Kou Qiang Yi Xue Za Zhi.* 2003;38(4):269–270
  52. de Beer BA, Schilder AG, Ingels K, Snik AF, Zielhuis GA, Graamans K. Hearing loss in young adults who had ventilation tube insertion in childhood. *Ann Otol Rhinol Laryngol.* 2004;113(6):438–444
  53. Stenstrom R, Pless IB, Bernard P. Hearing thresholds and tympanic membrane sequelae in children managed medically or surgically for otitis media with effusion. *Arch Pediatr Adolesc Med.* 2005;159(12):1151–1156
  54. Kadhim AL, Spilsbury K, Semmens JB, Coates HL, Lannigan FJ. Adenoidectomy for middle ear effusion: a study of 50,000 children over 24 years. *Laryngoscope.* 2007;117(3):427–433
  55. Motta G, Esposito E, Motta S, et al. Acute recurrent pharyngotonsillitis and otitis media [in Italian]. *Acta Otorhinolaryngol Ital.* 2006;26(5 suppl 84):30–55
  56. Reiter R, Haase S, Brosch S. Repaired cleft palate and ventilation tubes and their associations with cholesteatoma in children and adults. *Cleft Palate Craniofac J.* 2009;46(6):598–602
  57. Kobayashi H, Sakuma T, Yamada N, Suzuki H. Clinical outcomes of ventilation tube placement in children with cleft palate. *Int J Pediatr Otorhinolaryngol.* 2012;76(5):718–721
  58. Wolter NE, Dell SD, James AL, Campisi P. Middle ear ventilation in children with primary ciliary dyskinesia. *Int J Pediatr Otorhinolaryngol.* 2012;76(11):1565–1568
  59. Yousaf M, Inayatullah, Khan F. Medical versus surgical management of otitis media with effusion in children. *J Ayub Med Coll Abbottabad.* 2012;24(1):83–85
  60. Youssef TF, Ahmed MR. Laser-assisted myringotomy versus conventional myringotomy with ventilation tube insertion in treatment of otitis media with effusion: long-term follow-up. *Interv Med Appl Sci.* 2013;5(1):16–20
  61. Tian L, Chen X, Li G. The clinical curative effect of the low temperature plasma ablation adenoidectomy and tympanic membrane indwelling catheter in parallel or not used on childhood patients with secretory otitis media [in Chinese]. *Lin Chuang Er Bi Yan Hou Tou Jing Wai Ke Za Zhi.* 2015;29(5):415–417
  62. Li S, Zhang H, Wei Y, et al. Clinical comparative study on the treatment characteristics of secretory otitis media between cleft and non-cleft palate patients [in Chinese]. *Hua Xi Kou Qiang Yi Xue Za Zhi.* 2015;33(3):259–262
  63. Hu S, Patel NA, Shinhar S. Follow-up audiometry after bilateral myringotomy and tympanostomy tube insertion. *Int J Pediatr Otorhinolaryngol.* 2015;79(12):2068–2071
  64. Kuscu O, Gunaydin RO, Icen M, et al. The effect of early routine grommet insertion on management of otitis media with effusion in children with cleft palate. *J Craniomaxillofac Surg.* 2015;43(10):2112–2115
  65. Niclasen J, Obel C, Homøe P, Kørvel-Hanquist A, Dammeyer J.

- Associations between otitis media and child behavioural and learning difficulties: results from a Danish cohort. *Int J Pediatr Otorhinolaryngol.* 2016;84:12–20
66. Gonzalez C, Arnold JE, Woody EA, et al. Prevention of recurrent acute otitis media: chemoprophylaxis versus tympanostomy tubes. *Laryngoscope.* 1986;96(12):1330–1334
  67. Casselbrant ML, Kaleida PH, Rockette HE, et al. Efficacy of antimicrobial prophylaxis and of tympanostomy tube insertion for prevention of recurrent acute otitis media: results of a randomized clinical trial. *Pediatr Infect Dis J.* 1992;11(4):278–286
  68. El-Sayed Y. Treatment of recurrent acute otitis media chemoprophylaxis versus ventilation tubes. *Aust J Otolaryngol.* 1996;2(4):352–355
  69. Mattila PS, Joki-Erkkilä VP, Kilpi T, Jokinen J, Herva E, Puhakka H. Prevention of otitis media by adenoidectomy in children younger than 2 years. *Arch Otolaryngol Head Neck Surg.* 2003;129(2):163–168
  70. Hammarén-Malmi S, Saxen H, Tarkkanen J, Mattila PS. Adenoidectomy does not significantly reduce the incidence of otitis media in conjunction with the insertion of tympanostomy tubes in children who are younger than 4 years: a randomized trial. *Pediatrics.* 2005;116(1):185–189
  71. Kujala T, Alho OP, Luotonen J, et al. Tympanostomy with and without adenoidectomy for the prevention of recurrences of acute otitis media: a randomized controlled trial. *Pediatr Infect Dis J.* 2012;31(6):565–569
  72. Kujala T, Alho OP, Kristo A, et al. Quality of life after surgery for recurrent otitis media in a randomized controlled trial. *Pediatr Infect Dis J.* 2014;33(7):715–719
  73. Grindler DJ, Blank SJ, Schulz KA, Witsell DL, Lieu JE. Impact of otitis media severity on children's quality of life. *Otolaryngol Head Neck Surg.* 2014;151(2):333–340
  74. Brodsky L, Brookhauser P, Chait D, et al. Office-based insertion of pressure equalization tubes: the role of laser-assisted tympanic membrane fenestration. *Laryngoscope.* 1999;109(12):2009–2014
  75. Hoffmann KK, Thompson GK, Burke BL, Derkay CS. Anesthetic complications of tympanostomy tube placement in children. *Arch Otolaryngol Head Neck Surg.* 2002;128(9):1040–1043
  76. Isaacson G. Six Sigma tympanostomy tube insertion: achieving the highest safety levels during residency training. *Otolaryngol Head Neck Surg.* 2008;139(3):353–357
  77. Hörmann K, Roehrs M. Middle-ear findings in young cleft lip and palate children. Comparison of two treatment clinics [in German]. *Dtsch Z Mund Kiefer Gesichtschir.* 1991;15(2):149–152
  78. Suetake M, Kobayashi T, Takasaka T, Shinkawa H. Is change in middle ear air volume following ventilation tube insertion a reliable prognostic indicator? *Acta Otolaryngol Suppl.* 1990;471:73–80
  79. Gates GA, Avery CA, Cooper JC Jr, Prihoda TJ. Chronic secretory otitis media: effects of surgical management. *Ann Otol Rhinol Laryngol Suppl.* 1989;138:2–32
  80. Weigel MT, Parker MY, Goldsmith MM, Postma DS, Pillsbury HC. A prospective randomized study of four commonly used tympanostomy tubes. *Laryngoscope.* 1989;99(3):252–256
  81. Gates GA, Avery C, Prihoda TJ, Holt GR. Delayed onset post-tympanotomy otorrhea. *Otolaryngol Head Neck Surg.* 1988;98(2):111–115
  82. Debruyne F, Jorissen M, Poelmans J. Otorrhea during transtympanic ventilation. *Am J Otol.* 1988;9(4):316–317
  83. Tavin ME, Gordon M, Ruben RJ. Hearing results with the use of different tympanostomy tubes: a prospective study. *Int J Pediatr Otorhinolaryngol.* 1988;15(1):39–50
  84. Gates GA, Avery CA, Prihoda TJ, Cooper JC Jr. Effectiveness of adenoidectomy and tympanostomy tubes in the treatment of chronic otitis media with effusion. *N Engl J Med.* 1987;317(23):1444–1451
  85. Debruyne F, Jorissen M, Poelmans J. Follow-up of trans-tympanum tubes [in Dutch]. *Acta Otorhinolaryngol Belg.* 1986;40(4):666–677
  86. Rothera MP, Grant HR. Long-term ventilation of the middle ear using the Goode T-tube. *J Laryngol Otol.* 1985;99(4):335–337
  87. Levinson SR, Gill AJ, Teich L. Semipermeable membrane tubes: a prospective study. *Otolaryngol Head Neck Surg.* 1982;90(5):622–628
  88. Plotkin RP. Middle ear ventilation with the Castelli membrane tube. *Laryngoscope.* 1981;91(7):1173–1175
  89. Mandel EM, Casselbrant ML, Kurs-Lasky M. Acute otorrhea: bacteriology of a common complication of tympanostomy tubes. *Ann Otol Rhinol Laryngol.* 1994;103(9):713–718
  90. Brown JA. Management of ventilation tubes: preventing premature extrusion. *J S C Med Assoc.* 1993;89(9):427–430
  91. Debruyne F, Degroote M. One-year follow-up after tympanostomy tube insertion for recurrent acute otitis media. *ORL J Otorhinolaryngol Relat Spec.* 1993;55(4):226–229
  92. Owen MJ, Norcross-Nechay K, Howie VM. Brainstem auditory evoked potentials in young children before and after tympanostomy tube placement. *Int J Pediatr Otorhinolaryngol.* 1993;25(1–3):105–117
  93. Heaton JM, Mills RP. Otorrhea via ventilation tubes in adults and children. *Clin Otolaryngol Allied Sci.* 1993;18(6):496–499
  94. Siddiqui N, Toynton S, Mangat KS. Results of middle ear ventilation with 'Mangat' T-tubes. *Int J Pediatr Otorhinolaryngol.* 1997;40(2–3):91–96
  95. Golz A, Netzer A, Joachims HZ, Westerman ST, Gilbert LM. Ventilation tubes and persisting tympanic membrane perforations. *Otolaryngol Head Neck Surg.* 1999;120(4):524–527
  96. Valtonen H, Qvarnberg Y, Nuutinen J. Tympanostomy in young children with recurrent otitis media. A long-term follow-up study. *J Laryngol Otol.* 1999;113(3):207–211
  97. Rosenfeld RM, Bhaya MH, Bower CM, et al. Impact of tympanostomy tubes on child quality of life. *Arch Otolaryngol Head Neck Surg.* 2000;126(5):585–592
  98. Ah-Tye C, Paradise JL, Colborn DK. Otorrhea in young children after tympanostomy-tube placement

- for persistent middle-ear effusion: prevalence, incidence, and duration. *Pediatrics*. 2001;107(6):1251–1258
99. Valtonen H, Dietz A, Qvarnberg Y. Long-term clinical, audiologic, and radiologic outcomes in palate cleft children treated with early tympanostomy for otitis media with effusion: a controlled prospective study. *Laryngoscope*. 2005;115(8):1512–1516
  100. Pereira MB, Pereira DR, Costa SS. Tympanostomy tube sequelae in children with otitis media with effusion: a three-year follow-up study. *Rev Bras Otorrinolaringol (Engl Ed)*. 2005;71(4):415–420
  101. Spielmann PM, McKee H, Adamson RM, Thiel G, Schenk D, Hussain SS. Follow up after middle-ear ventilation tube insertion: what is needed and when? *J Laryngol Otol*. 2008;122(6):580–583
  102. O'Reilly RC, He Z, Bloedon E, et al. The role of extraesophageal reflux in otitis media in infants and children. *Laryngoscope*. 2008;118(7 pt 2 suppl 116):1–9
  103. Ida JB, Worley NK, Amedee RG. Gold laser adenoidectomy: long-term safety and efficacy results. *Int J Pediatr Otorhinolaryngol*. 2009;73(6):829–831
  104. Jung H, Lee SK, Cha SH, Byun JY, Park MS, Yeo SG. Current bacteriology of chronic otitis media with effusion: high rate of nosocomial infection and decreased antibiotic sensitivity. *J Infect*. 2009;59(5):308–316
  105. Kinnari TJ, Aarnisalo AA, Rihkanen H, Lundin J, Jero J. Can head position after anesthesia cause occlusion of the tympanostomy tube? *J Otolaryngol Head Neck Surg*. 2010;39(1):1–4
  106. Marzouk HA, Nathawad R, Hammerschlag MR, Weedon J, Bachman D, Goldstein NA. Methicillin-resistant *Staphylococcus aureus* colonization in otitis-prone children. *Arch Otolaryngol Head Neck Surg*. 2011;137(12):1217–1222
  107. Tuli BS, Parmar TL, Singh B. Evaluation of tympanostomy tubes in middle ear affections. *Indian J Otolaryngol Head Neck Surg*. 2001;53(3):217–220
  108. van Dongen TM, van der Heijden GJ, Freling HG, Venekamp RP, Schilder AG. Parent-reported otorrhea in children with tympanostomy tubes: incidence and predictors. *PLoS One*. 2013;8(7):e69062
  109. Saki N, Nikakhlagh S, Salehe F, Darabifard A. Incidence of complications developed after the insertion of ventilation tube in children under 6 years old in 2008-2009. *Iran J Otorhinolaryngol*. 2012;24(66):15–18
  110. Smillie I, Robertson S, Yule A, Wynne DM, Russell CJ. Complications of ventilation tube insertion in children with and without cleft palate: a nested case-control comparison. *JAMA Otolaryngol Head Neck Surg*. 2014;140(10):940–943
  111. Luo HN, Ma SJ, Sheng Y, et al. Pepsin deteriorates prognosis of children with otitis media with effusion who undergo myringotomy or tympanostomy tube insertion. *Int J Pediatr Otorhinolaryngol*. 2014;78(12):2250–2254
  112. Powell J, Powell S, Lennon M, Ho A, Murrant N. Paediatric ventilation tube insertion: our experience of seventy-five children in audiology-led follow-up. *Clin Otolaryngol*. 2015;40(4):385–389
  113. Birck HG, Mravec JJ. Myringostomy for middle ear effusions. Results of a two-year study. *Ann Otol Rhinol Laryngol*. 1976;85(2 suppl 25 pt 2):263–267
  114. Mackenzie IJ. Factors affecting the extrusion rates of ventilation tubes. *J R Soc Med*. 1984;77(9):751–753
  115. Eliachar I, Joachims HZ, Goldsher M, Golz A. Assessment of long-term middle ear ventilation. *Acta Otolaryngol*. 1983;96(1–2):105–112
  116. Jamal TS. Avoidance of postoperative blockage of ventilation tubes. *Laryngoscope*. 1995;105(8 pt 1):833–834
  117. Walker P. Ventilation tube duration versus site of placement. *Aust N Z J Surg*. 1997;67(8):571–572
  118. Roland PS, Kreisler LS, Reese B, et al. Topical ciprofloxacin/dexamethasone otic suspension is superior to ofloxacin otic solution in the treatment of children with acute otitis media with otorrhea through tympanostomy tubes. *Pediatrics*. 2004;113(1 pt 1). Available at: [www.pediatrics.org/cgi/content/full/113/1/e40](http://www.pediatrics.org/cgi/content/full/113/1/e40)
  119. Wallace HC, Newbegin CJ. Does ENT outpatient review at 1-week post ventilation tube insertion improve outcome at 1 month in paediatric patients? *Clin Otolaryngol Allied Sci*. 2004;29(6):595–597
  120. Allen J, Morton RP, Ahmad Z. Early post-operative morbidity after tympanostomy tube insertion. *J Laryngol Otol*. 2005;119(9):699–703
  121. Dohar J, Giles W, Roland P, et al. Topical ciprofloxacin/dexamethasone superior to oral amoxicillin/clavulanic acid in acute otitis media with otorrhea through tympanostomy tubes. *Pediatrics*. 2006;118(3). Available at: [www.pediatrics.org/cgi/content/full/118/3/e561](http://www.pediatrics.org/cgi/content/full/118/3/e561)
  122. Muenker G. Results after treatment of otitis media with effusion. *Ann Otol Rhinol Laryngol Suppl*. 1980;89(3 pt 2):308–311
  123. van Baarle PW, Wentges RT. Extrusion of transtympanic ventilating tubes, relative to the site of insertion. *ORL J Otorhinolaryngol Relat Spec*. 1975;37(1):35–40
  124. Kokko E, Palva T. Clinical results and complications of tympanostomy. *Ann Otol Rhinol Laryngol*. 1976;85(2 suppl 25 pt 2):277–279
  125. Paradise JL, Bluestone CD, Rogers KD, et al. Efficacy of adenoidectomy for recurrent otitis media in children previously treated with tympanostomy-tube placement. Results of parallel randomized and nonrandomized trials. *JAMA*. 1990;263(15):2066–2073
  126. MacKinnon DM. The sequel to myringotomy for exudative otitis media. *J Laryngol Otol*. 1971;85(8):773–794
  127. Daly KA, Hunter LL, Lindgren BR, Margolis R, Giebink GS. Chronic otitis media with effusion sequelae in children treated with tubes. *Arch Otolaryngol Head Neck Surg*. 2003;129(5):517–522
  128. Praveen CV, Terry RM. Does passive smoking affect the outcome of grommet insertion in children? *J Laryngol Otol*. 2005;119(6):448–454
  129. Khan F, Asif M, Farooqi GH, Shah SA, Sajid T, Ghani R. Management outcome of secretory otitis media. *J Ayub Med Coll Abbottabad*. 2006;18(1):55–58

130. Hammarén-Malmi S, Saxen H, Tarkkanen J, Mattila PS. Passive smoking after tympanostomy and risk of recurrent acute otitis media. *Int J Pediatr Otorhinolaryngol.* 2007;71(8):1305–1310
131. Fiebach A, Matschke RG. Duration and complications following grommet insertion in childhood [in Dutch]. *HNO.* 1987;35(2):61–66
132. Fior R, Veljak C. Late results and complications of tympanostomy tube insertion for prophylaxis of recurrent purulent otitis media in pediatric age. *Int J Pediatr Otorhinolaryngol.* 1984;8(2):139–146
133. Van Cauwenberge P, Cauwe F, Kluyskens P. The long-term results of the treatment with transtympanic ventilation tubes in children with chronic secretory otitis media. *Int J Pediatr Otorhinolaryngol.* 1979;1(2):109–116
134. Tos M, Poulsen G. Secretory otitis media. Late results of treatment with grommets. *Arch Otolaryngol.* 1976;102(11):672–675
135. Costa OA, Balieiro RO. Secretory otitis media in Brazilian children. *Scand Audiol Suppl.* 1986;26:93–94
136. Barfoed C, Rosborg J. Secretory otitis media. Long-term observations after treatment with grommets. *Arch Otolaryngol.* 1980;106(9):553–556
137. Levine S, Daly K, Giebink GS. Tympanic membrane perforations and tympanostomy tubes. *Ann Otol Rhinol Laryngol Suppl.* 1994;163:27–30
138. Hampton SM, Adams DA. Perforation rates after ventilation tube insertion: does the positioning of the tube matter? *Clin Otolaryngol Allied Sci.* 1996;21(6):548–549
139. Postma DS, Poole MD, Wu SM, Tober R. The impact of day care on ventilation tube insertion. *Int J Pediatr Otorhinolaryngol.* 1997;41(3):253–262
140. Valtonen HJ, Qvarnberg YH, Nuutinen J. Otolological and audiological outcomes five years after tympanostomy in early childhood. *Laryngoscope.* 2002;112(4):669–675
141. Valtonen H, Tuomilehto H, Qvarnberg Y, Nuutinen J. A 14-year prospective follow-up study of children treated early in life with tympanostomy tubes: part 1: clinical outcomes. *Arch Otolaryngol Head Neck Surg.* 2005;131(4):293–298
142. De Beer BA, Schilder AG, Zielhuis GA, Graamans K. Natural course of tympanic membrane pathology related to otitis media and ventilation tubes between ages 8 and 18 years. *Otol Neurotol.* 2005;26(5):1016–1021
143. Carignan M, Dorion D, Stephenson MF, Rouleau M. First myringotomy with insertion of a modified Goode T-Tube: changing the perforation paradigm. *J Otolaryngol.* 2006;35(5):287–291
144. Florentzson R, Finizia C. Transmyringeal ventilation tube treatment: a 10-year cohort study. *Int J Pediatr Otorhinolaryngol.* 2012;76(8):1117–1122
145. O’Niel MB, Cassidy LD, Link TR, Kerschner JE. Tracking tympanostomy tube outcomes in pediatric patients with otitis media using an electronic database. *Int J Pediatr Otorhinolaryngol.* 2015;79(8):1275–1278
146. Gundersen T, Tønning FM. Ventilating tubes in the middle ear. *Arch Otolaryngol.* 1976;102(4):198–199
147. Tos M, Stangerup SE, Larsen P. Dynamics of eardrum changes following secretory otitis. A prospective study. *Arch Otolaryngol Head Neck Surg.* 1987;113(4):380–385
148. Slack RW, Maw AR, Capper JW, Kelly S. Prospective study of tympanosclerosis developing after grommet insertion. *J Laryngol Otol.* 1984;98(8):771–774
149. Daly KA, Hunter LL, Levine SC, Lindgren BR, Giebink GS. Relationships between otitis media sequelae and age. *Laryngoscope.* 1998;108(9):1306–1310
150. Koc A, Uneri C. Sex distribution in children with tympanosclerosis after insertion of a tympanostomy tube. *Eur Arch Otorhinolaryngol.* 2001;258(1):16–19
151. Friedman EM, Sprecher RC, Simon S, Dunn JK. Quantitation and prevalence of tympanosclerosis in a pediatric otolaryngology clinic. *Int J Pediatr Otorhinolaryngol.* 2001;60(3):205–211
152. Bonding P, Lorenzen E. Chronic secretory otitis media—long-term results after treatment with grommets. *ORL J Otorhinolaryngol Relat Spec.* 1974;36(4):227–235
153. Bonding P, Lorenzen E. Cicatricial changes of the eardrum after treatment with grommets. *Acta Otolaryngol.* 1973;75(4):275–276
154. Li Y, Hunter LL, Margolis RH, et al. Prospective study of tympanic membrane retraction, hearing loss, and multifrequency tympanometry. *Otolaryngol Head Neck Surg.* 1999;121(5):514–522
155. Tos M, Poulsen G. Attic retractions following secretory otitis. *Acta Otolaryngol.* 1980;89(5–6):479–486
156. Golz A, Goldenberg D, Netzer A, et al. Cholesteatomas associated with ventilation tube insertion. *Arch Otolaryngol Head Neck Surg.* 1999;125(7):754–757
157. Spilsbury K, Ha JF, Semmens JB, Lannigan F. Cholesteatoma in cleft lip and palate: a population-based follow-up study of children after ventilation tubes. *Laryngoscope.* 2013;123(8):2024–2029
158. Djurhuus BD, Christensen K, Skyttke A, Faber CE. The impact of ventilation tubes in otitis media on the risk of cholesteatoma on a national level. *Int J Pediatr Otorhinolaryngol.* 2015;79(4):605–609
159. Rovers MM, Black N, Browning GG, Maw R, Zielhuis GA, Haggard MP. Grommets in otitis media with effusion: an individual patient data meta-analysis. *Arch Dis Child.* 2005;90(5):480–485
160. Boonacker CW, Rovers MM, Browning GG, Hoes AW, Schilder AG, Burton MJ. Adenoidectomy with or without grommets for children with otitis media: an individual patient data meta-analysis. *Health Technol Assess.* 2014;18(5):1–118
161. Rosenfeld RM, Shin JJ, Schwartz SR, et al. Clinical practice guideline: otitis media with effusion (update). *Otolaryngol Head Neck Surg.* 2016;154(suppl 1):S1–S41

**Effectiveness of Tympanostomy Tubes for Otitis Media: A Meta-analysis**  
Dale W. Steele, Gaelen P. Adam, Mengyang Di, Christopher H. Halladay, Ethan M.  
Balk and Thomas A. Trikalinos  
*Pediatrics* 2017;139;  
DOI: 10.1542/peds.2017-0125 originally published online May 16, 2017;

**Updated Information & Services**

including high resolution figures, can be found at:  
<http://pediatrics.aappublications.org/content/139/6/e20170125>

**References**

This article cites 157 articles, 11 of which you can access for free at:  
<http://pediatrics.aappublications.org/content/139/6/e20170125#BIBL>

**Subspecialty Collections**

This article, along with others on similar topics, appears in the following collection(s):  
**Ear, Nose & Throat Disorders**  
[http://www.aappublications.org/cgi/collection/ear\\_nose\\_-\\_throat\\_disorders\\_sub](http://www.aappublications.org/cgi/collection/ear_nose_-_throat_disorders_sub)  
**Otitis Media**  
[http://www.aappublications.org/cgi/collection/otitis\\_media\\_sub](http://www.aappublications.org/cgi/collection/otitis_media_sub)  
**Evidence-Based Medicine**  
[http://www.aappublications.org/cgi/collection/evidence-based\\_medicine\\_sub](http://www.aappublications.org/cgi/collection/evidence-based_medicine_sub)

**Permissions & Licensing**

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:  
<http://www.aappublications.org/site/misc/Permissions.xhtml>

**Reprints**

Information about ordering reprints can be found online:  
<http://www.aappublications.org/site/misc/reprints.xhtml>

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



# PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

## **Effectiveness of Tympanostomy Tubes for Otitis Media: A Meta-analysis**

Dale W. Steele, Gaelen P. Adam, Mengyang Di, Christopher H. Halladay, Ethan M. Balk and Thomas A. Trikalinos

*Pediatrics* 2017;139;

DOI: 10.1542/peds.2017-0125 originally published online May 16, 2017;

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/139/6/e20170125>

Data Supplement at:

<http://pediatrics.aappublications.org/content/suppl/2017/05/12/peds.2017-0125.DCSupplemental>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2017 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

