

Secular Trend of Age at Menarche in Chinese Adolescents Born From 1973 to 2004

Xin Meng, MD, Suyun Li, MD, PhD, Wenhou Duan, MD, Yanxin Sun, MPH, Chongqi Jia, MD, PhD

abstract

OBJECTIVES: To estimate secular trend and factors influencing the age at menarche for Chinese girls born from 1973 to 2004.

METHODS: Data were collected from the China Health and Nutrition Survey, a prospective open cohort and an ongoing nationwide health and nutrition survey, consisting of 3199 apparently healthy Chinese girls aged 6 to 18 years at entry from 1991 to 2011. Average age at menarche and its 95% confidence interval were estimated by weighted means of Kaplan-Meier survival analysis. Cox proportional hazard models were applied to identify the independent predictive factors of age at menarche.

RESULTS: The age at menarche declined from 14.25 in Chinese girls born before 1976 to 12.60 in girls born after 2000, with an estimated decline of 0.51 years per decade ($P < .001$). The downward trend of age at menarche for rural girls was greater than for urban girls (0.62 vs 0.35 years per decade; $P < .001$). Girls living in urban areas, living in central areas, with high BMI, and with high-energy diets had a high risk of early menarche, whereas girls living on the east coast who had high-carbohydrate diets had a low risk of early menarche.

CONCLUSIONS: The authors of this study found a continuous downward secular trend of age at menarche for Chinese girls in both urban and rural areas born from 1973 to 2004. Among all variables, urban residency, BMI, and energy intake are negatively associated with age at menarche, whereas carbohydrate intake is positively associated with age at menarche.



Department of Epidemiology, Shandong University, Jinan, Shandong, People's Republic of China

Dr Meng conceptualized and designed the study, conducted the initial analyses, and drafted and revised the initial manuscript; Dr Li reviewed and revised the manuscript; Dr Duan and Ms Sun conducted the initial analyses; Dr Jia conceptualized the study and critically reviewed the manuscript; and all authors approved the final manuscript as submitted.

DOI: <https://doi.org/10.1542/peds.2017-0085>

Accepted for publication May 12, 2017

Address correspondence to Chongqi Jia, MD, PhD, Department of Epidemiology, Shandong University, No. 44, Wenhuxi Rd, Jinan 250012, Shandong, People's Republic of China. E-mail: jiachongqi@sdu.edu.cn

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2017 by the American Academy of Pediatrics

FINANCIAL DISCLOSURE: The authors have indicated they have no financial relationships relevant to this article to disclose.

FUNDING: This research uses data from the China Health and Nutrition Survey. Major funding for the survey and data dissemination from 1991 to 2004 came from the National Institutes of Health (NIH) (grants P01-HD28076 and HD30880). Additional funding has come from the NIH (grant HD39183), the Carolina Population Center (in particular, the Carolina Population Center funded the

WHAT'S KNOWN ON THIS SUBJECT: Age at menarche for girls around the world has been decreasing, although the rate is slowing in industrialized countries.

WHAT THIS STUDY ADDS: There is a continuing downward trend in age at menarche for Chinese girls born from 1973 to 2004, which is greater in rural girls.

To cite: Meng X, Li S, Duan W, et al. Secular Trend of Age at Menarche in Chinese Adolescents Born From 1973 to 2004. *Pediatrics*. 2017;140(2):e20170085

Previous studies have found that the average age at menarche has decreased from ~16 to 13 years over the past 180 years, with some variation across different countries.¹ Although this trend toward earlier menarche may be slowing in industrialized countries,¹⁻³ it is still more significant in other countries.⁴⁻⁷ Little is known about the trends in menarche in China. China changed dramatically during the past few decades in many aspects, especially socioeconomic status, an important influencing factor of the onset of puberty.^{3,5,8}

The objectives of this study were to evaluate trends in the average age at menarche in China and to explore the demographic, socioeconomic, nutritional, and lifestyle characteristics associated with age at menarche in a large sample of Chinese girls born over the past 3 decades.

METHODS

China Health and Nutrition Survey

The China Health and Nutrition Survey (CHNS) is an ongoing open cohort established in 1989 and updated in 1989, 1991, 1993, 1997, 2000, 2004, 2006, 2009, 2011, and 2015.⁹ It is an international collaborative project between the Carolina Population Center at the University of North Carolina at Chapel Hill and the National Institute for Nutrition and Health at the Chinese Center for Disease Control and Prevention. Briefly, a 7-day period using a multistage, random cluster design was employed to obtain nationally representative information, taken from interviews and direct examinations, on the health and nutritional status of the Chinese population.⁹ Details of the CHNS have been described elsewhere.¹⁰⁻¹²

Sample

Girls aged 6 to 18 years were asked, "Have you ever had a menstrual period?" to define their menstrual status, and postmenarcheal girls were then asked, "How old were you when the periods started?" to identify the age of their first period. The questionnaires for girls aged 6 to 10 years were completed by a parent or guardian. Girls aged 11 to 18 years completed the questionnaires by themselves.

The outcome of interest was the attainment of menarche and the age at menarche assessed as month and year. The data were obtained from the aforementioned questions. To reduce potential misclassification, only the first reported ages at menarche in the panel data were used.

The survey included 8638 households with 35 703 participants among 15 provinces. With the exclusion of the surveys in 1989 and 2015 (because the data of 1989 on menarche were not contained and the data of 2015 were not yet available), 3483 girls aged 6 to 18 years were included. Among them, 284 girls with missing information on menarche were excluded. A total of 3199 Chinese girls born between 1973 and 2004 remained in the analyses. Birth years from 1973 to 1976 were combined because the total number of subjects included in the final analyses was small. Birth years from 2000 to 2004 were combined because <10% of girls born after 2000 had menarche during the interview,¹³ and statistical power was low if analyzed separately. On the basis of the birth years, girls were categorized in 5 groups, that is, 1973 to 1980, 1981 to 1985, 1986 to 1990, 1991 to 1995, and 1996 to 2004 (hereinafter referred to as "5 birth-year groups").

Predictors of Age at Menarche

BMI was categorized as 4 levels on the basis of the age-specified BMI

standard for Chinese teenagers,^{14,15} in which girls with different ages had different criteria.

Total energy and macronutrient (carbohydrate, protein, and fat) intake per day were estimated by using 3 24-hour recalls and by directly weighing food consumed during the survey^{9,10} and were categorized as high, normal, or low according to the Dietary Reference Intakes for Chinese teenagers.¹⁶ The carbohydrate and fat intakes are adequate when the supplying energy accounts for 50% to 65% and 20% to 30% of total energy,¹⁶ respectively. The standards of energy (kilocalories) and protein (grams) intakes are different for girls with different ages.¹⁶

As presented in Table 1, other variables (ethnicity, region, rural or urban residency, household income per person, parental education, living with parents, and exercise) were all assessed as categorical variables. The household income per person was divided into 4 categories by quartile. Region was categorized as central area (Henan, Hubei, Hunan), east coast (Beijing, Jiangsu, Shandong, Shanghai), northeastern area (Liaoning, Heilongjiang), and western area (Chongqing, Guangxi, Guizhou). Teenagers' physical activity was estimated by using the summary of exercise frequency at school and outside of campus, and was categorized into 3 categories (≤ 2 times per week, 3-7 times per week, ≥ 8 times per week).

Statistical Analysis

To address the complex sampling design, weights for the examination sample were used in all tests. Kaplan-Meier cumulative probabilities were used to estimate the restricted mean and SEs of age at menarche for each birth year and each category of predictors. Trends in age at menarche over birth year (years per decade) were estimated by

TABLE 1 Univariate and Multivariate Cox Proportional Hazard Models of Age at Menarche

Characteristics	Total	Mean Age at Menarche ^a	Unadjusted HR (95% CI)	Adjusted HR (95% CI) ^b
Total	3199	13.56		
Demographic variables				
Year of birth				
1973–1980	596	14.06	1.00	1.00
1981–1985	697	14.06	0.85 (0.75–0.96)*	0.77 (0.67–0.88)**
1986–1990	762	13.48	1.26 (1.15–1.38)**	1.12 (0.99–1.27)
1991–1995	482	13.00	1.69 (1.53–1.86)**	1.51 (1.33–1.71)**
1996–2004	662	12.75	2.13 (1.87–2.41)**	1.92 (1.70–2.17)**
<i>P</i> trend ^c			<.001	<.001
Region				
Northeastern area	491	13.45	1.00	1.00
East coast	664	13.66	0.91 (0.80–1.03)	0.83 (0.72–0.96)*
Central area	1096	13.47	1.03 (0.94–1.13)	1.13 (1.03–1.25)*
Western area	948	13.66	0.95 (0.84–1.08)	1.02 (0.91–1.15)
Ethnicity				
Han	2710	13.52	1.00	1.00
Minority	452	13.80	0.93 (0.83–1.05)	0.99 (0.87–1.12)
Missing	37	14.50	0.41 (0.17–0.94)*	0.35 (0.17–0.72)*
Socioeconomic variables				
Rural or urban residency				
Rural	2289	13.68	1.00	1.00
Urban	910	13.20	1.32 (1.18–1.48)**	1.26 (1.15–1.38)**
Household income per person				
Q1	785	14.03	1.00	1.00
Q2	786	13.60	1.19 (1.06–1.32)**	1.06 (0.94–1.20)
Q3	786	13.48	1.32 (1.19–1.47)**	1.03 (0.92–1.16)
Q4	784	13.12	1.59 (1.40–1.80)**	0.96 (0.82–1.13)
Missing	58	12.92	1.95 (1.40–2.73)**	1.55 (1.09–2.20)*
<i>P</i> trend ^c			<.001	.301
Paternal education (y)				
≤6	599	13.91	1.00	1.00
7–9	1240	13.58	1.18 (1.06–1.31)*	0.96 (0.86–1.06)
10–12	713	13.48	1.30 (1.12–1.52)**	0.98 (0.84–1.13)
≥13	67	13.37	1.38 (1.22–1.55)**	0.85 (0.72–1.00)*
Missing	171	13.12	1.53 (1.34–1.74)**	0.85 (0.63–1.16)
<i>P</i> trend ^c			<.001	.239
Maternal education (y)				
≤6	1101	13.87	1.00	1.00
7–9	1116	13.40	1.30 (1.18–1.43)**	1.09 (0.97–1.22)
10–12	512	13.42	1.39 (1.21–1.59)**	1.16 (1.02–1.32)*
≥13	161	13.10	1.55 (1.28–1.89)**	1.02 (0.80–1.29)
Missing	309	13.28	1.37 (1.17–1.61)**	0.72 (0.51–1.03)
<i>P</i> trend ^c			<.001	.231
Living with parents				
Without parents	180	12.87	1.00	1.00
Father or mother	295	13.26	0.78 (0.63–0.97)*	0.67 (0.47–0.95)*
Father and mother	2704	13.61	0.66 (0.54–0.79)**	0.55 (0.31–0.98)*
Missing	20	13.70	0.60 (0.39–0.92)*	0.79 (0.50–1.25)
<i>P</i> trend ^c			<.001	.072
Nutritional and lifestyle variables				
BMI				
Underweight	280	14.06	1.00	1.00
Normal	2290	13.54	1.40 (1.21–1.62)**	1.41 (1.23–1.61)**
Overweight	133	12.76	2.13 (1.66–2.72)**	2.05 (1.63–2.57)**
Obese	63	12.63	2.71 (1.83–4.01)**	2.33 (1.58–3.42)**
Missing	433	13.66	1.28 (1.04–1.57)*	1.30 (1.06–1.59)*
<i>P</i> trend ^c			<.001	<.001
Energy intake (kcal/d)				
Low	1563	13.48	1.00	1.00
Normal	982	13.61	0.91 (0.82–1.00)	1.04 (0.91–1.20)
High	534	13.51	1.04 (0.94–1.14)	1.29 (1.11–1.50)**
Missing	120	14.19	0.67 (0.53–0.85)**	0.69 (0.48–1.01)

TABLE 1 Continued

Characteristics	Total	Mean Age at Menarche ^a	Unadjusted HR (95% CI)	Adjusted HR (95% CI) ^b
<i>P</i> trend ^c			.856	.005
Protein intake (g/d)				
Low	856	13.51	1.00	1.00
Normal	652	13.68	0.91 (0.80–1.04)	0.93 (0.83–1.05)
High	1571	13.49	1.02 (0.93–1.16)	0.99 (0.88–1.10)
Missing	120	14.19	0.68 (0.54–0.86)*	—
<i>P</i> trend ^c			.444	.921
Fat intake (%)				
Low	931	14.01	1.00	1.00
Normal	957	13.33	1.52 (1.40–1.66)**	1.15 (0.97–1.36)
High	1191	13.28	1.62 (1.48–1.78)**	1.02 (0.82–1.27)
Missing	120	14.19	0.95 (0.74–1.23)	—
<i>P</i> trend ^c			<.001	.542
Carbohydrate intake (%)				
Low	638	13.18	1.00	1.00
Normal	1260	13.29	0.95 (0.87–1.04)	1.01 (0.90–1.13)
High	1181	13.93	0.62 (0.57–0.68)**	0.77 (0.63–0.94)*
Missing	120	14.19	0.56 (0.45–0.71)**	—
<i>P</i> trend ^c			<.001	.033
Exercise (times/wk)				
≤2	2057	13.72	1.00	1.00
3–7	734	13.26	1.26 (1.13–1.41)**	1.10 (0.99–1.23)
≥8	400	13.16	1.38 (1.21–1.58)**	1.06 (0.93–1.21)
Missing	8	13.39	1.14 (0.69–1.87)	1.32 (0.69–2.53)
<i>P</i> trend ^c			<.001	.176

Calculated with application of weights for the CHNS sample.

^a From Kaplan-Meier survival probabilities.

^b Adjusted for other covariates in the table.

^c Calculated by using the median value of each category as a single ordinal variable in the Cox regression model.

* $P < .05$.

** $P < .01$.

using the slope of linear regression model with mean age at menarche as the dependent variable and year of birth as the predictor. The trends of age at menarche were stratified and compared between rural and urban girls and Han and minority ethnicities. The statistical significance of these trends was determined by using tests for linear trends for year of birth in multivariate Cox proportional hazard models.

Cox proportional hazard models were used to examine the association of age at menarche with each predictor. Dummy variables were set for each predictor because all covariates were transferred into categorical data. Unadjusted and adjusted hazard ratios (HRs) and 95% confidence intervals (CIs) were estimated by using univariate and multivariate Cox proportional hazard models, respectively. Additionally, ties were handled by using the

Breslow method. The proportional hazards assumption in the Cox regression model was tested by using Schoenfeld residuals. Linear trends of HRs were tested by using the median value of each category as a single ordinal variable in the Cox regression model. Analyses were conducted with the statistical software Stata 13.1 (StataCorp, College Station, TX). Two-sided tests were used, and $P < .05$ was considered statistically significant.

RESULTS

The Age at Menarche

All the characteristics and other results are presented in Table 1. The weighted percentiles in the 5 birth-year groups for girls who had attained menarche in the last survey were 64.98, 34.68, 62.54, 72.71 and 31.81, respectively. Additionally, the weighted percentiles in the age groups of 6 to 9, 10 to 18, and 14

to 18 for girls who had attained menarche were 0.78, 40.32, and 81.72, respectively. Forty-seven percent girls had not had menarche.

Overall, for girls born during the 30-year period, the weighted mean age at menarche was estimated at 13.56 years. Age at menarche decreased substantially from 14.25 to 12.60 years, with an estimated decline of 0.51 years per decade ($P < .001$) (Fig 1). The results were also compared with other countries in Fig 2.

On average, during the 30-year period, urban girls had menarche 5.76 months (0.48 years) earlier than rural girls (at 13.20 vs 13.68 years). Although there were downward secular trends of age at menarche for both rural and urban girls, these 2 trends were significantly different ($P < .001$). The weighted mean age at menarche decreased by ~2.67 years for rural girls versus 1.34 years for urban girls (Fig 3). During the

30-year period, the mean age at menarche for rural girls decreased substantially from 14.51 years in the birth year of 1976 to 11.84 years in the birth year of 2000 (0.62 years per decade; $P < .001$). In contrast, the downward secular trend for urban girls was mild, with the mean age initially decreasing from 13.39 to 12.05 years (0.35 years per decade; $P < .001$). Besides, a significant difference between rural and urban girls born in the 1980s was found in mean age at menarche (14.00 vs 13.38 years), whereas mean ages were similar for those born after 1995 (12.75 vs 12.73 years).

On average, Han girls had menarche 3.36 months (0.28 years) earlier than minority girls (at 13.52 vs 13.80 years). The secular trends of age at menarche were not significant between Han girls (13.93–12.55 years; 0.52 years per decade, $P < .001$) and minority girls (14.33–11.80 years; 0.39 years per decade, $P < .001$).

Predictors of Age at Menarche

Results of Cox proportional hazard models are presented in Table 1. All variables in the models satisfy the proportional hazards assumption (all $P > .05$). After multivariate adjustment, the association of early menarche with household income per person, parental education, fat intake, exercise, and living with parents would lose statistical significance, whereas the association with girl's birth year, rural or urban residency, BMI, and carbohydrate intake remained significant. Although energy intake and region were not associated with age at menarche in univariate models, the associations changed after covariates' inclusion in multivariate model.

In detail, the risks of early menarche for girls born late (Fig 4) and girls with high BMI or high energy intake were higher, whereas the risks for




FIGURE 1 Mean age at menarche by year of birth in Chinese girls aged 6 to 18 born between 1973 and 2004.




FIGURE 2 Secular trends of age at menarche in 6 countries.^{8,17–23} Ages at menarche in the Netherlands and the United States were calculated as median age, whereas others were calculated as mean age.

girls with high carbohydrate intake was lower (all P trend $< .05$). Girls living in urban areas had a higher risk of early menarche (HR: 1.26, 95% CI: 1.15–1.38). The hazard of early menarche was significantly higher for

girls living in the central area (HR: 1.13, 95% CI: 1.03–1.25) and lower for girls living in the east coast area (HR: 0.83, 95% CI: 0.72–0.96) when compared with girls living in the northeastern area.




FIGURE 3 Mean age at menarche by year of birth in Chinese girls aged 6 to 18 born between 1973 and 2004, stratified by rural or urban residency.




FIGURE 4 Kaplan-Meier survival probability of age at menarche among Chinese girls aged 6 to 18 born between 1973 and 2004.

DISCUSSION

This study of a national cohort confirms a continuous downward secular trend of age at menarche through birth years of 1973 to 2004

among Chinese girls. Mean age at menarche declined from 14.25 to 12.60 years during the 30-year period, with an average decline of 0.51 years per decade. Moreover, the mean age at menarche for rural

girls decreased markedly (0.62 years per decade), but urban girls showed a smaller decrease (0.35 years per decade). However, secular trends of mean age at menarche between Han girls (0.52 years per decade) and minorities (0.39 years per decade) were not significant.

As presented in Fig 2, the downward trend was common around the world, although the average ages at menarche were different.^{8,17-23} Anderson et al^{24,25} found the mean age at menarche for US girls decreased from 12.75 years in 1963 through 1970 to 12.54 years in 1988 through 1994, and then to 12.34 years in 1999 through 2002. And the age at menarche for Western European girls declined by ~3 to 4 months per decade from 1830 to 1980.²⁶ Hosokawa et al²⁷ concluded that the mean age at menarche for Japanese girls had decreased by 1.6 years over the 50-year period since 1930, from 13.8 to 12.2 years. During the past decade, several researchers reported similar results on age at menarche for Chinese girls. Song et al^{28,29} analyzed a national sample of girls aged 9 to 18 years from the National Survey on the Constitution and Health of Chinese Students. They reported a decreased trend of mean age at menarche from 13.41 years in 1985 to 12.47 years in 2010²⁸ and a difference in mean age at menarche between rural and urban girls (12.92 vs 12.60 years).²⁹ Another retrospective study¹⁹ based on China Kadoorie Biobank found that mean age at menarche decreased from 15.5 years for girls born in 1925 to 14.2 years for girls born in 1978. Although the downward secular trend of age at menarche has recently slowed in Japan, the Netherlands, Germany, and Bulgaria,^{27,30} and remained stable in Belgium and Norway,³⁰ it has continued to decline in China, in both urban and rural areas.

The differences in age at menarche between urban and rural areas well establish that urban girls reach

menarche earlier than rural girls.^{30,31} Our study has also revealed a significant difference between urban and rural girls in age at menarche. A faster downward trend in age at menarche for rural girls is also evident. It is believed that urbanization influences the evolution of age at menarche, probably through increasing BMI.³¹ Studies^{28,32,33} have shown a continuous decrease in the magnitude of the association between age at menarche and rural or urban residency coinciding with the urbanization of China over the past 30 years. According to the official statistics, the rate of urbanization increased from 23.71% in 1985 to 29.04% in 1995 and, in addition, to 49.68% in 2010.^{28,33}

Our study has found that a high BMI was a risk factor of early menarche. Many researchers explained this by an effect of increased body fat on the hypothalamic-pituitary-gonadal axis.^{20,34–36} However, some researchers reported the limited influence of BMI on age at menarche in high-BMI groups¹⁸ and explained it by hyperandrogenism in girls with obesity.^{37–39} Tchernof and Després,⁴⁰ however, indicated that high plasma estradiol levels and low sex hormone binding globulin levels because of puberty were associated with adiposity, which meant that the high BMI resulted from the beginning of menarche. Previous studies have shown that higher energy, protein, and fat intake are positively associated with early menarche, and higher carbohydrate intake is negatively associated with early menarche.^{36,41–47} As shown by our results, high energy intake was positively associated with early menarche, and high carbohydrate intake was positively related to late menarche. The fact that energy intake was associated with age at menarche in multivariate analysis but not in univariate analysis can be explained by the effect of birth time. The association becomes significant

($P < .001$) when only adjusted for birth year. Ages at menarche are different among regions. Girls living in the central area have the biggest risk of early menarche, whereas girls living in the east coast region had menarche latest.

Additionally, the associations of early menarche with socioeconomic factors except rural or urban residency lose statistical significance after adjustment. This suggests that general social and economic status influence age at menarche through other covariates and that these covariates play a more minimal role in influencing the sexual maturation of Chinese girls. It supports the notion that socioeconomic factors are not independent predictors of age at menarche in well-developed countries.^{7,17,18}

Human evidence exists that exposure to endocrine-disrupting agents such as lead and polybrominated biphenyls can induce an earlier onset of menarche.^{39,48,49} However, neither exposure was available for analysis in our multivariate model, despite recognition that minority populations may have higher environmental contaminant exposure. It is possible that girls born late might be more likely to be exposed to these agents and, as a result, had menarche earlier than girls born early.

Strengths of the study are as follows. Firstly, this research used data from a national representative sample with a diverse population and a prospective follow-up study design; thus, the results are more likely to be generalizable to the China population. Secondly, compared with other studies around the world, not only demographic and socioeconomic variables but also nutritional and lifestyle variables were included as factors. And this study used convincing BMI and nutrition category standards formulated by the Chinese government. Thirdly, differing from other cross-sectional studies, our study was conducted in a birth

cohort, which reduced recall bias and increased accuracy.

Several limitations should be kept in mind when interpreting our findings. First, analysis of retrospective data might bring recall bias to our study. Height, weight, and other variables were collected a period after the menarche. The second concern is the large amount of missing data. Approximately half of the girls failed to record their age at menarche because they were too young to have their menarche at last survey or didn't want to answer the questions. Approximately one-fourth of girls were last surveyed before 12 years old. Additionally, some of the factors this study wanted to analyze had missing data. To address this concern, this study generated new categories containing missing data for each variable and calculated the HRs and 95% CIs of the missing groups; however, it still might bring bias to our study.

CONCLUSIONS

Data from the CHNS indicated that the average age at menarche for Chinese girls born from 1973 to 2004 both in urban and rural areas had a continuous downward secular trend. In general, both the downward trend and average age at menarche for rural girls are greater than for urban girls. Additionally, urban residency, BMI, and energy intake are negatively associated with age at menarche, whereas carbohydrate intake is positively associated with age at menarche. Additional studies are warranted to elucidate factors, particularly nutrition and exposure to endocrine-disrupting agents, which may account for the downward secular trend in age at menarche.

ACKNOWLEDGMENTS

This research uses data from the CHNS. Thanks to the National

Institute of Nutrition and Food Safety at the Chinese Center for Disease Control and Prevention, the Carolina Population Center at the University of North Carolina at Chapel Hill, the National Institutes of Health, and the Fogarty

International Center for the CHNS data collection and for the analysis of files from 1989 to 2006 and both parties, plus the China-Japan Friendship Hospital, Ministry of Health, for supporting the 2009 CHNS and future surveys.

ABBREVIATIONS

CHNS: China Health and Nutrition Survey
HR: hazard ratio
CI: confidence interval

1989 China Health and Nutrition Survey), the Ford Foundation, the National Science Foundation (grant INT-9215399), the National Institute of Nutrition and Food Safety (formerly named the Institute of Nutrition and Food Hygiene), and the Chinese Center for Disease Control and Prevention (formerly named the Chinese Academy of Preventive Medicine). Funded by the National Institutes of Health (NIH).

POTENTIAL CONFLICT OF INTEREST: The authors have indicated they have no potential conflicts of interest to disclose.

REFERENCES

1. Parent AS, Teilmann G, Juul A, Skakkebaek NE, Toppari J, Bourguignon JP. The timing of normal puberty and the age limits of sexual precocity: variations around the world, secular trends, and changes after migration. *Endocr Rev.* 2003;24(5):668–693
2. Ulijaszek SJ, Evans E, Miller DS. Age at menarche of European, Afro-Caribbean and Indo-Pakistani schoolgirls living in London. *Ann Hum Biol.* 1991;18(2):167–175
3. Vercauteren M, Susanne C. The secular trend of height and menarche in Belgium: are there any signs of a future stop? *Eur J Pediatr.* 1985;144(4):306–309
4. Apraiz AG. Influence of family size and birth order on menarcheal age of girls from Bilbao city (Biscay, Basque country). *Am J Hum Biol.* 1999;11(6):779–783
5. Kahl H, Schaffrath Rosario A, Schlaud M. [Sexual maturation of children and adolescents in Germany. Results of the German Health Interview and Examination Survey for children and adolescents (KiGGS)]. *Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz.* 2007;50(5–6):677–685
6. Padez C. Social background and age at menarche in Portuguese university students: a note on the secular changes in Portugal. *Am J Hum Biol.* 2003;15(3):415–427
7. Papadimitriou A, Fytanidis G, Douros K, Bakoula C, Nicolaidou P, Fretzayas A. Age at menarche in contemporary Greek girls: evidence for levelling-off of the secular trend. *Acta Paediatr.* 2008;97(6):812–815
8. Euling SY, Herman-Giddens ME, Lee PA, et al. Examination of US puberty-timing data from 1940 to 1994 for secular trends: panel findings. *Pediatrics.* 2008;121(suppl 3):172–191
9. Carolina Population Center. China Health and Nutrition Survey. Available at: www.cpc.unc.edu/projects/china. 2011. Accessed March 20, 2017
10. He K, Du S, Xun P, et al. Consumption of monosodium glutamate in relation to incidence of overweight in Chinese adults: China Health and Nutrition Survey (CHNS). *Am J Clin Nutr.* 2011;93(6):1328–1336
11. Xu X, Byles JE, Shi Z, Hall JJ. Evaluation of older Chinese people's macronutrient intake status: results from the China Health and Nutrition Survey. *Br J Nutr.* 2015;113(1):159–171
12. Popkin BM, Du S, Zhai F, Zhang B. Cohort profile: The China Health and Nutrition Survey—monitoring and understanding socio-economic and health change in China, 1989–2011. *Int J Epidemiol.* 2010;39(6):1435–1440
13. Jansen EC, Herrán OF, Villamor E. Trends and correlates of age at menarche in Colombia: results from a nationally representative survey. *Econ Hum Biol.* 2015;19:138–144
14. National Health and Family Planning Commission of PRC. Screening standard for malnutrition of school-age children and adolescents. Available at: www.nhfpc.gov.cn/ ewebeditor/uploadfile/2014/07/20140704142652587.pdf. Accessed March 20, 2017
15. National Health and Family Planning Commission of PRC. Comprehensive evaluation of children and adolescents development. Available at: www.nhfpc.gov.cn/ewebeditor/uploadfile/2015/04/20150407151300845.pdf. Accessed March 20, 2017
16. Cheng YY. Chinese dietary reference intakes (2013 revised edition). *Acta Nutrimenta Sinica.* 2014(4):313–317
17. Talma H, Schönbeck Y, van Dommelen P, Bakker B, van Buuren S, Hirasings RA. Trends in menarcheal age between 1955 and 2009 in the Netherlands. *PLoS One.* 2013;8(4):e60056
18. Lee MH, Kim SH, Oh M, Lee KW, Park MJ. Age at menarche in Korean adolescents: trends and influencing factors. *Reprod Health.* 2016;13(1):121
19. Lin LL, Zheng B, Lyu J, et al; China Kadoorie Biobank (CKB) Collaborative Group. [Association between age at menarche and height and leg length in adult women: findings from survey in 10 areas in China]. *Zhonghua Liu Xing Bing Xue Za Zhi.* 2016;37(11):1454–1458
20. Harris MA, Prior JC, Koehoorn M. Age at menarche in the Canadian population: secular trends and relationship to adulthood BMI. *J Adolesc Health.* 2008;43(6):548–554
21. Cho GJ, Park HT, Shin JH, et al. Age at menarche in a Korean population: secular trends and influencing factors. *Eur J Pediatr.* 2010;169(1):89–94

22. Rigon F, Bianchin L, Bernasconi S, et al. Update on age at menarche in Italy: toward the leveling off of the secular trend. *J Adolesc Health*. 2010;46(3):238–244
23. Krieger N, Kosheleva A, Waterman PD, Chen JT, Beckfield J, Kiang MV. 50-year trends in US socioeconomic inequalities in health: US-born black and white Americans, 1959–2008. *Int J Epidemiol*. 2014;43(4):1294–1313
24. Anderson SE, Dallal GE, Must A. Relative weight and race influence average age at menarche: results from two nationally representative surveys of US girls studied 25 years apart. *Pediatrics*. 2003;111(4 pt 1):844–850
25. Anderson SE, Must A. Interpreting the continued decline in the average age at menarche: results from two nationally representative surveys of U.S. girls studied 10 years apart. *J Pediatr*. 2005;147(6):753–760
26. Danubio ME, Sanna E. Secular changes in human biological variables in Western countries: an updated review and synthesis. *J Anthropol Sci*. 2008;86:91–112
27. Hosokawa M, Imazeki S, Mizunuma H, Kubota T, Hayashi K. Secular trends in age at menarche and time to establish regular menstrual cycling in Japanese women born between 1930 and 1985. *BMC Womens Health*. 2012;12:19
28. Song Y, Ma J, Wang HJ, et al. Trends of age at menarche and association with body mass index in Chinese school-aged girls, 1985–2010. *J Pediatr*. 2014;165(6):1172–1177.e1
29. Song Y, Ma J, Hu PJ, Zhang B. [Geographic distribution and secular trend of menarche in 9–18 year-old Chinese Han girls]. *Beijing Da Xue Xue Bao*. 2011;43(3):360–364
30. de Muinich Keizer SM, Mul D. Trends in pubertal development in Europe. *Hum Reprod Update*. 2001;7(3):287–291
31. Pasquet P, Biyong AM, Rikong-Adie H, Befidi-Mengue R, Garba MT, Froment A. Age at menarche and urbanization in Cameroon: current status and secular trends. *Ann Hum Biol*. 1999;26(1):89–97
32. Zhang KH, Song S. Rural–urban migration and urbanization in China: evidence from time-series and cross-section analyses. *China Econ Rev*. 2003;14(4):386–400
33. Sicular T, Ximing Y, Gustafsson B, Shi L. The urban-rural income gap and inequality in China. *Rev Income Wealth*. 2007;53(1):93–126
34. Karapanou O, Papadimitriou A. Determinants of menarche. *Reprod Biol Endocrinol*. 2010;8:115
35. Himes JH, Obarzanek E, Baranowski T, Wilson DM, Rochon J, McClanahan BS. Early sexual maturation, body composition, and obesity in African-American girls. *Obes Res*. 2004;12(suppl):64S–72S
36. Blell M, Pollard TM, Pearce MS. Predictors of age at menarche in the newcastle thousand families study. *J Biosoc Sci*. 2008;40(4):563–575
37. Burt Solorzano CM, McCartney CR. Obesity and the pubertal transition in girls and boys. *Reproduction*. 2010;140(3):399–410
38. Sadrzadeh S, Klip WA, Broekmans FJ, et al; OMEGA Project group. Birth weight and age at menarche in patients with polycystic ovary syndrome or diminished ovarian reserve, in a retrospective cohort. *Hum Reprod*. 2003;18(10):2225–2230
39. Wolff MS, Landrigan PJ. Organochlorine chemicals and children’s health. *J Pediatr*. 2002;140(1):10–13
40. Tchernof A, Després JP. Sex steroid hormones, sex hormone-binding globulin, and obesity in men and women. *Horm Metab Res*. 2000;32(11–12):526–536
41. Cameron JL. Nutritional determinants of puberty. *Nutr Rev*. 1996;54(2 pt 2):S17–S22
42. Rogol AD, Clark PA, Roemmich JN. Growth and pubertal development in children and adolescents: effects of diet and physical activity. *Am J Clin Nutr*. 2000;72(suppl 2):521S–528S
43. Rogers IS, Northstone K, Dunger DB, Cooper AR, Ness AR, Emmett PM. Diet throughout childhood and age at menarche in a contemporary cohort of British girls. *Public Health Nutr*. 2010;13(12):2052–2063
44. Maclure M, Travis LB, Willett W, MacMahon B. A prospective cohort study of nutrient intake and age at menarche. *Am J Clin Nutr*. 1991;54(4):649–656
45. Koprowski C, Ross RK, Mack WJ, Henderson BE, Bernstein L. Diet, body size and menarche in a multiethnic cohort. *Br J Cancer*. 1999;79(11–12):1907–1911
46. Berkey CS, Gardner JD, Frazier AL, Colditz GA. Relation of childhood diet and body size to menarche and adolescent growth in girls. *Am J Epidemiol*. 2000;152(5):446–452
47. Kralj-Cercek L. The influence of foods, body build, and social origin on the age at menarche. *Hum Biol*. 1956;28(4):393–406
48. Blanck HM, Marcus M, Tolbert PE, et al. Age at menarche and tanner stage in girls exposed in utero and postnatally to polybrominated biphenyl. *Epidemiology*. 2000;11(6):641–647
49. McKinney JD, Waller CL. Polychlorinated biphenyls as hormonally active structural analogues. *Environ Health Perspect*. 1994;102(3):290–297

Secular Trend of Age at Menarche in Chinese Adolescents Born From 1973 to 2004

Xin Meng, Suyun Li, Wenhou Duan, Yanxin Sun and Chongqi Jia
Pediatrics 2017;140;

DOI: 10.1542/peds.2017-0085 originally published online July 18, 2017;

Updated Information & Services

including high resolution figures, can be found at:
<http://pediatrics.aappublications.org/content/140/2/e20170085>

References

This article cites 45 articles, 5 of which you can access for free at:
<http://pediatrics.aappublications.org/content/140/2/e20170085#BIBL>

Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):
Endocrinology
http://www.aappublications.org/cgi/collection/endocrinology_sub
Puberty
http://www.aappublications.org/cgi/collection/puberty_sub
Public Health
http://www.aappublications.org/cgi/collection/public_health_sub

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
<http://www.aappublications.org/site/misc/Permissions.xhtml>

Reprints

Information about ordering reprints can be found online:
<http://www.aappublications.org/site/misc/reprints.xhtml>

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®



PEDIATRICS[®]

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Secular Trend of Age at Menarche in Chinese Adolescents Born From 1973 to 2004

Xin Meng, Suyun Li, Wenhou Duan, Yanxin Sun and Chongqi Jia
Pediatrics 2017;140;

DOI: 10.1542/peds.2017-0085 originally published online July 18, 2017;

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/140/2/e20170085>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 345 Park Avenue, Itasca, Illinois, 60143. Copyright © 2017 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN[®]

