

Adolescent Bariatric Surgery and Thiamine Deficiency: What Do We Know So Far?

We read with great interest the recent article by Armstrong-Javors et al¹ published in the December 2016 issue of *Pediatrics* that describes a case of Wernicke encephalopathy (WE) 2 months after Roux-en-Y gastric bypass (RYGB). The authors report a rare case of WE resulting from thiamine deficiency in a 15-year-old white girl who underwent RYGB. According to the article, WE was suspected because of the progressive neurologic symptoms (nystagmus, irritability, ataxia) and frequent vomiting. The aforementioned case is important because it reports not only a rare but also an acute complication after bariatric surgery with notably increased mortality and morbidity. Furthermore, this review revealed that the overwhelming majority of the cases of WE after bariatric surgery occurred during recent years. There are 3 solid reasons for the aforementioned increase.

Firstly, adolescent obesity has dramatically increased worldwide in recent decades, and the prevention strategies for obesity are failing. Because pharmacological and behavioral treatment options have had mainly poor results in the treatment of obesity, bariatric surgery has emerged as an alternative treatment option. Bariatric surgery in adult populations has proved effective with superior outcomes in terms of weight loss and remission of obesity-associated comorbidities; however, long-term results and complications after bariatric surgery in obese adolescents are not yet available. According to Armstrong-Javors et al,¹ there are only 9 definitive adolescent cases of WE after bariatric surgery reported in the literature. Our review of the literature revealed that there is also a dramatic increase of the published cases of WE after bariatric procedures in adult populations during recent years.^{2,3} Because bariatric surgery in pediatric patients has been

used increasingly within the last few years, there is a consequential rise in complications, mainly severe micronutrient deficiency, protein-calorie malnutrition, gastrointestinal obstruction, and pulmonary embolism.

Secondly, RYGB is the most commonly performed procedure in adolescents with obesity in the United States, and the risk of severe micronutrient deficiencies after malabsorptive procedures such as RYGB is significantly higher than restrictive bariatric procedures such as sleeve gastrectomy and laparoscopic adjustable gastric band. The RYGB reduces the absorption of thiamine mainly because it bypasses the duodenum and more than 30 cm of jejunum beyond the ligament of Treitz. Moreover, small intestinal bacterial overgrowth because of altered gut ecology is associated with the development of thiamine deficiency and WE.⁴ Last but not least, the majority of patients undergoing bariatric procedures do not receive adequate and appropriate nutrition supplements, including supplemental vitamin B₁, not only after but also before the operation, and for this reason, these patients are more likely to develop WE.⁵

Overall, bariatric surgery in adolescents should be offered only within authorized adolescent obesity prevention programs provided by interdisciplinary adolescent obesity teams, and only within specialist centers, to minimize detrimental complications and improve outcomes. Furthermore, patients and family should be informed and educated about the risks and benefits, expected complications, potential eating behavior changes, careful nutritional follow-up, and lifelong requirement for multivitamin supplementation.

Antonios Athanasiou
General Surgeon, Department of Surgery, Mercy
University Hospital, Cork, Ireland
E-mail: antwnis_athanasiou@hotmail.com

Michael Spartalis
Cardiologist

Eleftherios Spartalis
General Surgeon

CONFLICT OF INTEREST: The authors have indicated they have no potential conflicts of interest to disclose.

REFERENCES

1. Armstrong-Javors A, Pratt J, Kharasch S. Wernicke encephalopathy in adolescents after bariatric surgery: case report and review. *Pediatrics*. 2016;138(6):e20161039
2. Athanasiou A, Moris D, Davakis S, Spartalis E. Wernicke encephalopathy after restrictive bariatric surgery: a significant increase of a rare complication. *Am Surg*. 2016;82(12):374–375
3. Athanasiou A, Angelou A, Diamantis T. Wernicke's encephalopathy after sleeve gastrectomy. Where do we stand today? A reappraisal. *Surg Obes Relat Dis*. 2014;10(3):563
4. Lakhani SV, Shah HN, Alexander K, Finelli FC, Kirkpatrick JR, Koch TR. Small intestinal bacterial overgrowth and thiamine deficiency after Roux-en-Y gastric bypass surgery in obese patients. *Nutr Res*. 2008;28(5):293–298
5. Inge TH, Krebs NF, Garcia VF, et al. Bariatric surgery for severely overweight adolescents: concerns and recommendations. *Pediatrics*. 2004;114(1):217–223

doi:10.1542/peds.2017-1633A

Authors' Response

Thank you for your thoughtful commentary. Although we agree with most of your discussion, especially your conclusion that adolescent bariatric surgery should be performed in multidisciplinary adolescent centers, there are a few points that we would like to emphasize.

Firstly, most bariatric procedures performed in adolescents in the United States and worldwide today are sleeve gastrectomy (SG) rather than Roux-en-Y gastric bypass (RYGB) procedures. Although there are limited statistical data in adolescents, this transition from bypass to sleeve procedures seems to be mirroring that of the adult population. Although 37% of weight reduction surgeries in 2011 were RYGBs and 18% were SGs, by 2015,

Adolescent Bariatric Surgery and Thiamine Deficiency: What Do We Know So Far?

Antonios Athanasiou, Michael Spartalis and Eleftherios Spartalis
Pediatrics 2017;140;

DOI: 10.1542/peds.2017-1633A originally published online July 31, 2017;

Updated Information & Services

including high resolution figures, can be found at:
<http://pediatrics.aappublications.org/content/140/2/e20171633A>

References

This article cites 5 articles, 2 of which you can access for free at:
<http://pediatrics.aappublications.org/content/140/2/e20171633A#BIBL>

Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):
Administration/Practice Management
http://www.aappublications.org/cgi/collection/administration:practice_management_sub

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
<http://www.aappublications.org/site/misc/Permissions.xhtml>

Reprints

Information about ordering reprints can be found online:
<http://www.aappublications.org/site/misc/reprints.xhtml>

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®



PEDIATRICS[®]

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Adolescent Bariatric Surgery and Thiamine Deficiency: What Do We Know So Far?

Antonios Athanasiou, Michael Spartalis and Eleftherios Spartalis

Pediatrics 2017;140;

DOI: 10.1542/peds.2017-1633A originally published online July 31, 2017;

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/140/2/e20171633A>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 345 Park Avenue, Itasca, Illinois, 60143. Copyright © 2017 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN[®]

