

# Momentary Parental Stress and Food-Related Parenting Practices

Jerica M. Berge, PhD, MPH, LMFT, CFLE,<sup>a</sup> Allan Tate, MPH,<sup>a</sup> Amanda Trofholz, MPH, RD,<sup>a</sup> Angela R. Fertig, PhD,<sup>b</sup> Michael Miner, PhD,<sup>a</sup> Scott Crow, MD,<sup>c,d</sup> Dianne Neumark-Sztainer, PhD, RD<sup>e</sup>

abstract

**BACKGROUND:** Research suggests that stress and depressed mood are associated with food-related parenting practices (ie, parent feeding practices, types of food served at meals). However, current measures of parental stress, depressed mood, and food-related parenting practices are typically survey-based and assessed as static/unchanging characteristics, failing to account for fluctuations across time and context. Identifying momentary factors that influence parent food-related parenting practices will facilitate the development of effective interventions aimed at promoting healthy food-related parenting practices. In this study, we used ecological momentary assessment to examine the association between momentary factors (eg, stress, depressed mood) occurring early in the day and food-related parenting practices at the evening meal.

**METHODS:** Children aged 5 to 7 years and their families ( $N = 150$ ) from 6 racial and/or ethnic groups ( $n = 25$  each African American, Hispanic/Latino, Hmong, American Indian, Somali, and white families) were recruited for this mixed-methods study through primary care clinics.

**RESULTS:** Higher stress and depressed mood earlier in the day predicted pressure-to-eat feeding practices and fewer homemade foods served at meals the same night. Effect modification was found for certain racial and/or ethnic groups with regard to engaging in pressure-to-eat feeding practices (ie, American Indian, Somali) or serving fewer homemade meals (ie, African American, Hispanic/Latino) in the face of high stress or depressed mood.

**CONCLUSIONS:** Clinicians may want to consider discussing with parents the influence stress and depressed mood can have on everyday food-related parenting practices. Additionally, future researchers should consider using real-time interventions to reduce parental stress and depressed mood to promote healthy parent food-related parenting practices.



<sup>a</sup>Division of Epidemiology and Community Health, School of Public Health, Departments of <sup>a</sup>Family Medicine and Community Health and <sup>c</sup>Psychiatry, University of Minnesota Medical School, University of Minnesota, Minneapolis, Minnesota; <sup>b</sup>Medica Research Institute, Minneapolis, Minnesota; and <sup>d</sup>The Emily Program, St Paul, Minnesota

Dr Berge is the principal investigator on the grant and conceptualized the manuscript, assisted with data analysis and interpretation, and wrote all drafts of the manuscript; Mr Tate conducted the data analysis and critically reviewed the manuscript; Ms Trofholz assisted with data acquisition and interpretation of the data and critically reviewed the manuscript; Dr Fertig assisted with the conceptualization of the manuscript and the data analysis and critically reviewed the manuscript; Dr Miner assisted with the data analysis of the manuscript and critically reviewed the manuscript; Dr Crow assisted with the conceptualization of the manuscript and critically reviewed the manuscript; Dr Neumark-Sztainer assisted in conceptualizing the manuscript, contributed to the design of the study, and critically reviewed the manuscript; and all authors gave final approval of this version to be published and agree to be accountable for all aspects of the work regarding the accuracy or integrity of any part of the work.

**DOI:** <https://doi.org/10.1542/peds.2017-2295>

**WHAT'S KNOWN ON THIS SUBJECT:** Stress and depressed mood have been hypothesized to influence food-related parenting practices. However, stress, mood, and parenting practices are often measured as static and/or unchanging characteristics, failing to account for fluctuations across time and context.

**WHAT THIS STUDY ADDS:** In this EMA study, we examined the association between momentary stress and depressed mood and food-related parenting practices. Higher parental stress and depressed mood earlier in the day predicted pressure-to-eat feeding practices and fewer homemade foods served at dinner the same night.

**To cite:** Berge JM, Tate A, Trofholz A, et al. Momentary Parental Stress and Food-Related Parenting Practices. *Pediatrics*. 2017;140(6):e20172295

Previous studies have revealed that food-related parenting practices such as parent feeding practices and healthfulness of foods served at family meals are associated with child weight and weight-related outcomes.<sup>1–12</sup> For example, controlling parent feeding practices such as restriction and pressure-to-eat have been found to be associated with overweight status,<sup>6–8</sup> unhealthy diet quality,<sup>7–9</sup> lower satiety responsiveness,<sup>10,11</sup> and unhealthy weight control behaviors<sup>12</sup> in children. Additionally, researchers have suggested that serving unhealthy foods at family meals (eg, energy dense foods, high-fat foods, sugar-sweetened beverages) is associated with more unhealthy diet quality and overweight status in children.<sup>1–5</sup> Given the potential harmful outcomes associated with controlling parent feeding practices and unhealthy food served at family meals, it is important to find predictors of these food-related parenting practices. Two important factors that may increase a parent's potential to engage in controlling feeding practices or to serve unhealthy foods at family meals are stress and depressed mood.

There is a large body of research that indicates elevated levels of stress contribute to a parent's own unhealthy dietary intake,<sup>13,14</sup> unhealthful food preparation,<sup>13</sup> weight status,<sup>14</sup> and weight control behaviors (eg, emotional eating).<sup>14–17</sup> In addition, previous studies have revealed that depressed mood is associated with a parent's excess weight and unhealthier lifestyle.<sup>18</sup> However, little is known about whether a parent's stress levels or depressed mood are associated with engaging in more or less controlling parent feeding practices with their children or serving more or less unhealthy foods at meals.<sup>19</sup> Of the limited previous studies, results have revealed that maternal depressed mood was

associated with increased risk of engaging in pressure-to-eat feeding practices<sup>20</sup> and that stress was associated with parent controlling feeding practices with children.<sup>19</sup> In addition, the authors of 1 study showed that parent depressed mood was associated with serving less healthy foods to adolescents at family meals.<sup>4</sup> Although the authors of these previous studies suggest that there may be a connection between parental stress and depressed mood and food-related parenting practices, there are limitations with the research. For example, researchers for past studies have measured stress, depressed mood, and food-related parenting practices using self-report or static measures, which are prone to social desirability, lack of specificity, and assume food-related parenting practices do not fluctuate. Additionally, few studies have been focused on populations from diverse ethnic and/or racial and low socioeconomic status backgrounds, which is important given the high prevalence of both obesity and stress in these populations.<sup>21–23</sup>

With this study, we build on and expand the previous research in which researchers examined associations between parental stress and depressed mood and parent food-related parenting practices by using ecological momentary assessment (EMA) of both the exposure variables (ie, parental stress, depressed mood) and outcome variables (ie, parent feeding practices, healthfulness of foods served at dinner). EMA is a powerful technique that uses a smartphone-based Web application to record behaviors and/or ratings of stress, anxiety, hunger, etc. These recordings are made in the moment and in the participant's daily life, allowing for observing behaviors as they unfold in the real world, moment-by-moment, to capture dynamic changes in behavior over time and across contexts.<sup>24–28</sup> This approach avoids

the limitations and biases inherent in retrospective recall. For example, momentary mechanisms preceding food-related parent feeding practices such as stress or depressed mood can be identified by using EMA and then targeted in interventions or discussed with parents during well-child visits to help them manage these feelings to reduce harmful food-related parenting practices. In addition, we include parents from minority, low-income, and immigrant population households, which will allow for understanding how stress and depressed mood operate similarly or differently with regard to food-related parenting practices in diverse families.

The main hypothesis tested in the current study is: high parental stress and depressed mood experienced earlier in the day is associated with controlling parent feeding practices (ie, restriction, pressure-to-eat) and less healthful foods (ie, preprepared foods, fast food) being served at family meals the same evening. In addition, an exploratory hypothesis being tested is: associations found between stress, depressed mood, and parent food-related practices will be modified by race and/or ethnicity. Results from the current study will help inform future interventions regarding momentary influences of food-related parenting practices. For example, understanding whether momentary stress and/or depressed mood are associated with a parent being more likely to engage in controlling feeding practices or to serving unhealthy foods at dinner the same day may inform the development of interventions that can intervene in real time to help parents engage in healthful food-related parenting practices in the face of high stress or depressive mood.

## METHODS

Data for the current study are from *Family Matters*, a National Institutes

of Health–funded study.<sup>29</sup> *Family Matters* is a 5-year incremental (Phase I = 2014–2016; Phase II = 2017–2019), mixed-methods (eg, video-recorded tasks, EMA, interviews, surveys), longitudinal study designed to identify novel risk and protective factors for childhood obesity in the home environments of racially and/or ethnically diverse and primarily low-income children. Phase I included an in-depth, mixed-methods, cross-sectional examination of the family home environment of diverse families ( $n = 150$ ). Phase II will be a longitudinal epidemiologic cohort study with diverse families ( $n = 1200$ ).

Data in the current study are from Phase I of the *Family Matters* study. In Phase I, a mixed-methods analysis of the home environments of children ages 5 to 7 years from 6 racial and/or ethnic groups including African American, Hispanic/Latino, Hmong, American Indian, Somali, and white ( $n = 25$  from each racial and/or ethnic group) was conducted to identify individual, dyadic, and familial risk and protective factors for childhood obesity. The University of Minnesota’s Institutional Review Board Human Subjects Committee approved all protocols used in both phases of the *Family Matters* study.

### Recruitment and Eligibility Criteria

Eligible children ( $n = 150$ ) and their families were recruited from the Minneapolis and St Paul, Minnesota areas between 2015 and 2016 via a letter sent to them by their family physician. Children were eligible to participate in the study if they were between the ages of 5 to 7 years, had a sibling between the ages of 2 to 12 years living in the same home, lived with their parent or primary guardian >50% of the time, shared at least 1 meal/day with the parent or primary guardian, and were from 1 of 6 racial and/or ethnic categories (African American, Hispanic/Latino, Hmong, American Indian, Somali, and

white). The sample was intentionally stratified by race and/or ethnicity and weight status (overweight or obese = BMI  $\geq$ 85th percentile; nonoverweight = BMI >5th percentile and <85th percentile) of the study child to identify potential weight- and/or race and/or ethnic-specific home environment factors related to obesity risk.

### Procedures and Data Collection

A 10-day in-home observation was conducted with each family, including 2 in-home visits and an 8-day direct observational period in between home visits. The observational components included: (1) an interactive observational family task<sup>30,31</sup> using a family board game with activities around family meal planning, meal preparation, and family physical activity to measure family functioning and parenting practices; (2) EMA<sup>24</sup> surveys measuring parental stress, depressed mood, parent feeding practices, food preparation, parent modeling of eating and physical activity, and child dietary intake, physical activity, and sedentary behaviors; (3) child and parent accelerometry; (4) three 24-hour child dietary recalls; (5) a home food inventory; (6) built environment block audit; (7) objectively measured height and weight on all family members; (8) a parent-completed online survey; and (9) a parent interview. All study materials were translated into Spanish, Somali, and Hmong, and bilingual staff were available at all home visits, allowing families to participate in their preferred language. Because of the level of study involvement required, families were compensated with the iPad mini used to record EMA data (~\$300) and additional gift card opportunities (up to \$100) if all elements of the study were completed (eg, 8 complete days of EMA, 3 complete dietary recalls, 4 complete days of accelerometry). This also increased

the likelihood that the equipment was not misused or stolen.

The study sample included diverse families who were equally distributed across the 6 racial and/or ethnic groups recruited in the study (African American, Hispanic/Latino, Hmong, American Indian, Somali, white) (Table 1). Additionally, families were from low-income households, with 70% of families earning <\$35 000 per year. The majority of participants were mothers (91%) who were ~35 years old (mean = 34.5; SD = 7.1) with children aged 6 years (mean = 6.4; SD = 0.08). Over half of the mothers worked full- or part-time, and 61% had a high school diploma or less. Approximately half of the mothers were married, and 64% of households had 2 parents.

### Measures

Multiple daily measures of EMA over 8 days were collected on parents. Standardized EMA data collection protocols from previous studies<sup>24</sup> were used in the study including: (1) signal contingent, (2) event contingent, and (3) end-of-day EMA messaging.<sup>24</sup> iPad minis were provided to parents to enter responses to the EMA surveys during the 8-day observation period. Screen shots of example EMA questions are shown in Fig 1.

Signal contingent recordings were researcher initiated and were used in a stratified random manner so that each parent was prompted via a text message to fill out a survey 4 times a day within a 3-hour time block (eg, 7–10 AM, 11–2 PM, 3–6 PM, 7–10 PM). The surveys expired after 1 hour. The timing of EMA prompts was adjusted for parent shift work and wake times to accommodate parents’ differing life situations. The signal contingent recordings allowed for examining different contexts that occurred day-to-day, moment-by-moment, in families’ lives. Questions asked on the signal contingent surveys included parent modeling of eating,

physical activity and sedentary behavior, parental stress levels and depressed mood, and parent feeding practices.

Event contingent recordings were self-initiated by parents whenever an eating occasion (ie, child and at least 1 other person were eating) occurred. In addition, the parent had to be present for the meal and indicate who was eating the meal together (eg, child, parent, sibling). Parents were asked to fill out information about the type of food served at the meal occasion, what the child actually ate, parent feeding practices, child eating behaviors, pickiness, meal atmosphere, food preparation and planning, and other meal logistics (eg, how long the meal lasted, where it occurred).

The end-of-day recording was completed before sleep to capture any events not reported since the last recording and to get end-of-day measures. All EMA responses were time-stamped. Participants were assigned additional days of EMA if several EMA prompts were missed within a day to obtain a minimum of 8 full days of EMA data with at least 4 complete EMA responses per day (ie, at least 2 signal contingent responses; at least 1 event contingent response; 1 end-of-day response).

All exposure (ie, parental stress, parental depressed mood) and outcome variables (types of food served at dinner, parent feeding practices) are presented in Table 2.

### Statistical Analysis

Descriptive and inferential statistics were used to examine how morning stress and depressed mood related to evening feeding practices and separately to the daily composition of food types (homemade, preprepared, and fast food) served at evening dinner and snacking meal occasions. The 2 predictor variables (stress and depressive symptoms before noon) were analyzed as

**TABLE 1** Family Matters Study Phase I Demographic Characteristics (N = 150)

Participant Characteristics	Primary Caregiver (N = 150), n (%)	Child (N = 150), n (%)
Female sex	137 (91)	71 (47)
Age in years (SD)	34.5 (7.1)	6.4 (0.8)
Adult BMI (BMI percentile child)	30.9 (7.2)	75.9 (23.1)
Weight status		
Nonoverweight	35 (23)	77 (51)
Overweight or obese	115 (77)	73 (49)
Race and/or ethnicity		
Asian (Hmong)	25 (17)	25 (17)
Black or African American	22 (15)	25 (17)
Hispanic/Latino	23 (15)	25 (17)
American Indian or Alaskan native	21 (14)	25 (17)
Somali	25 (17)	25 (17)
White	27 (18)	25 (17)
Multiracial or other	7 (5)	—
Household Characteristics		
Household structure		
1 parent (no other adults)	37 (25)	—
1 parent (w/ other adults)	18 (12)	—
2 parents (no other adults)	78 (52)	—
2 parents (w/ other adults)	17 (11)	—
Primary caregiver marital status		
Married	78 (51)	—
Committed dating relationship or engaged	31 (21)	—
Casually dating	2 (1)	—
Separated or divorced	6 (4)	—
Widowed	1 (1)	—
Single or never married	31 (21)	—
Missing	1 (1)	—
Educational attainment		
Middle school or junior high	15 (10)	—
Some high school	17 (11)	—
High school or GED	60 (40)	—
Vocational, technical, trade certificate program, or associate degree	28 (18)	—
College degree (eg, bachelor, masters, or doctorate)	25 (16)	—
Other	4 (3)	—
Missing	1 (1)	—
Primary caregiver work status		
Working full-time	63 (42)	—
Working part-time	32 (21)	—
Stay-at-home caregiver	25 (17)	—
Currently unemployed, seeking work	18 (12)	—
Not working for pay (unable to work, retired, student, etc)	11 (7)	—
Not applicable	1 (1)	—
Household income		
<\$20 000	50 (33)	—
\$20 000–\$34 999	55 (37)	—
\$35 000–\$49 999	16 (11)	—
\$50 000–\$74 999	12 (8)	—
\$75 000–\$99 999	7 (5)	—
\$100 000 or more	9 (6)	—
Missing	1 (1)	—
Household income source (all that apply)		
Wages from self	85 (57)	—
Wages from other guardian	34 (23)	—
Another family member	14 (10)	—
Unemployment compensation	4 (3)	—
Worker's compensation	0 (0)	—
Social Security	11 (7)	—
Public assistance (eg, MFIP)	34 (23)	—



**TABLE 1** Continued

Participant Characteristics	Primary Caregiver (N = 150), n (%)	Child (N = 150), n (%)
Alimony and/or child support	17 (11)	—
Other sources	21 (14)	—

Percentages may not sum to exactly 100% because of rounding. GED, General Education Diploma; MFIP, Minnesota Family Investment Program; —, not applicable.

continuous random variables. The outcome variables (evening feeding practices and proportion of food types present at those meal occasions) were evaluated with generalized estimating equations with a binomial variance family and logistic link and with Gaussian variance family and identity link respectively. Robust SEs were used and the covariance structure was

initially set to independent. These analyses were performed for the full sample, and separately for each of the 6 race and/or ethnic groups to evaluate differences in subpopulation response to level of stress or mood (ie, effect moderation). All models were adjusted for family level race (regardless of whether the observation day occurred on a weekday or weekend) and the


primary caregiver’s and child’s age, sex, and weight status and/or BMI. All analyses were performed in Stata 13.1SE (StataCorp, College Station, TX).

**RESULTS**


**Stress and Depressed Mood Levels**

Overall, population average stress ( $0.67 \pm 0.95$ ) and depressed mood ( $0.45 \pm 0.85$ ) were low in this sample. Average stress levels were highest among African Americans ( $0.91 \pm 1.1$ ), whites ( $0.79 \pm 0.80$ ), and Hispanics/Latinos ( $0.68 \pm 1.0$ ) and lowest among American Indians ( $0.48 \pm 0.90$ ), Somalis ( $0.55 \pm 0.85$ ), and

**A**



**B**



**FIGURE 1**

Screenshots of signal- and event-contingent EMA survey questions answered by Family Matters participants (names have been changed to protect confidentiality). A, Signal contingent questions. B, Event contingent questions.

**TABLE 2** Exposure and Outcomes Variables Used in the Analysis

Exposure Variables	
Parental stress	Parental stress was measured via signal contingent EMA surveys by using items adapted from the daily health diary. <sup>32</sup> Parents were asked about their current level of stress (ie, how stressed are you feeling right now?), the main source of stress (ie, what is the main source of your stress?; response options, eg, a lot of work to get done at job or school, conflicts or arguments with my spouse or romantic partner), and their perceived ability to cope with stress <sup>32</sup> (ie, right now, how certain do you feel that you can handle all the things that you have to do today?) since they woke up or since the last survey. EMA-reported stress before noon (ie, morning stress) was analyzed as a continuous random variable with Likert scale values ranging from 0 to 4 (0, “not at all”; 1, “a little”; 2, “moderately”; 3, “quite a bit”; 4, “extremely”). Multiple morning stress observations on signal contingent EMA surveys were averaged by participant and by observation day if the stress measure occurred before noon to ensure temporal ordering of the predictor variable.
Parent depressed mood	Parent depressed mood was measured during signal contingent EMA surveys by using an item adapted from Kessler-6 measure of depressive symptoms. <sup>33</sup> Parents were asked about their current level of depressed mood (ie, how sad or depressed are you feeling right now?). EMA-reported depressed mood before noon (ie, morning stress) was also analyzed as a continuous random variable with Likert scale values ranging from 0 to 4 (0, “not at all”; 1, “a little”; 2, “moderately”; 3, “quite a bit”; 4, “extremely”). Temporal ordering was handled for morning depressive symptoms in the same way as the morning stress measure.
Outcome Variables	
Parent feeding practices	Parent restriction and pressure-to-eat parent feeding practices were measured during event contingent (ie, meal occasions) EMA surveys by using 2 items modeled after the child feeding questionnaire. <sup>34</sup> Parent restriction (ie, did you have to make sure [child’s name] didn’t eat too much food at this meal?) and pressure-to-eat (ie, did you have to encourage [child’s name] to eat more food at this meal?) feeding practices at meal occasions were measured as a dichotomous variable (0, “no”; 1, “yes”). Dinner and snacking event contingent meal occasions occurring at 4 PM or later were included for analysis (ie, breakfast and lunch meal occasions were excluded to avoid reverse causation). Evening feeding practices were averaged at the participant and observation day level and treated as a binomial outcome variable ranging between 0 and 1 in all quantitative analyses. If a participant reported pressuring feeding practices at 2 of 3 meal occasions, the binomial outcome variable at the day level would be equal to 0.667 (ie, 2 meal occasions in which the feeding practice was observed divided by the 3 total meals occurring that day after 4 PM at dinner or snacking occasions).
Type of foods served at meals	The type of foods served at meals was assessed during event contingent EMA surveys (ie, which best describes the type of food served at the meal?), adapted from previous survey research questions. Response options included: homemade, preprepared or fast food. Parents could report multiple types of foods served at any one meal occasion. <sup>4,35</sup> The proportion of evening meal occasions (at the participant, observation day level) in which the food type was present was operationalized as a binomial outcome variable in the same way as the parent feeding practices variable (ie, range 0–1). Like the example provided above, if the daily proportion of fast food at evening dinner and snacking meal occasions was equal to 0.667, then two-thirds of 1 day’s evening meal occasions had fast food item present. Parents could report multiple types of foods served at any 1 meal occasion. Lunch meals were excluded to ensure temporal ordering of early parental stress and later-day parent feeding practices and type of foods served at meals.

Hmong ( $0.61 \pm 0.92$ ) participants. Depressed mood showed some similar and some different subpopulation patterns. African Americans ( $0.65 \pm 1.0$ ), Somalis ( $0.55 \pm 0.90$ ), and Hmong ( $0.48 \pm 0.80$ ) had a higher average level of depressed mood, whereas whites ( $0.29 \pm 0.46$ ), American Indians ( $0.36 \pm 0.85$ ), and Hispanics/Latinos ( $0.42 \pm 0.95$ ) reported lower levels of depressed mood. African Americans reported the most stressed and depressed mood and American Indians reported the lowest stress and second lowest depressed mood relative to the rest of the sample.

### Parent Feeding Practices

With regard to parent feeding practices, a 1-unit increase in

stress earlier in the day was associated with 45% greater odds of parents engaging in pressure-to-eat feeding practices at dinner the same night, after controlling for sociodemographics, weight status, and weekday or weekend observation (Table 3). Additionally, a 1-unit increase in depressed mood earlier in the day was associated with 42% greater odds of parents engaging in pressure-to-eat parent feeding practices at dinner the same night (Table 3). There were no significant associations found between parental stress, depressed mood, and parent food restriction. Effect modification results indicated that the relationship between parental stress, depressed mood, and pressure-to-eat feeding practices was strongest for American Indian

parents (odds ratio [OR]: 2.64, 95% confidence interval [CI]: 1.49 to 4.67,  $P = .001$ ) and for Somali parents (OR: 2.02, 95% CI: 1.05 to 3.91,  $P = .036$ ; Table 3).

### Foods Served at Family Meals

With regard to types of foods served at dinner, a 1-unit increase in stress earlier in the day was associated with a decrease in the proportion of dinner meals in which homemade foods were served (mean response:  $-0.05$ , 95% CI:  $-0.08$  to  $-0.01$ ,  $P = .008$ ), after controlling for sociodemographics, BMI, and weekday or weekend observation (Table 4). In addition, a 1-unit increase in depressed mood earlier in the day was negatively associated with serving homemade foods (mean response:  $-0.06$ , 95% CI:  $-0.09$

**TABLE 3** Bivariate and Adjusted Associations Between Morning Stress and Mood Levels With Evening Food Pressuring and Restrictive Feeding Practices (N = 150 Caregivers; N = 1044 Meal Occasions)

Independent Predictor: Stress	OR	95% CI	P	Independent Predictor: Mood	OR	95% CI	P
Full sample adjusted analyses							
Pressure to eat	1.45	(1.08 to 1.94)	.013*	Pressure to eat	1.42	(1.01 to 1.99)	.043*
Restriction	1.26	(0.96 to 1.64)	.091	Restriction	1.24	(0.96 to 1.58)	.096
Stratified by race and/or ethnicity							
African American (N = 25, N = 170 meals)				African American (N = 25, N = 170 meals)			
Pressure to eat	1.46	(0.45 to 4.74)	.531	Pressure to eat	1.05	(0.51 to 2.14)	.894
Restriction	1.02	(0.56 to 1.84)	.951	Restriction	1.04	(0.56 to 1.93)	.892
White (N = 25, N = 182 meals)				White (N = 25, N = 182 meals)			
Pressure to eat	0.90	(0.56 to 1.44)	.661	Pressure to eat	0.49	(0.19 to 1.26)	.140
Restriction	0.47	(0.19 to 1.16)	.100	Restriction	0.81	(0.16 to 4.21)	.807
Hmong (N = 25, N = 185 meals)				Hmong (N = 25, N = 185 meals)			
Pressure to eat	1.14	(0.67 to 1.95)	.633	Pressure to eat	1.09	(0.55 to 2.13)	.811
Restriction	1.42	(0.78 to 2.60)	.253	Restriction	1.27	(0.70 to 2.31)	.438
Hispanic/Latino (N = 25, N = 178 meals)				Hispanic/Latino (N = 25, N = 178 meals)			
Pressure to eat	1.12	(0.76 to 1.63)	.566	Pressure to eat	1.09	(0.61 to 1.97)	.764
Restriction	0.97	(0.60 to 1.56)	.888	Restriction	0.97	(0.48 to 1.94)	.925
American Indian (N = 25, N = 162 meals)				American Indian (N = 25, N = 162 meals)			
Pressure to eat	2.64	(1.49 to 4.67)	.001*	Pressure to eat	2.40	(1.29 to 4.49)	.006*
Restriction	1.19	(0.61 to 2.32)	.614	Restriction	1.24	(0.76 to 2.00)	.390
Somali (N = 25, N = 167 meals)				Somali (N = 25, N = 167 meals)			
Pressure to eat	2.02	(1.05 to 3.91)	.036*	Pressure to eat	1.55	(0.79 to 3.02)	.203
Restriction	1.49	(0.80 to 2.77)	.210	Restriction	1.06	(0.63 to 1.80)	.819

Adjusted models include covariates: primary caregiver and/or child age, sex, and weight status; family race; weekend or weekday observation. Interpretation example: a 1-unit increase in morning stress was associated with 45% greater odds of pressure-to-eat feeding practices (OR: 1.45, 95% CI: 1.08 to 1.94, P = .013) the same night at dinner, after controlling for all other covariates in the adjusted models. Effect modification interpretation: the relationship between stress and pressure-to-eat feeding practices was strongest for American Indian caregivers (OR: 2.64, 95% CI: 1.49 to 4.67, P = .001) and for Somali caregivers (OR: 2.02, 95% CI: 1.05 to 3.91, P = .36).

\* P < .05.

to  $-0.02$ ,  $P = .004$ ) and positively associated with serving preprepared foods (mean response:  $0.05$ , 95% CI:  $0.01$  to  $0.08$ ,  $P = .025$ ) at dinner the same night (Table 4).

Effect modification results indicated that the relationship between parental stress, depressed mood, and the proportion of dinner meals with homemade food was strongest for African American parents and Hispanic/Latino parents. For example, a 1-unit increase in stress earlier in the day was negatively associated with serving homemade foods at dinner meals (mean response:  $-0.10$ , 95% CI:  $-0.16$  to  $-0.04$ ,  $P = .001$ ) and a 1-unit increase in depressed mood earlier in the day was negatively associated with serving homemade food at dinner (mean response:  $-0.09$ , 95% CI:  $-0.16$  to  $-0.02$ ,  $P = .008$ ) and positively associated with serving preprepared foods at dinner (mean response:  $0.10$ , 95% CI:  $0.03$  to

$0.17$ ,  $P = .008$ ) for African American parents (Table 4).

## DISCUSSION

Overall, the results of the current study indicated that parental stress and depressed mood experienced earlier in the day were associated with food-related parenting practices at dinner the same night, with some exceptions. The results of the current study both confirm and extend previous research. The findings in the current study related to parental depressed mood support previous studies revealing that maternal depressed mood is associated with parent pressure-to-eat feeding practices.<sup>20</sup> Findings related to stress extend previous studies in the field whose results reveal that stress earlier in the day is associated with parent engagement in pressure-to-eat feeding practices the same night but not parent restriction. One potential explanation of this

finding is that restriction of foods may require more parental energy to engage in than pressure-to-eat; thus, a parent who is experiencing stress or depressive symptoms may not engage in restriction as readily as pressure-to-eat feeding practices. Additionally, results indicate that certain populations (ie, American Indian, Somali) are at higher risk of engaging in pressure-to-eat feeding practices when experiencing high stress levels or depressed mood.

Results of the current study also extend previous research by showing that high parental stress and depressed mood experienced earlier in the day are associated with a lower prevalence of homemade foods being served at dinner meals and a higher prevalence of preprepared foods being served at dinner meals the same night. Additionally, results reveal that certain racial and/or ethnic groups (ie, African American, Hispanic/Latino) are at higher risk

**TABLE 4** Adjusted Associations of Morning Stress and Mood on the Proportion of Food Categories Served at Snack and Dinner Evening Meal Occasions

Independent Predictor: Stress	Mean	95% CI	P	Independent Predictor: Mood	Mean	95% CI	P
Full sample adjusted analyses							
Fast food	0.01	(-0.01 to 0.03)	.149	Fast food	0.00	(-0.02 to 0.02)	.967
Preprepared foods	0.02	(-0.01 to 0.06)	.182	Preprepared foods	0.05	(0.01 to 0.08)	.025*
Homemade foods	-0.05	(-0.08 to -0.01)	.008*	Homemade foods	-0.06	(-0.09 to -0.02)	.004*
Stratified by race and/or ethnicity							
African American				African American			
Fast food	0.04	(-0.01 to 0.08)	.116	Fast food	-0.02	(-0.08 to 0.04)	.485
Preprepared foods	0.03	(-0.04 to 0.11)	.389	Preprepared foods	0.10	(0.03 to 0.17)	.008*
Homemade foods	-0.10	(-0.16 to -0.04)	.001*	Homemade foods	-0.09	(-0.16 to -0.02)	.008*
White				White			
Fast food	0.01	(-0.02 to 0.04)	.509	Fast food	0.03	(-0.05 to 0.11)	.466
Preprepared foods	0.03	(-0.04 to 0.09)	.427	Preprepared foods	-0.02	(-0.13 to 0.09)	.755
Homemade foods	-0.02	(-0.09 to 0.05)	.555	Homemade foods	0.00	(-0.14 to 0.14)	.967
Hmong				Hmong			
Fast food	-0.01	(-0.03 to 0.01)	.319	Fast food	-0.02	(-0.05 to 0.00)	.080
Preprepared foods	0.04	(-0.03 to 0.10)	.270	Preprepared foods	0.04	(-0.04 to 0.13)	.296
Homemade foods	-0.02	(-0.09 to 0.05)	.527	Homemade foods	-0.02	(-0.11 to 0.07)	.682
Hispanic/Latino				Hispanic/Latino			
Fast food	0.00	(-0.04 to 0.04)	.897	Fast food	-0.01	(-0.04 to 0.02)	.522
Preprepared foods	0.05	(-0.04 to 0.14)	.306	Preprepared foods	0.12	(0.04 to 0.20)	.003*
Homemade foods	-0.07	(-0.17 to 0.03)	.161	Homemade foods	-0.12	(-0.19 to -0.06)	<.001*
American Indian				American Indian			
Fast food	-0.04	(-0.07 to 0.00)	.038*	Fast food	-0.01	(-0.06 to 0.04)	.655
Preprepared foods	0.07	(0.00 to 0.14)	.053	Preprepared foods	0.04	(-0.04 to 0.12)	.344
Homemade foods	-0.04	(-0.11 to 0.02)	.223	Homemade foods	-0.05	(-0.12 to 0.02)	.164
Somali				Somali			
Fast food	0.07	(-0.01 to 0.15)	.069	Fast food	0.04	(-0.03 to 0.12)	.254
Preprepared foods	-0.04	(-0.14 to 0.05)	.342	Preprepared foods	-0.02	(-0.11 to 0.06)	.631
Homemade foods	-0.03	(-0.11 to 0.04)	.381	Homemade foods	-0.02	(-0.09 to 0.05)	.572

Adjusted models include covariates: primary caregiver age and BMI, family race, weekend or weekday observation. Interpretation example: a 1-unit increase in depressed mood earlier in the day was negatively associated with serving homemade foods (mean response: -0.06, 95% CI: -0.09 to -0.02,  $P = .004$ ) but was positively associated with serving preprepared foods the same night at dinner (mean response: 0.05, 95% CI: 0.01 to 0.08,  $P = .025$ ), after controlling for all other covariates in the adjusted models. Effect modification interpretation example: in the African American subpopulation analysis, a 1-unit increase in stress levels earlier in the day was negatively associated with serving homemade foods the same night at dinner (mean response: -0.10, 95% CI: -0.16 to -0.04,  $P = .001$ ).

\*  $P < .05$ .

of serving less homemade food when experiencing high levels of stress or depressive symptoms.

Results from the current study may inform future research and recommendations for health care clinicians that work with children and families. For example, future intervention researchers may want to consider using momentary intervention techniques such as ecological momentary intervention methods to intervene in real time with parents to help them engage in healthful food-related parenting practices in the face of high stress or depressed mood. Additionally, intervention researchers may want to consider tailoring interventions with specific populations (ie, African American, Hispanic/Latino, American

Indian, Somali) to address stress and depressed mood to increase healthful food-related parenting practices.

In addition, results from the current study may be used by health care clinicians to identify specific groups of children who may be at higher risk of experiencing pressure-to-eat feeding practices or for being served less healthy foods at dinner. For example, in many primary care clinics, parents are screened for depressed mood. When parents report high levels of depressed mood, health care clinicians may want to educate parents and/or provide resources for parents regarding food-related parenting practices in the face of high stress and/or depressed mood. Furthermore, health care clinicians may want to consider educating

parents during well-child visits regarding the momentary influences that stress and depressed mood can have on behaviors such as what types of food one serves at dinner or parent feeding practices. Many parents may be unaware that their stress levels or depressed mood could influence what they serve their child for dinner or their own parent feeding practices.

There were both strengths and limitations of the current study. Strengths of the current study include the use of EMA to measure behaviors at multiple time points within and across days over an 8-day period. Additionally, EMA was used to measure both exposure (ie, parental stress and depressed mood) and outcome variables (parent feeding practices, types of food



served at meals) across time and context. Furthermore, the sample included racially and/or ethnically and socioeconomically diverse participants as well as immigrant populations. There were also limitations of the current study; 1 includes the use of items from scales that have not been used with EMA or immigrant populations. Additionally, although the overall sample size was relatively small ( $n = 150$ ), this study was a mixed-methods study that allowed for in-depth measurements of family behaviors that would not have been feasible with a larger sample size. Additionally, because EMA was conducted over an 8-day period, with 4 surveys each day, there are over 1000 data points, which increases the precision of our analyses.

## CONCLUSIONS

Results of the current study indicated that high levels of parental stress and depressed mood earlier in the day predicted pressure-to-eat feeding practices, less homemade food, and more preprepared foods at meals the same night. Additionally, these findings were stronger for some racial and/or ethnic groups. Recommendations for future research include developing interventions using ecological momentary intervention to target momentary factors in real time that influence food-related feeding practices such as stress and depressed mood to promote healthful food-related parenting

practices. Health care providers may also want to use study findings to guide their anticipatory guidance with parents during well-child visits regarding the influence that stress and depressed mood can have on everyday food-related parenting practices. In addition, given the study findings, these messages may need to be tailored to specific groups (eg, African American, Hispanic/Latino, American Indian, Somali).

## ABBREVIATIONS

CI: confidence interval  
EMA: ecological momentary assessment  
OR: odds ratio

Accepted for publication Sep 14, 2017

Address correspondence to Jerica M. Berge, PhD, MPH, LMFT, CFLE, Department of Family Medicine and Community Health, University of Minnesota Medical School, 717 Delaware St SE, Room 425, Minneapolis, MN 55414. E-mail: jberge@umn.edu

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2017 by the American Academy of Pediatrics

**FINANCIAL DISCLOSURE:** The authors have indicated they have no financial relationships relevant to this article to disclose.

**FUNDING:** Supported by grant R01HL126171 from the National Heart, Lung, and Blood Institute (principal investigator: Dr Berge). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Heart, Lung, and Blood Institute or the National Institutes of Health. Funded by the National Institutes of Health (NIH).

**POTENTIAL CONFLICT OF INTEREST:** The authors have indicated they have no potential conflicts of interest to disclose.

## REFERENCES

1. Larson N, Eisenberg ME, Berge JM, Arcan C, Neumark-Sztainer D. Ethnic/racial disparities in adolescents' home food environments and linkages to dietary intake and weight status. *Eat Behav.* 2015;16:43–46
2. Loth KA, MacLehose RF, Larson N, Berge JM, Neumark-Sztainer D. Food availability, modeling and restriction: how are these different aspects of the family eating environment related to adolescent dietary intake? *Appetite.* 2016;96:80–86
3. Cullen KW, Baranowski T, Owens E, Marsh T, Rittenberry L, de Moor C. Availability, accessibility, and preferences for fruit, 100% fruit juice, and vegetables influence children's dietary behavior. *Health Education Behav.* 2003;30(5):615–626
4. Neumark-Sztainer D, MacLehose R, Loth K, Fulkerson JA, Eisenberg ME, Berge J. What's for dinner? Types of food served at family dinner differ across parent and family characteristics. *Public Health Nutr.* 2014;17(1):145–155
5. Trofholz AC, Tate AD, Draxten ML, et al. What's being served for dinner? An exploratory investigation of the associations between the healthfulness of family meals and child dietary intake. *J Acad Nutr Diet.* 2017;117(1):102–109
6. Loth KA, MacLehose RF, Fulkerson JA, Crow S, Neumark-Sztainer D. Food-related parenting practices and adolescent weight status: a population-based study. *Pediatrics.* 2013;131(5). Available at: [www.pediatrics.org/cgi/content/full/131/5/e1443](http://www.pediatrics.org/cgi/content/full/131/5/e1443)
7. Birch LL, Fisher JO. Mothers' child-feeding practices influence daughters' eating and weight. *Am J Clin Nutr.* 2000;71(5):1054–1061
8. Birch LL, Davison KK. Family environmental factors influencing the developing behavioral controls of food intake and childhood overweight. *Pediatr Clin North Am.* 2001;48(4):893–907
9. Fisher JO, Mitchell DC, Smiciklas-Wright H, Birch LL. Parental influences

- on young girls' fruit and vegetable, micronutrient, and fat intakes. *J Am Diet Assoc.* 2002;102(1):58–64
10. Birch LL, Fisher JO, Davison KK. Learning to overeat: maternal use of restrictive feeding practices promotes girls' eating in the absence of hunger. *Am J Clin Nutr.* 2003;78(2):215–220
  11. Fisher JO, Birch LL. Restricting access to palatable foods affects children's behavioral response, food selection, and intake. *Am J Clin Nutr.* 1999;69(6):1264–1272
  12. Loth KA, MacLehose RF, Fulkerson JA, Crow S, Neumark-Sztainer D. Are food restriction and pressure-to-eat parenting practices associated with adolescent disordered eating behaviors? *Int J Eat Disord.* 2014;47(3):310–314
  13. Devine CM, Connors MM, Sobal J, Bisogni CA. Sandwiching it in: spillover of work onto food choices and family roles in low- and moderate-income urban households. *Soc Sci Med.* 2003;56(3):617–630
  14. Wardle J, Steptoe A, Oliver G, Lipsey Z. Stress, dietary restraint and food intake. *J Psychosom Res.* 2000;48(2):195–202
  15. Arnow B, Kenardy J, Agras WS. The Emotional Eating Scale: the development of a measure to assess coping with negative affect by eating. *Int J Eat Disord.* 1995;18(1):79–90
  16. Loth K, van den Berg P, Eisenberg ME, Neumark-Sztainer D. Stressful life events and disordered eating behaviors: findings from Project EAT. *J Adolesc Health.* 2008;43(5):514–516
  17. Lattimore PJ. Stress-induced eating: an alternative method for inducing ego-threatening stress. *Appetite.* 2001;36(2):187–188
  18. Castillo F, Francis L, Wylie-Rosett J, Isasi CR. Depressive symptoms are associated with excess weight and unhealthier lifestyle behaviors in urban adolescents. *Child Obes.* 2014;10(5):400–407
  19. El-Behadli AF, Sharp C, Hughes SO, Obasi EM, Nicklas TA. Maternal depression, stress and feeding styles: towards a framework for theory and research in child obesity. *Br J Nutr.* 2015;113(suppl):S55–S71
  20. Goulding AN, Rosenblum KL, Miller AL, et al. Associations between maternal depressive symptoms and child feeding practices in a cross-sectional study of low-income mothers and their young children. *Int J Behav Nutr Phys Act.* 2014;11:75
  21. Ogden CL, Carroll MD, Kit BK, Flegal KM. Prevalence of obesity and trends in body mass index among US children and adolescents, 1999–2010. *JAMA.* 2012;307(5):483–490
  22. Ogden CL, Carroll MD, Kit BK, Flegal KM. Prevalence of childhood and adult obesity in the United States, 2011–2012. *JAMA.* 2014;311(8):806–814
  23. Ogden CL, Carroll MD, Lawman HG, et al. Trends in obesity prevalence among children and adolescents in the United States, 1988–1994 through 2013–2014. *JAMA.* 2016;315(21):2292–2299
  24. Shiffman S, Stone AA, Hufford MR. Ecological momentary assessment. *Annu Rev Clin Psychol.* 2008;4:1–32
  25. Heron KE, Smyth JM. Ecological momentary interventions: incorporating mobile technology into psychosocial and health behaviour treatments. *Br J Health Psychol.* 2010;15(pt 1):1–39
  26. Dunton GF, Intille SS, Wolch J, Pentz MA. Children's perceptions of physical activity environments captured through ecological momentary assessment: a validation study. *Prev Med.* 2012;55(2):119–121
  27. Dunton GF, Liao Y, Intille SS, Spruijt-Metz D, Pentz M. Investigating children's physical activity and sedentary behavior using ecological momentary assessment with mobile phones. *Obesity (Silver Spring).* 2011;19(6):1205–1212
  28. De Young KP, Lavender JM, Crosby RD, et al. Bidirectional associations between binge eating and restriction in anorexia nervosa. An ecological momentary assessment study. *Appetite.* 2014;83:69–74
  29. Berge JM, Trofholz A, Tate AD, et al. Examining unanswered questions about the home environment and childhood obesity disparities using an incremental, mixed-methods, longitudinal study design: the Family Matters study. *Contemp Clin Trials.* 2017;62:61–76
  30. Melby JN, Conger RD. The Iowa family interaction rating scales: instrument summary. In: Kerig PK, Lindahl KM, eds. *Family Observational Coding Systems: Resources for Systemic Research.* Mahway, NJ: Lawrence Erlbaum Associates; 2001:33–58
  31. Berge JM, Mendenhall TJ, Doherty WJ. Using Community-based Participatory Research (CBPR) to target health disparities in families. *Fam Relat.* 2009;58(4):475–488
  32. Dunton GF, Liao Y, Dzubur E, et al. Investigating within-day and longitudinal effects of maternal stress on children's physical activity, dietary intake, and body composition: protocol for the MATCH study. *Contemp Clin Trials.* 2015;43:142–154
  33. Kessler RC, Green JG, Gruber MJ, et al. Screening for serious mental illness in the general population with the K6 screening scale: results from the WHO World Mental Health (WMH) survey initiative. *Int J Methods Psychiatr Res.* 2010;19(suppl 1):4–22
  34. Birch LL, Fisher JO, Grimm-Thomas K, Markey CN, Sawyer R, Johnson SL. Confirmatory factor analysis of the child feeding questionnaire: a measure of parental attitudes, beliefs and practices about child feeding and obesity proneness. *Appetite.* 2001;36(3):201–210
  35. Boutelle KN, Fulkerson JA, Neumark-Sztainer D, Story M, French SA. Fast food for family meals: relationships with parent and adolescent food intake, home food availability and weight status. *Public Health Nutr.* 2007;10(1):16–23

**Momentary Parental Stress and Food-Related Parenting Practices**  
Jerica M. Berge, Allan Tate, Amanda Trofholz, Angela R. Fertig, Michael Miner,  
Scott Crow and Dianne Neumark-Sztainer  
*Pediatrics* 2017;140;  
DOI: 10.1542/peds.2017-2295 originally published online November 22, 2017;

**Updated Information & Services**

including high resolution figures, can be found at:  
<http://pediatrics.aappublications.org/content/140/6/e20172295>

**References**

This article cites 34 articles, 4 of which you can access for free at:  
<http://pediatrics.aappublications.org/content/140/6/e20172295#BIBL>

**Subspecialty Collections**

This article, along with others on similar topics, appears in the following collection(s):

**Obesity**

[http://www.aappublications.org/cgi/collection/obesity\\_new\\_sub](http://www.aappublications.org/cgi/collection/obesity_new_sub)

**Public Health**

[http://www.aappublications.org/cgi/collection/public\\_health\\_sub](http://www.aappublications.org/cgi/collection/public_health_sub)

**Permissions & Licensing**

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:

<http://www.aappublications.org/site/misc/Permissions.xhtml>

**Reprints**

Information about ordering reprints can be found online:  
<http://www.aappublications.org/site/misc/reprints.xhtml>

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



# PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

## **Momentary Parental Stress and Food-Related Parenting Practices**

Jerica M. Berge, Allan Tate, Amanda Trofholz, Angela R. Fertig, Michael Miner,  
Scott Crow and Dianne Neumark-Sztainer

*Pediatrics* 2017;140;

DOI: 10.1542/peds.2017-2295 originally published online November 22, 2017;

The online version of this article, along with updated information and services, is  
located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/140/6/e20172295>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2017 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

