

Early Child Care and Education: A Key Component of Obesity Prevention in Infancy

Sandra G. Hassink, MD, MS

We are now in the fourth decade of the childhood obesity epidemic, and our increased understanding of the impact of stress, early environments, early feeding, and activity and/or inactivity patterns on obesity risk has focused our attention on our youngest children.¹ During these early years, risk factors for obesity are associated with later obesity and racial and/or ethnic disparities² in childhood, adolescence, and adulthood.³

Pre- and postnatal risk factors for obesity include higher maternal prepregnancy BMI, maternal excess gestational weight gain, gestational diabetes, prenatal tobacco exposure, high infant birth weight, and rapid weight gain in infancy, child care attendance, low strength of maternal-infant relationship, low socioeconomic status, shortened infant sleep, inappropriate bottle use, introduction of solid food before age 4 months, infant antibiotic exposure, sugar sweetened beverage intake, infant feeding patterns high in energy density, and a nighttime feeding pattern. Protective factors include caregiver responsive feeding practices and breastfeeding.²⁻⁷

Recommendations to address these risk factors for obesity by prevention in early child care have been made by the Institute of Medicine,⁴ the US Department of Agriculture,⁸ the American Academy of Pediatrics, and American Public Health Association.⁹ Child care has been the focus for obesity prevention for many reasons:

1. Sixty-one percent (12 500 000) of US preschool-aged children are in some care arrangement other than parental care,¹⁰ and of the 63% of mothers with young children in the labor force, ~80% place their children in nonparental care arrangements for an average of ~40 hours a week.¹¹
2. Children who attend center-based child care can get as much as 67% of their daily dietary intake while in care.¹²
3. Both center- and home-based child care are subject to state and sometimes local or federal regulatory control, representing an opportunity to influence nutrition and feeding practices in a systematic way.
4. Certain child care settings can either promote or reduce obesity risk. Time in home child care during infancy has been shown to be associated with increased BMI z score at age 3 years,¹³ whereas Head Start attendance was associated with a decrease in BMI z score among children who already were overweight or obese.¹⁴
5. Child care centers have been the site of successful multicomponent intervention trials to prevent excess weight gain, particularly for children at higher risk such as those in higher weight categories and children with lower socioeconomic status.¹⁵

In this issue of *Pediatrics*, Benjamin-Neelon et al¹⁶ focused on 10 infant

American Academy of Pediatrics Institute for Healthy Childhood Weight, Wilmington, Delaware

Opinions expressed in these commentaries are those of the author and not necessarily those of the American Academy of Pediatrics or its Committees.

DOI: <https://doi.org/10.1542/peds.2017-2846>

Accepted for publication Aug 25, 2017

Address correspondence to Sandra G. Hassink, MD, MS, American Academy of Pediatrics Institute for Healthy Childhood Weight, 2602 Pennington Dr, Wilmington, DE 19810. E-mail: shassink@aap.net

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2017 by the American Academy of Pediatrics

FINANCIAL DISCLOSURE: The author has indicated she has no financial relationships relevant to this article to disclose.

FUNDING: No external funding.

POTENTIAL CONFLICT OF INTEREST: The author has indicated she has no potential conflicts of interest to disclose.

COMPANION PAPER: A companion to this article can be found online at www.pediatrics.org/cgi/doi/10.1542/peds.2017-2076.

To cite: Hassink SG. Early Child Care and Education: A Key Component of Obesity Prevention in Infancy. *Pediatrics*. 2017;140(6):e20172846

feeding standards from the American Academy of Pediatrics' *Caring for Our Children* guidelines for out-of-home child care programs (2011)⁹ and reviewed progress in states' regulation of early infant feeding practices between 2008 and 2016. Encouragingly, all but 2 states had increased regulations concerning infant feeding in most categories in both center and home care sites.

This improvement in the early childhood day care regulatory environment is welcome and an essential step toward implementation of obesity prevention in the infant population. However, there remain barriers to implementation at the individual child care level, such as addressing a caregiver's cultural beliefs and practices related to child feeding and reported difficulties working with parents to provide consistent messages in the child's environment as well as additional training needed for specific child feeding policies.¹⁷ Ethnic differences have also been found to affect attitudes about feeding policies.¹⁸ On the basis of obesity intervention trials in child care, multilevel interventions that are focused on the child care center environment, policies and practices, staff training, and personal health in addition to child engagement, parental support, and outreach would be a goal for future obesity-prevention interventions and is consistent with obesity prevention recommendations.¹⁵

As we think about implementation of obesity prevention strategies, it is worth looking at where our youngest children spend their time. A Census Bureau report (2011)¹⁸ found that of 61% of the 20 400 000 children <5 years old in child care, 24% were regularly cared for by a grandparent, 18% by their father, 13% were in center care, 7% cared for by other relatives, 11% by nonrelatives, 6% in nursery or preschool, 5% in family day care, and 3% were cared for

by siblings. Children of employed mothers were more likely to be in multiple child care arrangements than children of unemployed mothers (53% and 39%). Children of employed mothers who were in multiple child care arrangements were more likely to be cared for by grandparents, a nonrelative in the child's home, other nonrelative care, or cared for in a nursery, preschool, or family day care than in a day care center. Children of nonemployed mothers in multiple child care arrangements were less likely to be in nursery or preschool and more likely to be taken care of by a nonrelative in the child's home.¹⁹

Given the multiple child care arrangements experienced by preschool-aged children and the fact that the greatest association between being overweight and having obesity is related to care by relatives and nonrelatives in informal care,²⁰ there is a need to develop an integrated, child-centered approach for obesity prevention that can take into consideration the multiple different child care environments and individuals involved in the nutritional well-being of young children.

There are many organizations and providers focused on nutrition education, developing parenting, feeding, and nutrition skills, and increasing access to healthy food and many venues in which parents and families can access this information, including the pediatricians' office, Women, Infants, and Children Food and Nutrition Service program, Supplemental Nutrition Assistance Program education, community parenting programs, Internet sites, and books.

On the basis of our current progress, we are poised to take the next steps in obesity prevention and treatment of our infants. First, we need to take a population approach to building the foundations for parent and caregiver skills, knowledge, and resources for infant feeding across all sectors.

This includes developing simple, public, and consistent messaging about evidence-based strategies for obesity prevention in this age group. It is important to understand individual and cultural barriers to implementing healthy infant feeding practices and address them in ways that empower all caregivers. We need to consider the constellation of individuals and multiple sites that care for and feed infants and create approaches that can more effectively provide outreach and education to grandparents, fathers, and relative and nonrelative caregivers (as well as care givers at child care centers) and develop ways to effectively connect to the infant's family and to each other. Attention to the nutritional needs of infants should become a national priority, essential to ensuring the foundations of health for all children and the health of our population.

REFERENCES

1. Ogden CL, Carroll MD, Kit BK, Flegal KM. Prevalence of childhood and adult obesity in the United States, 2011-2012. *JAMA*. 2014;311(8):806-814
2. Woo Baidal JA, Locks LM, Cheng ER, Blake-Lamb TL, Perkins ME, Taveras EM. Risk factors for childhood obesity in the first 1,000 days: a systematic review. *Am J Prev Med*. 2016;50(6):761-779
3. Singh AS, Mulder C, Twisk JW, van Mechelen W, Chinapaw MJ. Tracking of childhood overweight into adulthood: a systematic review of the literature. *Obes Rev*. 2008;9(5):474-488
4. Institute of Medicine. *Early Childhood Obesity Prevention Policies*. In: Birch LL, Parker L, Burns A, eds. Washington, DC: The National Academies Press; 2011
5. Rose CM, Birch LL, Savage JS. Dietary patterns in infancy are associated with child diet and weight outcomes at 6 years. *Int J Obes*. 2017;41(5):783-788
6. Wallby T, Lagerberg D, Magnusson M. Relationship between breastfeeding and early childhood obesity: results

- of a prospective longitudinal study from birth to 4 years. *Breastfeed Med.* 2017;12:48–53
7. Cheng TS, Loy SL, Toh JY, et al. Predominantly nighttime feeding and weight outcomes in infants. *Am J Clin Nutr.* 2016;104(2):380–388
 8. Food and Nutrition Service, US Department of Agriculture. Child and Adult Care Food Program: meal pattern revisions related to the Healthy, Hunger-Free Kids Act of 2010. Final rule. *Fed Regist.* 2016;81(79):24347–24383
 9. American Academy of Pediatrics; American Public Health Association; National Resource Center for Health and Safety in Child Care and Early Education. *Caring for Our Children: National Health and Safety Performance Standards; Guidelines for Early Care and Education Programs.* 3rd ed. Elk Grove Village, IL: American Academy of Pediatrics; 2011
 10. Mamedova S, Redford J. *Early Childhood Program Participation, From the National Household Education Surveys Program of 2012.* Washington, DC: National Center for Education Statistics, Institute of Education Sciences, United States Department of Education; 2015
 11. Larson N, Ward D, Neelon SB, Story M. What role can child-care settings play in obesity prevention? A review of the evidence and call for research efforts. *J Am Diet Assoc.* 2011;111(9):1343–1362
 12. Radcliffe BC, Cameron CV, Baade PD. Nutrient intakes of young children: implications for long-day child-care nutrition recommendations. *Nutr Diet.* 2002;59(3):187–190
 13. Benjamin SE, Rifas-Shiman SL, Taveras EM, et al. Early child care and adiposity at ages 1 and 3 years. *Pediatrics.* 2009;124(2):555–562
 14. Lumeng JC, Kaciroti N, Sturza J, et al. Changes in body mass index associated with head start participation. *Pediatrics.* 2015;135(2). Available at: www.pediatrics.org/cgi/content/full/135/2/e449
 15. Sisson SB, Krampe M, Anundson K, Castle S. Obesity prevention and obesogenic behavior interventions in child care: a systematic review. *Prev Med.* 2016;87:57–69
 16. Benjamin-Neelon SE, Gonzalez-Nahm S, Grossman E, et al. State variations in infant feeding regulations for child care. *Pediatrics.* 2017;140(6):e20172076
 17. Tovar A, Mena NZ, Risica P, Gorham G, Gans KM. Nutrition and physical activity environments of home-based child care: what Hispanic providers have to say. *Child Obes.* 2015;11(5):521–529
 18. Tovar A, Risica P, Mena N, Lawson E, Ankoma A, Gans KM. An assessment of nutrition practices and attitudes in family child-care homes: implications for policy implementation. *Prev Chronic Dis.* 2015;12:E88
 19. Laughlin L. *Who's Minding the Kids? Child Care Arrangements: Spring 2011. Current Population Reports.* Washington, DC: U.S. Census Bureau; 2013:70–135
 20. Alberdi G, McNamara AE, Lindsay KL, et al. The association between childcare and risk of childhood overweight and obesity in children aged 5 years and under: a systematic review. *Eur J Pediatr.* 2016;175(10):1277–1294

Early Child Care and Education: A Key Component of Obesity Prevention in Infancy

Sandra G. Hassink

Pediatrics 2017;140;

DOI: 10.1542/peds.2017-2846 originally published online November 1, 2017;

Updated Information & Services

including high resolution figures, can be found at:
<http://pediatrics.aappublications.org/content/140/6/e20172846>

References

This article cites 16 articles, 4 of which you can access for free at:
<http://pediatrics.aappublications.org/content/140/6/e20172846#BIBL>

Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):

Nutrition

http://www.aappublications.org/cgi/collection/nutrition_sub

Obesity

http://www.aappublications.org/cgi/collection/obesity_new_sub

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:

<http://www.aappublications.org/site/misc/Permissions.xhtml>

Reprints

Information about ordering reprints can be found online:
<http://www.aappublications.org/site/misc/reprints.xhtml>

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®



PEDIATRICS[®]

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Early Child Care and Education: A Key Component of Obesity Prevention in Infancy

Sandra G. Hassink

Pediatrics 2017;140;

DOI: 10.1542/peds.2017-2846 originally published online November 1, 2017;

The online version of this article, along with updated information and services, is
located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/140/6/e20172846>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 345 Park Avenue, Itasca, Illinois, 60143. Copyright © 2017 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN[®]

