

Postdischarge Interventions to Prevent Pediatric Readmissions: Lost in Translation?

James C. Gay, MD, MMHC

Like it or not, the use of readmission rates to reflect the quality of hospital care has trickled down from the Medicare world to pediatrics.^{1–3} The National Quality Forum, Agency for Healthcare Research and Quality, the Centers for Medicare and Medicaid Services, US News and World Report, and many states use readmission rates as the basis for quality measures or financial penalties.^{4–10} Although the validity of this approach for pediatric hospitalizations is unclear, the negative impact readmissions can have on our patients and their families provides ample incentive for us to seek preventive interventions. But which interventions will actually translate into fewer readmissions?

Improved discharge planning, follow-up telephone calls, and home visits have been shown to reduce readmissions for some adult populations.^{11–19} The importance of successful hospital-to-home transitions is well recognized,^{18,20} and although readmission rates for children are much lower than for adults,^{1,3,21,22} poor discharge planning and follow-up are potential causes for preventable pediatric readmissions.²³ Children with established medical homes seem to have fewer readmissions,²⁴ yet some patients with early postdischarge follow-up visits may be readmitted more often.^{25,26}

Obtaining definitive data on specific interventions to reduce readmissions is challenging. However, in this issue of *Pediatrics*, Auger et al²⁷ have narrowed this knowledge gap with a meticulously

designed randomized control trial of postdischarge home nursing visits. In their study, the authors randomly assigned 1500 children to either a single nurse-led home visit or no visit within 4 days of discharge. Outcome measures included 30-day unplanned emergency department visits and hospital readmissions and assessment of parental feelings and perceptions. In an intention-to-treat analysis, the authors found that overall hospital resource use actually increased in the intervention group (17.8% vs 14.0% in controls). Analyzing strict adherence to the study protocol revealed no statistical difference in health care use between groups. Postdischarge coping scores and time to return to “normal routine” were also similar between groups, although parents in the intervention group did recall more “red flags” at 14 days.

Although readers may conclude that home nursing visits lead to increased pediatric health care use, there may be caveats to these findings. The authors appropriately acknowledge that the nurses conducting the follow-up visits were different from those responsible for the discharge preparations and, thus, did not observe the patients’ clinical status at discharge. Could this have led visiting nurses to recommend further care for more patients than nurses familiar with a child at discharge? Perhaps, and additional studies would be necessary to understand the impact of a different strategy. Alternatively, if the intervention patients were simply

Department of Pediatrics, Vanderbilt University Medical Center, and Monroe Carell Jr. Children’s Hospital at Vanderbilt, Nashville, Tennessee

Opinions expressed in these commentaries are those of the author and not necessarily those of the American Academy of Pediatrics or its Committees.

DOI: <https://doi.org/10.1542/peds.2018-1190>

Accepted for publication Apr 19, 2018

Address correspondence to James C. Gay, MD, MMHC, Department of Pediatrics, Monroe Carell Jr. Children’s Hospital at Vanderbilt, 11204 Doctors’ Office Tower, 2220 Children’s Way, Nashville, TN 37232-9225. E-mail: james.gay@vanderbilt.edu

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2018 by the American Academy of Pediatrics

FINANCIAL DISCLOSURE: The author has indicated he has no financial relationships relevant to this article to disclose.

FUNDING: No external funding.

POTENTIAL CONFLICT OF INTEREST: The author has indicated he has no potential conflicts of interest to disclose.

COMPANION PAPER: A companion to this article can be found online at www.pediatrics.org/cgi/doi/10.1542/peds.2017-3919.

To cite: Gay JC. Postdischarge Interventions to Prevent Pediatric Readmissions: Lost in Translation?. *Pediatrics*. 2018;142(1):e20181190

sicker than control patients, their need for medical attention in the immediate postdischarge period may have been greater. The authors' careful methodology to ensure equal severity of illness of both groups makes this unlikely.

At best, nurse-led follow-up visits were ineffective at decreasing postdischarge hospital resource use for the study population. Readers should note that 30-day readmission rates in the intervention (7.7%) and control (5.5%) groups were similar to the ~6.5% unplanned readmission rate reported for children^{21,22}; thus, the study population likely represented a good cross section of inpatients with common medical conditions. In contrast, readmission rates for children with medical complexity (CMC) or multiple chronic conditions are more than twice the overall rates, and these readmissions may be more amenable to interventions.^{21,28} Nurse-led home visits for CMC may identify postdischarge problems, potentially averting some readmissions,²⁹ and the use of home health nursing services may also reduce readmissions in this population.³⁰ Researchers conducting future studies of postdischarge home nursing visits may well consider CMC as another target population.

In this study, the intervention group used more hospital resources after discharge than control patients. If both groups had similar health care needs, then increased emergency visits and readmissions by the intervention group may not have been necessary to achieve the same clinical outcomes. Although the study was not designed to address this issue, the health outcomes of patients who do and do not use hospital services are equally important when determining the most cost-effective care. Further studies are needed to address this question.

Auger et al²⁷ deserve commendation for their approach to these important

questions. Future studies of different patient groups and interventions should include similarly rigorous methods. I am also left with the notion that some readmissions may be prevented if we just find better ways to care for some patients in nonhospital settings. Personally, I remain skeptical about the value of readmission rates as quality measures for children's hospitals. But preventable readmissions should be prevented, not because it will help hospitals look better than their peers in the ratings or save them from financial penalties but because it's the best thing to do for patients and their families. While we're at it, we should probably extend this to all preventable admissions in the first place. So, let's keep at it, because our patients and their families deserve nothing less.

ABBREVIATION

CMC: children with medical complexity

REFERENCES

1. Jencks SF, Williams MV, Coleman EA. Rehospitalizations among patients in the Medicare fee-for-service program [published correction appears in *N Engl J Med*. 2011;364(16):1582]. *N Engl J Med*. 2009;360(14):1418–1428
2. Jha AK, Orav EJ, Epstein AM. Public reporting of discharge planning and rates of readmissions. *N Engl J Med*. 2009;361(27):2637–2645
3. Berry JG, Toomey SL, Zaslavsky AM, et al. Pediatric readmission prevalence and variability across hospitals [published correction appears in *JAMA*. 2013;309(10):986]. *JAMA*. 2013;309(4):372–380
4. National Quality Forum. All-cause admissions and readmissions measures—final report. Available at: www.qualityforum.org/Publications/2015/04/All-Cause_Admissions_and_Readmissions_Measures_-_Final_Report.aspx. Accessed April 12, 2018
5. Agency for Healthcare Research and Quality. Designing and delivering whole-person transitional care: the hospital guide to reducing Medicaid readmissions. Available at: www.ahrq.gov/professionals/systems/hospital/medicaidreadmitguide. Accessed April 12, 2018
6. Centers for Medicare and Medicaid Services. Readmissions reduction program (HRRP). Available at: www.cms.gov/Medicare/Medicare-Fee-for-Service-Payment/AcuteInpatientPPS/Readmissions-Reduction-Program.html. Accessed April 12, 2018
7. Texas External Quality Review Organization. Potentially preventable readmissions in Texas Medicaid and CHIP programs: measurement period: fiscal year 2013. Available at: <https://hhs.texas.gov/sites/default/files/documents/about-hhs/process-improvement/medicaid-chip-qei/PPR-FY2013.pdf>. Accessed April 12, 2018
8. Illinois Department of Healthcare and Family Services. Report 1: potentially preventable readmissions (PPR) summary report by facility SFY 2013 hospital target and actual readmission rates—revised. Available at: <https://www.illinois.gov/hfs/SiteCollectionDocuments/PPRReport.pdf>. Accessed April 12, 2018
9. Ohio Department of Medicaid. Modernize hospital payments. Available at: www.medicaid.ohio.gov/RESOURCES/ReportsandResearch/ModernizeHospitalPayments.aspx. Accessed April 12, 2018
10. US News and World Report. FAQ: how and why we rank and rate hospitals. Available at: <https://health.usnews.com/health-care/best-hospitals/articles/faq-how-and-why-we-rank-and-rate-hospitals>. Accessed April 10, 2018
11. Naylor MD, Brooten D, Campbell R, et al. Comprehensive discharge planning and home follow-up of hospitalized elders: a randomized clinical trial. *JAMA*. 1999;281(7):613–620
12. Stewart S, Vandenbroek AJ, Pearson S, Horowitz JD. Prolonged beneficial effects of a home-based intervention on unplanned readmissions and mortality among patients with congestive heart failure. *Arch Intern Med*. 1999;159(3):257–261

13. Harrison MJ, Kushner KE, Benzies K, Kimak C, Jacobs P, Mitchell BF. In-home nursing care for women with high-risk pregnancies: outcomes and cost. *Obstet Gynecol.* 2001;97(6):982–987
14. Jack BW, Chetty VK, Anthony D, et al. A reengineered hospital discharge program to decrease rehospitalization: a randomized trial. *Ann Intern Med.* 2009;150(3):178–187
15. Verhaegh KJ, MacNeil-Vroomen JL, Eslami S, Geerlings SE, de Rooij SE, Buurman BM. Transitional care interventions prevent hospital readmissions for adults with chronic illnesses. *Health Aff (Millwood).* 2014;33(9):1531–1539
16. Van Spall HGC, Rahman T, Mytton O, et al. Comparative effectiveness of transitional care services in patients discharged from the hospital with heart failure: a systematic review and network meta-analysis. *Eur J Heart Fail.* 2017;19(11):1427–1443
17. Jackson C, Kasper EW, Williams C, DuBard CA. Incremental benefit of a home visit following discharge for patients with multiple chronic conditions receiving transitional care. *Popul Health Manag.* 2016;19(3):163–170
18. Coleman EA, Parry C, Chalmers S, Min SJ. The care transitions intervention: results of a randomized controlled trial. *Arch Intern Med.* 2006;166(17):1822–1828
19. Harrison PL, Hara PA, Pope JE, Young MC, Rula EY. The impact of postdischarge telephonic follow-up on hospital readmissions. *Popul Health Manag.* 2011;14(1):27–32
20. Berry JG, Blaine K, Rogers J, et al. A framework of pediatric hospital discharge care informed by legislation, research, and practice. *JAMA Pediatr.* 2014;168(10):955–962; quiz 965–966
21. Berry JG, Gay JC, Joynt Maddox K, et al. Age trends in 30 day hospital readmissions: US national retrospective analysis. *BMJ.* 2018;360:k497
22. Gay JC, Agrawal R, Auger KA, et al. Rates and impact of potentially preventable readmissions at children’s hospitals. *J Pediatr.* 2015;166(3):613–619.e5
23. Toomey SL, Peltz A, Loren S, et al. Potentially preventable 30-day hospital readmissions at a children’s hospital. *Pediatrics.* 2016;138(2):e20154182
24. Collier RJ, Klitzner TS, Saenz AA, Lerner CF, Nelson BB, Chung PJ. The medical home and hospital readmissions. *Pediatrics.* 2015;136(6). Available at: www.pediatrics.org/cgi/content/full/136/6/e1550
25. Brittan MS, Sills MR, Fox D, et al. Outpatient follow-up visits and readmission in medically complex children enrolled in Medicaid. *J Pediatr.* 2015;166(4):998–1005.e1
26. Collier RJ, Klitzner TS, Lerner CF, Chung PJ. Predictors of 30-day readmission and association with primary care follow-up plans. *J Pediatr.* 2013;163(4):1027–1033
27. Auger KA, Simmons JM, Tubbs-Cooley HL, et al. Postdischarge nurse home visits and reuse: the hospital to home outcomes trial. *Pediatrics.* 2018;142(1):e20173919
28. Berry JG, Hall DE, Kuo DZ, et al. Hospital utilization and characteristics of patients experiencing recurrent readmissions within children’s hospitals. *JAMA.* 2011;305(7):682–690
29. Wells S, O’Neill M, Rogers J, et al. Nursing-led home visits post-hospitalization for children with medical complexity. *J Pediatr Nurs.* 2017;34:10–16
30. Gay JC, Thurm CW, Hall M, et al. Home health nursing care and hospital use for medically complex children. *Pediatrics.* 2016;138(5):e20160530

Postdischarge Interventions to Prevent Pediatric Readmissions: Lost in Translation?

James C. Gay

Pediatrics 2018;142;

DOI: 10.1542/peds.2018-1190 originally published online June 22, 2018;

Updated Information & Services

including high resolution figures, can be found at:
<http://pediatrics.aappublications.org/content/142/1/e20181190>

References

This article cites 23 articles, 6 of which you can access for free at:
<http://pediatrics.aappublications.org/content/142/1/e20181190#BIBL>

Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):
Hospital Medicine
http://www.aappublications.org/cgi/collection/hospital_medicine_sub
Continuity of Care Transition & Discharge Planning
http://www.aappublications.org/cgi/collection/continuity_of_care_transition_-_discharge_planning_sub

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
<http://www.aappublications.org/site/misc/Permissions.xhtml>

Reprints

Information about ordering reprints can be found online:
<http://www.aappublications.org/site/misc/reprints.xhtml>

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Postdischarge Interventions to Prevent Pediatric Readmissions: Lost in Translation?

James C. Gay

Pediatrics 2018;142;

DOI: 10.1542/peds.2018-1190 originally published online June 22, 2018;

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/142/1/e20181190>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2018 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

