

# “GAPPS” in Patient Safety

Ricardo A. Quinonez, MD,<sup>a</sup> Alan R. Schroeder, MD<sup>b</sup>

Nearly 2 decades after the pervasive issue of iatrogenic harm was thrust into the public sphere by the landmark publication of *To Err is Human* by the National Academy of Medicine (formerly the Institute of Medicine),<sup>1</sup> collective efforts to improve patient safety have become ubiquitous, particularly in the inpatient setting. Although researchers in several pediatric-focused multi-institutional collaboratives have begun to address this issue and have made laudable gains,<sup>2,3</sup> there is concern that patient harm may still be a significant and underestimated reality.<sup>4–6</sup> One potential factor behind underreporting of adverse events (AEs) is the reality that most reporting systems rely on voluntary or passive error reporting.<sup>7</sup> Tools that use “triggers,” including some that are pediatric-specific, are used to actively detect AEs via automated processes and have been found to detect errors at a higher rate than the usual passive methods.<sup>8–10</sup>

In this issue of *Pediatrics*, Stockwell et al<sup>11</sup> report the use of their Global Assessment of Pediatric Patient Safety (GAPPS) trigger tool used to measure trends in AEs over a 5-year period (from January 2007 to December 2012). The authors randomly selected and retrospectively reviewed 3790 individual patient admissions and used GAPPS to filter out AEs from 16 teaching and nonteaching children’s institutions. Their findings are sobering. A total of 414 AEs representing 19.1 out of 1000 patient days were found, with 210 of these AEs deemed as preventable (9.5 out of 1000 patient days). There was no change in the rate of AEs over the 5-year period.

These results are contrary to other more encouraging recent findings. One such large-scale effort, the Children’s Hospitals’ Solutions for Patient Safety (SPS), was used to report significant reductions in hospital-acquired conditions (HACs), presumably as a result of wide-scale adoption of best practice prevention bundles.<sup>2</sup> Several factors may have influenced these discordant findings. First, different hospitals were involved in the 2 efforts, although there was some overlap. Second, the SPS time period was slightly later (2011–2015 instead of 2007–2012). Finally, the measured safety events were different; the types of safety events measured by GAPPS are more inclusive, whereas SPS was specifically focused on HACs. If the latter explanation is indeed correct, it serves as a cautionary reminder that patient safety extends well beyond HACs and that improvements in specific safety events do not necessarily translate to improvements in overall hospital safety.

Even if authors of a more updated analysis of patient safety using GAPPS were to demonstrate more encouraging findings, we must be mindful that, although GAPPS can be used to capture a wide variety of harm, these harms are still discrete, readily evident, and easily quantifiable. GAPPS uses triggers that suggest an AE occurred (eg, naloxone administration).<sup>12</sup> However, children in the hospital setting are exposed to other more insidious harms. The harm from unnecessary care delivered to children may not be as easy to detect, because the effects may be subtle or may only be apparent through long-term outcomes. An unnecessary computed tomographic scan, for

<sup>a</sup>Section of Pediatric Hospital Medicine, Texas Children’s Hospital and Baylor College of Medicine, Houston, Texas; and <sup>b</sup>Department of Pediatrics, College of Medicine, Stanford University, Palo Alto, California

Opinions expressed in these commentaries are those of the authors and not necessarily those of the American Academy of Pediatrics or its Committees.

**DOI:** <https://doi.org/10.1542/peds.2018-0954>

Accepted for publication Apr 30, 2018

Address correspondence to Ricardo A. Quinonez, MD, Section of Pediatric Hospital Medicine, Texas Children’s Hospital, 1102 Bates Ave, FC 1860, Houston, TX 77030. E-mail: raquinon@texaschildrens.org

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2018 by the American Academy of Pediatrics

**FINANCIAL DISCLOSURE:** The authors have indicated they have no financial relationships relevant to this article to disclose.

**FUNDING:** No external funding.

**POTENTIAL CONFLICT OF INTEREST:** The authors have indicated they have no potential conflicts of interest to disclose.

**COMPANION PAPER:** A companion to this article can be found online at [www.pediatrics.org/cgi/doi/10.1542/peds.2017-3360](http://www.pediatrics.org/cgi/doi/10.1542/peds.2017-3360).

**To cite:** Quinonez RA and Schroeder AR. “GAPPS” in Patient Safety. *Pediatrics*. 2018;142(2):e20180954

example, increases the lifetime risk of leukemia<sup>13</sup> and may unleash a prolonged diagnostic cascade when subtle abnormalities are detected, but would not be considered an AE.

Stockwell et al<sup>11</sup> report that children in nonteaching institutions experienced a lower rate of AEs. Given that >70% of hospitalized children may receive care outside of large freestanding children's hospitals,<sup>14</sup> often in a nonteaching setting, this finding has some potential relevance. The authors speculate that differences in complexity and severity of patients may account for this finding. Although an adjusted analysis for chronic conditions revealed that the difference still remained, the authors appropriately recognize that their adjustment cannot fully account for the differences in patient population or severity and that 1 possible explanation for this finding is that children at nonteaching institutions receive safer care.

If hospitalized children are indeed safer at nonteaching institutions, 1 potential explanation is that they simply receive less care. A previous single-center study revealed that a staff-only model for care of children who are medically complex had lower costs and length of stay than a teaching service in which similar children were admitted.<sup>15</sup> Presumably, the lower costs seen in this study were driven, at least in part, by more efficient, less interventional care. Authors of a larger national database study evaluated staff-only visits versus supervised resident-staff emergency department visits.<sup>16</sup> After adjusting for severity and acuity, supervised encounters had an increased number of admissions, longer median emergency department length of stay, and higher rates of advanced imaging. The possibility that supply-sensitive care in larger teaching institutions could drive care that is less safe merits further study.

Stockwell et al's<sup>11</sup> work is a significant addition to the patient safety literature. It adds an important pediatric contribution to a number of publications in which suboptimal results in preventing harm in the hospital setting are suggested.<sup>4–6,17,18</sup> Although focused efforts on HACs may have had some successes, HACs are only a small part of the patient safety story. Moving forward, widespread adoption of trigger tools such as GAPPS that can be used to more sensitively identify AEs may lead to improvement in AE reporting. Other creative solutions such as family reporting of AEs may augment current reporting efforts in hospitalized children as well.<sup>19,20</sup> Whether increased reporting translates to improved overall safety remains unclear.

Even if we are to achieve a meaningful reduction in reportable harm, the potential for unmeasured harm still looms. We must be careful about making the false conclusion that hospital care is safe on the basis of our available metrics. Stockwell et al's<sup>11</sup> results can be used to reinforce the notion that the only dependable way to avoid harm from an intervention is to avoid the intervention in the first place.<sup>17</sup> If a child has an opioid-related respiratory arrest or a life-threatening bleed after a tonsillectomy, do we ever address whether the tonsillectomy was indicated? When a peripherally inserted central catheter becomes infected, how often do we question the necessity of the catheter in the first place? Given that even safely delivered health care can cause harm, we agree with recent calls to label unnecessary care as a reportable safety event.<sup>21</sup> Only by identifying all sources of harm and equally weighing errors of omission with errors of commission can we hope for a future in which hospitals are truly safe.

## ABBREVIATIONS

AE: adverse event  
GAPPS: Global Assessment of Pediatric Patient Safety  
HAC: hospital-acquired condition  
SPS: Solutions for Patient Safety

## REFERENCES

1. Kohn LT, Corrigan JM, Donaldson MS, eds. *To Err is Human: Building a Safer Health System*. 1st ed. Washington, DC: National Academy of Sciences; 2000
2. Lyren A, Brillli RJ, Zieker K, Marino M, Muething S, Sharek PJ. Children's hospitals' Solutions for Patient Safety collaborative impact on hospital-acquired harm. *Pediatrics*. 2017;140(3):e20163494
3. Schaffzin JK, Harte L, Marquette S, et al. Surgical site infection reduction by the Solutions for Patient Safety hospital engagement network. *Pediatrics*. 2015;136(5). Available at: [www.pediatrics.org/cgi/content/full/136/5/e1353](http://www.pediatrics.org/cgi/content/full/136/5/e1353)
4. James JT. A new, evidence-based estimate of patient harms associated with hospital care. *J Patient Saf*. 2013;9(3):122–128
5. Landrigan CP, Parry GJ, Bones CB, Hackbarth AD, Goldmann DA, Sharek PJ. Temporal trends in rates of patient harm resulting from medical care. *N Engl J Med*. 2010;363(22):2124–2134
6. Makary MA, Daniel M. Medical error—the third leading cause of death in the US. *BMJ*. 2016;353:i2139
7. Resar RK, Rozich JD, Classen D. Methodology and rationale for the measurement of harm with trigger tools. *Qual Saf Health Care*. 2003;12(suppl 2):ii39–ii45
8. Hibbert PD, Molloy CJ, Hooper TD, et al. The application of the global trigger tool: a systematic review. *Int J Qual Health Care*. 2016;28(6):640–649
9. Takata GS, Mason W, Taketomo C, Logsdon T, Sharek PJ. Development, testing, and findings of a pediatric-focused trigger tool to identify medication-related harm in US children's hospitals. *Pediatrics*.

- 2008;121(4). Available at: [www.pediatrics.org/cgi/content/full/121/4/e927](http://www.pediatrics.org/cgi/content/full/121/4/e927)
10. Unbeck M, Lindemalm S, Nydert P, et al. Validation of triggers and development of a pediatric trigger tool to identify adverse events. *BMC Health Serv Res*. 2014;14:655
  11. Stockwell DC, Landrigan CP, Toomey SL, et al; GAPPs Study Group. Adverse events in hospitalized pediatric patients. *Pediatrics*. 2018;142(2):e20173360
  12. Landrigan CP, Stockwell D, Toomey SL, et al. Performance of the Global Assessment of Pediatric Patient Safety (GAPPs) tool. *Pediatrics*. 2016;137(6):e20154076
  13. Pearce MS, Salotti JA, Little MP, et al. Radiation exposure from CT scans in childhood and subsequent risk of leukaemia and brain tumours: a retrospective cohort study. *Lancet*. 2012;380(9840):499–505
  14. Leyenaar JK, Ralston SL, Shieh MS, Pekow PS, Mangione-Smith R, Lindenauer PK. Epidemiology of pediatric hospitalizations at general hospitals and freestanding children's hospitals in the United States. *J Hosp Med*. 2016;11(11):743–749
  15. Bekmezian A, Chung PJ, Yazdani S. Staff-only pediatric hospitalist care of patients with medically complex subspecialty conditions in a major teaching hospital. *Arch Pediatr Adolesc Med*. 2008;162(10):975–980
  16. Pitts SR, Morgan SR, Schragger JD, Berger TJ. Emergency department resource use by supervised residents vs attending physicians alone. *JAMA*. 2014;312(22):2394–2400
  17. Schroeder AR, Harris SJ, Newman TB. Safely doing less: a missing component of the patient safety dialogue. *Pediatrics*. 2011;128(6). Available at: [www.pediatrics.org/cgi/content/full/128/6/e1596](http://www.pediatrics.org/cgi/content/full/128/6/e1596)
  18. Sharek PJ, Classen D. The incidence of adverse events and medical error in pediatrics. *Pediatr Clin North Am*. 2006;53(6):1067–1077
  19. Khan A, Coffey M, Litterer KP, et al; and the Patient and Family Centered I-PASS Study Group. Families as partners in hospital error and adverse event surveillance. *JAMA Pediatr*. 2017;171(4):372–381
  20. Khan A, Furtak SL, Melvin P, Rogers JE, Schuster MA, Landrigan CP. Parent-reported errors and adverse events in hospitalized children. *JAMA Pediatr*. 2016;170(4):e154608
  21. Zapata JA, Lai AR, Moriates C. Is excessive resource utilization an adverse event? *JAMA*. 2017;317(8):849–850

## "GAPPS" in Patient Safety

Ricardo A. Quinonez and Alan R. Schroeder

*Pediatrics* 2018;142;

DOI: 10.1542/peds.2018-0954 originally published online July 13, 2018;

### Updated Information & Services

including high resolution figures, can be found at:  
<http://pediatrics.aappublications.org/content/142/2/e20180954>

### References

This article cites 20 articles, 8 of which you can access for free at:  
<http://pediatrics.aappublications.org/content/142/2/e20180954#BIBL>

### Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):  
**Administration/Practice Management**  
[http://www.aappublications.org/cgi/collection/administration:practice\\_management\\_sub](http://www.aappublications.org/cgi/collection/administration:practice_management_sub)  
**Safety**  
[http://www.aappublications.org/cgi/collection/safety\\_sub](http://www.aappublications.org/cgi/collection/safety_sub)  
**Hospital Medicine**  
[http://www.aappublications.org/cgi/collection/hospital\\_medicine\\_sub](http://www.aappublications.org/cgi/collection/hospital_medicine_sub)  
**Patient Education/Patient Safety/Public Education**  
[http://www.aappublications.org/cgi/collection/patient\\_education:patient\\_safety:public\\_education\\_sub](http://www.aappublications.org/cgi/collection/patient_education:patient_safety:public_education_sub)

### Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:  
<http://www.aappublications.org/site/misc/Permissions.xhtml>

### Reprints

Information about ordering reprints can be found online:  
<http://www.aappublications.org/site/misc/reprints.xhtml>

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



# PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

## "GAPPS" in Patient Safety

Ricardo A. Quinonez and Alan R. Schroeder

*Pediatrics* 2018;142;

DOI: 10.1542/peds.2018-0954 originally published online July 13, 2018;

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/142/2/e20180954>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2018 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

