

# Public Health Surveillance of Prenatal Opioid Exposure in Mothers and Infants

Margaret A. Honein, PhD, MPH, Coleen Boyle, PhD, MS hyg, Robert R. Redfield, MD

The US opioid crisis is the public health emergency of our time and requires urgent public health action to monitor and protect the most vulnerable Americans. We have witnessed a startling death toll in 2017 with 70 237 drug overdose deaths in the United States, of which two-thirds involved opioids.<sup>1</sup> The devastating consequences of this epidemic for mothers and infants have received less attention. Increases in opioid use and misuse in pregnancy have paralleled the increases in the general population; at delivery hospitalization, there were 4 times as many women with an opioid use disorder in 2014 compared with 1999.<sup>2</sup> One of the most immediate and visible impacts of the opioid crisis on infants is the drug withdrawal in the newborn period, termed neonatal abstinence syndrome (NAS). On the basis of 2014 data, 1 newborn was diagnosed with NAS every 15 minutes in the United States, totaling about 32 000 infants annually with associated hospital costs estimated at \$563 million.<sup>3</sup>

## INFORMING CLINICAL CARE AND LINKAGE TO SERVICES FOR PREGNANT WOMEN AND INFANTS

Because prevention and treatment of opioid use disorder in pregnancy is important to both maternal and infant health, there is an urgent need to expand prevention efforts and reduce barriers for medication-assisted treatment before, during, and after pregnancy. The stigma of substance exposure during pregnancy and fears of punitive consequences can be obstacles to women receiving the treatment and care they need, and studies with inadequate control of cofactors can exaggerate the specific effects of an individual substance as occurred with prenatal cocaine exposure.<sup>4</sup> Opioid use disorder during pregnancy is a medical condition that requires care and treatment and can have adverse consequences for the infant, much like preeclampsia, which increases the risk of birth defects and other adverse pregnancy outcomes.<sup>5</sup> Current recommendations for pregnant women with opioid use disorder are to avoid withdrawal during pregnancy and be provided appropriate opioid agonist treatment.<sup>6</sup> Increasing and targeting prevention efforts and providing appropriate treatment and

*Centers for Disease Control and Prevention, Atlanta, Georgia*

Dr Honein conceptualized and drafted the initial manuscript and reviewed and revised the manuscript; Drs Boyle and Redfield critically reviewed the manuscript for important intellectual content; and all authors approved the final manuscript as submitted and agree to be accountable for all aspects of the work.

The findings and conclusions in this report are those of the authors and do not necessarily represent the official position of the Centers for Disease Control and Prevention.

**DOI:** <https://doi.org/10.1542/peds.2018-3801>

Accepted for publication Dec 20, 2018

Address correspondence to Margaret A. Honein, PhD, MPH, National Center on Birth Defects and Developmental Disabilities, Centers for Disease Control and Prevention, 4770 Buford Hwy NE, Mailstop S106-3, Atlanta, GA 30341. E-mail: [mrh7@cdc.gov](mailto:mrh7@cdc.gov)

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2019 by the American Academy of Pediatrics

**FINANCIAL DISCLOSURE:** The authors have indicated they have no financial relationships relevant to this article to disclose.

**FUNDING:** All authors completed this work as full-time government employees; no other specific funding support was provided.

**POTENTIAL CONFLICT OF INTEREST:** The authors have indicated they have no potential conflicts of interest to disclose.

**To cite:** Honein MA, Boyle C, Redfield RR. Public Health Surveillance of Prenatal Opioid Exposure in Mothers and Infants. *Pediatrics*. 2019;143(3):e20183801

adequate access to care could improve both maternal and infant health in the midst of this public health crisis.

Public health surveillance is needed to fill critical knowledge gaps and inform clinical guidance for the care of pregnant women with a treated or untreated opioid use disorder and their infants. Updated, evidence-based clinical guidance might help mitigate adverse infant and childhood outcomes by providing treatment that is tailored to the specific prenatal exposures, including both pharmacologic and nonpharmacologic treatments for infants who are exposed. Progress could be monitored by using a cascade of care model (Table 1) with metrics obtained from longitudinal linked public health surveillance.

### UNDERSTANDING THE FULL IMPACT OF PRENATAL OPIOID EXPOSURE ON INFANTS AND CHILDREN

Understanding the effects of prenatal opioid and polysubstance exposure requires assessing the multiple forms and combinations in which exposure can occur. Prenatal opioid exposure can include opioid medications for pain management or treatment of opioid use disorder, misuse of prescription or illicit opioid drugs, and polysubstance exposure to both opioids and other potentially harmful substances. Although NAS is a well-recognized consequence of prenatal opioid exposure, the full impact of

exposure to these psychoactive medications on the developing fetus and particularly the fetal brain is unknown; the effects might extend far beyond diagnoses that are apparent in the first days and weeks of life. Of significant concern is that polysubstance exposure during pregnancy can include alcohol, a known human teratogen. The lessons learned from prenatal alcohol exposure might be relevant for opioids. Although there is no known safe level of alcohol exposure in pregnancy, sustained and heavy use of alcohol in pregnancy can cause fetal alcohol spectrum disorders. Fetal alcohol spectrum disorders can include a range of physical problems (eg, facial features, central nervous system abnormalities) as well as neurologic (eg, executive functioning deficits), behavioral (eg, emotional lability), and learning disabilities, which are lifelong and influenced by timing and extent of exposure in pregnancy.<sup>7</sup> We are concerned that fetal opioid exposure (and fetal polysubstance exposure) might also result in a range of adverse outcomes and that not all will be readily apparent at birth. Prenatal alcohol exposure has been associated with onset of alcohol use disorders when the prenatally exposed children reach adolescence and young adulthood,<sup>8</sup> and it is suggested in animal studies that prenatal alcohol can also increase the risk of addiction to other drugs of abuse.<sup>9</sup> Public health surveillance could help inform whether a similar risk of future

addiction affects children with prenatal exposure to opioids. As was true in evaluating prenatal cocaine exposure, full consideration must be given to the postnatal environment of children with prenatal opioid exposures, which might include social and economic complexities that adversely impact child development.

The full impact of opioid exposure during pregnancy on the health of our nation's children, including the impact on birth defects, is not yet known. The birth prevalence of gastroschisis, a severe defect of the abdominal wall, has been increasing for more than a decade with no clear explanation. Young maternal age is the strongest and most consistent risk factor for gastroschisis. Prescription opioid exposure in early pregnancy has been associated with several structural birth defects, including gastroschisis, and authors of 1 recent study suggested a possible association between preconception opioid use and autism.<sup>10-12</sup> The Centers for Disease Control and Prevention released a report linking county-level opioid prescribing data with population-based birth defects surveillance data of gastroschisis from 20 states that include nearly 50% of US births.<sup>13</sup> Although the authors of the report use an ecological design with appropriate cautions for not overinterpreting this association, the prevalence of gastroschisis was 1.6 times higher in counties with high opioid prescribing rates (5.1 per 10 000 live births; 95% confidence interval: 4.9-5.3) compared with counties with low opioid prescribing rates (3.2 per 10 000 live births; 95% confidence interval: 3.1-3.4). The report sounds an early alarm for the need to increase our public health surveillance on the full range of fetal, infant, and childhood outcomes potentially related to these exposures. More research is needed to better understand the possible role of opioid exposure or related substance

**TABLE 1** Proposed Cascade of Care for Prenatal Opioid Exposure and the Impact on Mothers and Infants

Cascade of Care
Reproductive-aged women with opioid use disorder at risk for pregnancy
Pregnant women with opioid use disorder
Pregnant women with opioid use disorder linked to treatment
Pregnant women with opioid use disorder in treatment throughout pregnancy and postpartum period
Women with opioid use disorder in a previous pregnancy who are receiving appropriate treatment and care preconceptionally
Infants with prenatal opioid exposure receiving appropriate neonatal assessment and care
Infants with prenatal opioid exposure linked to early intervention services
Infants with prenatal opioid exposure monitored for behavioral and learning disabilities

exposure on the occurrence of gastroschisis and other birth defects and the underlying mechanisms for these potential effects. Areas of the United States with high opioid use might need to plan for services for infants and children impacted with a broader range of adverse outcomes, far beyond the impact of NAS. Current surveillance efforts are limited to monitoring prenatal opioid use, opioid use disorder at delivery, and diagnosis of NAS; these efforts lack information on the full range of outcomes that could be obtained from longitudinal mother-infant linked surveillance approaches.

### A PUBLIC HEALTH CALL TO ACTION

There is an urgent need to build the evidence base and define the full spectrum of adverse infant and childhood outcomes associated with sustained prenatal opioid exposure and monitor the linkage to needed care and services. Longitudinal linked surveillance of mother-infant pairs with follow-up into childhood will help clarify the physical and developmental health outcomes linked to prenatal opioid exposure and help target effective programs and services to mitigate adverse effects of the substance exposure. Some preliminary data suggest that infants with a diagnosis of NAS are more likely to need special education services in preschool and early school age than infants without this diagnosis, but better data are needed to clarify the nature of the developmental problems and how they might relate to in utero exposures to opioids and other substances as well as the postnatal environment.<sup>14</sup> The Centers for Disease Control and Prevention collaborated with clinical and public health partners to launch mother-infant linked surveillance during the Zika virus emergency to rapidly understand the impact of infection during pregnancy and translate findings to action in the form of

clinical guidance.<sup>15</sup> We are committed to building on this model to continue to address the health threats to and needs of mothers and infants through a network of public health surveillance.

### ABBREVIATION

NAS: neonatal abstinence syndrome

### REFERENCES

1. Hedegaard H, Miniño AM, Warner M. *Drug Overdose Deaths in the United States, 1999–2017. NCHS Data Brief, No 329.* Hyattsville, MD: National Center for Health Statistics; 2018
2. Haight SC, Ko JY, Tong VT, Bohm MK, Callaghan WM. Opioid use disorder documented at delivery hospitalization - United States, 1999-2014. *MMWR Morb Mortal Wkly Rep.* 2018;67(31):845–849
3. Winkelman TNA, Villapiano N, Kozhimannil KB, Davis MM, Patrick SW. Incidence and costs of neonatal abstinence syndrome among infants with Medicaid: 2004-2014. *Pediatrics.* 2018;141(4):e20173520
4. Frank DA, Augustyn M, Knight WG, Pell T, Zuckerman B. Growth, development, and behavior in early childhood following prenatal cocaine exposure: a systematic review. *JAMA.* 2001; 285(12):1613–1625
5. Gabbay-Benziv R, Reece EA, Wang F, Yang P. Birth defects in pregestational diabetes: defect range, glycemic threshold and pathogenesis. *World J Diabetes.* 2015;6(3):481–488
6. The American College of Obstetricians and Gynecologists. ACOG committee opinion: opioid use and opioid use disorder in pregnancy. Available at: <https://www.acog.org/Clinical-Guidance-and-Publications/Committee-Opinions/Committee-on-Obstetric-Practice/Opioid-Use-and-Opioid-Use-Disorder-in-Pregnancy>. Accessed December 30, 2018
7. Senturias Y, Asamoah A. Fetal alcohol spectrum disorders: guidance for recognition, diagnosis, differential diagnosis and referral. *Curr Probl*

*Pediatr Adolesc Health Care.* 2014;44(4): 88–95

8. Alati R, Al Mamun A, Williams GM, O’Callaghan M, Najman JM, Bor W. In utero alcohol exposure and prediction of alcohol disorders in early adulthood: a birth cohort study. *Arch Gen Psychiatry.* 2006;63(9): 1009–1016
9. Wang R, Shen YL, Hausknecht KA, et al. Prenatal ethanol exposure increases risk of psychostimulant addiction. *Behav Brain Res.* 2019;356:51–61
10. Rubenstein E, Young JC, Croen LA, et al. Brief report: maternal opioid prescription from preconception through pregnancy and the odds of autism spectrum disorder and autism features in children [published online ahead of print August 21, 2018] [published correction appears in *J Autism Dev Disord.* 2018]. *J Autism Dev Disord.* doi:doi:10.1007/s10803-018-3721-8
11. Broussard CS, Rasmussen SA, Reefhuis J, et al; National Birth Defects Prevention Study. Maternal treatment with opioid analgesics and risk for birth defects. *Am J Obstet Gynecol.* 2011;204(4):314.e1–314.e11
12. Lind JN, Interrante JD, Ailes EC, et al. Maternal use of opioids during pregnancy and congenital malformations: a systematic review. *Pediatrics.* 2017;139(6):e20164131
13. Short TD, Stallings EB, Isenburg J, et al. Gastroschisis Trends and Ecologic Link to Opioid Prescription Rates — United States, 2006–2015. *MMWR Morb Mortal Wkly.* In press 2019. Available at: [https://www.cdc.gov/mmwr/volumes/68/wr/mm6802a2.htm?s\\_cid=mm6802a2\\_w](https://www.cdc.gov/mmwr/volumes/68/wr/mm6802a2.htm?s_cid=mm6802a2_w)
14. Fill MA, Miller AM, Wilkinson RH, et al. Educational disabilities among children born with neonatal abstinence syndrome. *Pediatrics.* 2018;142(3): e20180562
15. Rice ME, Galang RR, Roth NM, et al. Vital signs: Zika-associated birth defects and neurodevelopmental abnormalities possibly associated with congenital Zika virus infection - U.S. territories and freely associated states, 2018. *MMWR Morb Mortal Wkly Rep.* 2018;67(31): 858–867

## Public Health Surveillance of Prenatal Opioid Exposure in Mothers and Infants

Margaret A. Honein, Coleen Boyle and Robert R. Redfield

*Pediatrics* 2019;143;

DOI: 10.1542/peds.2018-3801 originally published online January 17, 2019;

<b>Updated Information &amp; Services</b>	including high resolution figures, can be found at: <a href="http://pediatrics.aappublications.org/content/143/3/e20183801">http://pediatrics.aappublications.org/content/143/3/e20183801</a>
<b>References</b>	This article cites 12 articles, 3 of which you can access for free at: <a href="http://pediatrics.aappublications.org/content/143/3/e20183801#BIBL">http://pediatrics.aappublications.org/content/143/3/e20183801#BIBL</a>
<b>Subspecialty Collections</b>	This article, along with others on similar topics, appears in the following collection(s): <b>Fetus/Newborn Infant</b> <a href="http://www.aappublications.org/cgi/collection/fetus:newborn_infant_sub">http://www.aappublications.org/cgi/collection/fetus:newborn_infant_sub</a> <b>Birth Defects</b> <a href="http://www.aappublications.org/cgi/collection/birth_defects_sub">http://www.aappublications.org/cgi/collection/birth_defects_sub</a> <b>Gynecology</b> <a href="http://www.aappublications.org/cgi/collection/gynecology_sub">http://www.aappublications.org/cgi/collection/gynecology_sub</a> <b>Maternal and Fetal Medicine</b> <a href="http://www.aappublications.org/cgi/collection/maternal_fetal_medicine_sub">http://www.aappublications.org/cgi/collection/maternal_fetal_medicine_sub</a>
<b>Permissions &amp; Licensing</b>	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: <a href="http://www.aappublications.org/site/misc/Permissions.xhtml">http://www.aappublications.org/site/misc/Permissions.xhtml</a>
<b>Reprints</b>	Information about ordering reprints can be found online: <a href="http://www.aappublications.org/site/misc/reprints.xhtml">http://www.aappublications.org/site/misc/reprints.xhtml</a>

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



# PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

## **Public Health Surveillance of Prenatal Opioid Exposure in Mothers and Infants**

Margaret A. Honein, Coleen Boyle and Robert R. Redfield

*Pediatrics* 2019;143;

DOI: 10.1542/peds.2018-3801 originally published online January 17, 2019;

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/143/3/e20183801>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2019 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

