

The Pediatrician's Role in Eliminating Racial and Ethnic Disparities in Sleep-Related Infant Deaths

Jeffrey D. Colvin, MD, JD,^a Rachel Y. Moon, MD^b

The infant mortality rate in the United States ranks 33rd among countries belonging to the Organization for Economic Cooperation and Development,¹ and there are persistent racial and ethnic disparities, with non-Hispanic black infants dying at a rate 2 to 3 times that of non-Hispanic white infants.² In this issue of *Pediatrics*, Hirai et al³ provide insight into racial and ethnic differences in infant sleep practices that have important implications for racial and ethnic disparities for sleep-related infant deaths, including sudden infant death syndrome and accidental suffocation and strangulation in bed, which are the third leading cause of infant mortality in the United States.

Hirai et al³ leveraged recently added questions to the Centers for Disease Control and Prevention's Pregnancy Risk Assessment Monitoring System (PRAMS) to examine differences in sleep position, items in the sleep environment, bed-sharing, and room-sharing by race and ethnicity. Forty-seven states currently partner with the Centers for Disease Control and Prevention to administer the PRAMS survey to a randomly selected subpopulation of women in their state who have recently delivered. PRAMS previously only included questions on the infant's sleep position. States began to include questions about bed-sharing and items in the sleep environment in 2009. In 2016, those questions, along with additional survey items about room-sharing and safe sleep anticipatory

guidance from health care providers, were included for all participating states. This allowed the authors to more comprehensively examine the sleep environment of a representative population of infants from a majority of states.

They found a mixture of good and bad news. First, the bad news: one-third of non-Hispanic black infants were not usually placed on their back to sleep; 75% were not solely placed to sleep in a crib, bassinet, or portable crib; and two-thirds had soft objects in the sleep environment. The authors also found that many mothers reported that they never received the full range of safe sleep anticipatory guidance from their health care provider. Approximately 15% of mothers reported never being told to place their infant in a crib, bassinet, or portable crib to sleep and remove all soft objects from the sleep environment; fewer than 50% were advised to room share without bed-sharing. Even if this anticipatory guidance was given but not remembered by the mothers, these numbers should certainly alert our profession to the need to repeatedly discuss safe sleep. Because there are multiple opportunities to discuss safe sleep in the newborn nursery and at the frequent well-child visits and weight checks in early infancy, this is one area in which we can make a difference in infant mortality.

The authors also had good news. They found that parents listen to their health care providers' advice on safe sleep for

^aDepartment of Pediatrics, Children's Mercy Kansas City and School of Medicine, University of Missouri-Kansas City, Kansas City, Missouri; and ^bDepartment of Pediatrics, School of Medicine, University of Virginia, Charlottesville, Virginia

Opinions expressed in these commentaries are those of the authors and not necessarily those of the American Academy of Pediatrics or its Committees.

DOI: <https://doi.org/10.1542/peds.2019-2310>

Accepted for publication Aug 9, 2019

Address correspondence to Rachel Y. Moon, MD, Department of Pediatrics, University of Virginia School of Medicine, PO Box 800386, Charlottesville, VA 22908. E-mail: rymoon@virginia.edu

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2019 by the American Academy of Pediatrics

FINANCIAL DISCLOSURE: The authors have indicated they have no financial relationships relevant to this article to disclose.

FUNDING: No external funding.

POTENTIAL CONFLICT OF INTEREST: The authors have indicated they have no potential conflicts of interest to disclose.

COMPANION PAPER: A companion to this article can be found online at www.pediatrics.org/cgi/doi/10.1542/peds.2019-1286.

To cite: Colvin JD and Moon RY. The Pediatrician's Role in Eliminating Racial and Ethnic Disparities in Sleep-Related Infant Deaths. *Pediatrics*. 2019;144(5):e20192310

their infant. Consistent with previous, smaller studies,⁴⁻⁷ mothers who received safe sleep anticipatory guidance from their health care provider were 10% to 30% more likely to place their infant on their back for sleep; place their infant in a crib, bassinet, or portable playpen; room share without bed-sharing; and have no soft objects in the sleep environment. This makes the provision of repeated safe sleep anticipatory guidance all the more important. The most effective guidance will also likely specifically address the primary drivers of parent behavior: perceptions of infant comfort and safety, the mistaken belief that their infant is not at risk, and misplaced confidence in parental vigilance to prevent a sleep-related death.^{4,5,8-12} Unfortunately, when parents receive inconsistent advice from health care providers, they are more likely to disregard the guidance of all health care providers, instead relying on the advice of friends and family.^{7,11} This is particularly important because sleep-deprived parents may make decisions about infant sleep position, location, and bedding simply on the basis of their desire for the infant to sleep longer.^{4,5,8-12} Thus, consistent, evidence-based infant sleep recommendations from all health care providers will be important in eliminating disparities in sleep practices and sleep-related infant mortality.

However, although evidence-based guidance on infant safe sleep from health care providers is essential in eliminating these disparities, it likely will not be sufficient. In previous research, it has been found that the influence of family and friends and the overall weight of social norms are incredibly important.^{7,13,14} To overcome those influences, repeated, consistent messaging after the infant

leaves the hospital and clinic will be needed.¹⁵ Furthermore, racial and ethnic health disparities cannot be separated from socioeconomic health disparities in infant mortality. This will require the voices of pediatricians outside of the examination room, as well. The advocacy of pediatricians is essential to advance policies, such as paid parental leave and home visiting programs, to support families with infants so that parents and other child care providers get the support, respite, and consistent health messaging needed to eliminate racial and ethnic disparities in infant sleep-related deaths in the United States.

ABBREVIATION

PRAMS: Pregnancy Risk Assessment Monitoring System

REFERENCES

1. Organization for Economic Cooperation and Development. Infant mortality rates. 2019. Available at: <https://data.oecd.org/healthstat/infant-mortality-rates.htm>. Accessed July 8, 2019
2. Mathews TJ, Driscoll AK. *Trends in Infant Mortality in the United States, 2005-2014*. Washington, DC: National Center for Health Statistics; 2017
3. Hirai AH, Kortsmid K, Kaplan L, et al. Prevalence and factors associated with safe infant sleep practices. *Pediatrics*. 2019;144(5):e20191286
4. Colson ER, Geller NL, Heeren T, Corwin MJ. Factors associated with choice of infant sleep position. *Pediatrics*. 2017; 140(3):e20170596
5. Colson ER, Rybin D, Smith LA, et al. Trends and factors associated with infant sleeping position: the national infant sleep position study, 1993-2007. *Arch Pediatr Adolesc Med*. 2009;163(12): 1122-1128

6. Robida D, Moon RY. Factors influencing infant sleep position: decisions do not differ by SES in African-American families. *Arch Dis Child*. 2012;97(10): 900-905
7. Moon RY, Mathews A, Oden R, Carlin R. A qualitative analysis of how mothers' social networks are established and used to make infant care decisions. *Clin Pediatr (Phila)*. 2019;58(9):985-992
8. Ajao TI, Oden RP, Joyner BL, Moon RY. Decisions of black parents about infant bedding and sleep surfaces: a qualitative study. *Pediatrics*. 2011; 128(3):494-502
9. Herman S, Adkins M, Moon RY. Knowledge and beliefs of African-American and American Indian parents and supporters about infant safe sleep. *J Community Health*. 2015;40(1):12-19
10. Joyner BL, Oden RP, Ajao TI, Moon RY. Where should my baby sleep: a qualitative study of African American infant sleep location decisions. *J Natl Med Assoc*. 2010;102(10):881-889
11. Moon RY, Oden RP, Joyner BL, Ajao TI. Qualitative analysis of beliefs and perceptions about sudden infant death syndrome in African-American mothers: implications for safe sleep recommendations. *J Pediatr*. 2010; 157(1):92-97.e2
12. Oden RP, Joyner BL, Ajao TI, Moon RY. Factors influencing African American mothers' decisions about sleep position: a qualitative study. *J Natl Med Assoc*. 2010;102(10):870-872, 875-880
13. Moon RY, Carlin RF, Cornwell B, et al. Implications of mothers' social networks for risky infant sleep practices. *J Pediatr*. 2019;212: 151-158.e2
14. Moon RY, Corwin MJ, Kerr S, et al. Mediators of improved adherence to infant safe sleep using a mobile health intervention. *Pediatrics*. 2019;143(5): e20182799
15. Moon RY, Hauck FR, Colson ER, et al. The effect of nursing quality improvement and mobile health interventions on infant sleep practices: a randomized clinical trial. *JAMA*. 2017;318(4):351-359

The Pediatrician's Role in Eliminating Racial and Ethnic Disparities in Sleep-Related Infant Deaths

Jeffrey D. Colvin and Rachel Y. Moon

Pediatrics 2019;144;

DOI: 10.1542/peds.2019-2310 originally published online October 21, 2019;

Updated Information & Services	including high resolution figures, can be found at: http://pediatrics.aappublications.org/content/144/5/e20192310
References	This article cites 13 articles, 5 of which you can access for free at: http://pediatrics.aappublications.org/content/144/5/e20192310#BIBL
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Fetus/Newborn Infant http://www.aappublications.org/cgi/collection/fetus:newborn_infant_sub SIDS http://www.aappublications.org/cgi/collection/sids_sub
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.aappublications.org/site/misc/Permissions.xhtml
Reprints	Information about ordering reprints can be found online: http://www.aappublications.org/site/misc/reprints.xhtml

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®



PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

The Pediatrician's Role in Eliminating Racial and Ethnic Disparities in Sleep-Related Infant Deaths

Jeffrey D. Colvin and Rachel Y. Moon

Pediatrics 2019;144;

DOI: 10.1542/peds.2019-2310 originally published online October 21, 2019;

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/144/5/e20192310>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 345 Park Avenue, Itasca, Illinois, 60143. Copyright © 2019 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®

