

Skin-to-Skin Care and COVID-19

Carolyn Boscia, MD^a

As an internal medicine and pediatric hospitalist in the home stretch of my third pregnancy, my daily focus is on squeezing into scrubs, following strict infection control procedures, and keeping up a good pace and extra hydration on rounds. But third trimester brings some broader concerns during this first coronavirus disease 2019 (COVID-19) season, and as I follow the medical literature, I find myself wondering what would happen if I were to be infected when I deliver my baby? How would I weigh the risk of transmitting the virus against the known costs of separation from my infant during the first days of life? And in a rapidly evolving clinical landscape, how should I as a physician counsel families asking similar questions?

Numbers from New York suggest that nearly 90% of women positive for COVID-10 who deliver in the hospital setting are asymptomatic at the time of presentation.¹ Given this and the fact that masks are not tolerable for most women during the second stage of labor, many hospitals around the country are moving toward universal COVID-19 screening on labor and delivery units to protect health care workers and patients alike. For women who test positive for COVID-19, current American Academy of Pediatrics guidelines recommend physical separation from their infant when space allows, unless they choose rooming-in despite being counseled on risk.² On the other hand, the World Health Organization's recommendations for women positive for COVID-19 encourage breastfeeding initiation within an hour of birth and routine newborn care with added emphasis on respiratory and hand hygiene.³ As clinicians, we are tasked with understanding the data behind these guidelines and translating them into family-centered practice.

The risk of COVID-19 disease in newborns due to maternal transmission is undefined but appears to be low. In China, a country with >80 000 symptomatic patients with COVID-19, there have been only 6 reported cases in infants born to COVID-infected women, and all 6 made a full recovery without requiring intubation.⁴ China adopted an early policy of mother-infant separation, and we cannot know what these numbers would look like had they not done so. In the United States, there have been 2 reported infant (not newborn) fatalities associated with a positive COVID-19 polymerase chain reaction result, although neither has a formal cause of death yet declared.^{5,6} There has been no clear evidence of vertical

^aThe University of Vermont Medical Center and Departments of Family Medicine, Pediatrics, and Internal Medicine, Larner College of Medicine, The University of Vermont, Burlington, Vermont

Dr Boscia drafted, reviewed, and revised the manuscript; she approved the final manuscript as submitted and agrees to be accountable for all aspects of the work.

DOI: <https://doi.org/10.1542/peds.2020-1836>

Accepted for publication May 15, 2020

Address correspondence to Carolyn Boscia, MD, Pediatric Hospitalist Division, Smith 5, The University of Vermont Medical Center, 111 Colchester Ave, Burlington, VT 05401. E-mail: carolyn.boscia@uvmhealth.org

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2020 by the American Academy of Pediatrics

FINANCIAL DISCLOSURE: The author has indicated she has no financial relationships relevant to this article to disclose.

FUNDING: No external funding.

POTENTIAL CONFLICT OF INTEREST: The author has indicated she has no potential conflicts of interest to disclose.

To cite: Boscia C. Skin-to-Skin Care and COVID-19. *Pediatrics*. 2020;146(2):e20201836

transmission. In the severe acute respiratory syndrome coronavirus outbreak, infants of infected mothers did not develop infection in the neonatal period, although some were born prematurely or small for gestational age because of severe maternal illness during pregnancy.⁷

There are several reasons why neonates could be relatively protected from infection. It is suggested in preliminary data that infants born to mothers positive for COVID-19 may benefit from some degree of passive immune protection at birth, with transplacental immunoglobulin G acting as a natural form of convalescent plasma transfusion.⁸ Thus far, there has been no COVID-19 viral shedding detected in breast milk of infected women, and although we do not yet know whether breastfeeding is protective in this setting, it has known immune benefits in other viral respiratory infections.^{9,10} Even immune system immaturity may be protective for neonates because much COVID-19 morbidity in older patients appears to be driven by the domino effect of cytokine storm.¹¹

Still, because data are limited, separation for mothers positive for COVID-19 and their infants may seem the safest approach. But we must weigh the potential benefits of isolation against the real costs, for both women and infants, of losing postpartum contact. Skin-to-skin contact in the first hours of life is associated with reduced postpartum hemorrhage risk, decreased rates of postpartum depression and anxiety, and increased odds of successful breastfeeding.¹²⁻¹⁴ Breastfeeding is far easier to establish with an oxytocin-inducing newborn than with a cold plastic pump and has myriad health benefits ranging from decreased risks of breast and ovarian cancer for women to development of a diverse intestinal microbiome for infants and lower infection risk in the first year of life.¹⁵⁻¹⁷ As any NICU

parent can attest, the impact of postpartum separation on parents' mental health and parent-child bonding can be enormous; this is part of the reason why NICUs feel kangaroo care is so important even for tiny infants who are otherwise kept in the strict quarantine environment of an isolette.¹⁸

There are other strategies beyond separation that would decrease COVID-associated risks for the newborns in our communities. The worst perinatal outcomes in this pandemic have been associated not with newborn infection but with maternal morbidity and mortality and premature birth due to maternal COVID-19. In a large cohort from China, researchers reported a preterm birth rate of 21% among women positive for COVID-19, 3 times higher than the national average.^{19,20} Current circumstances reaffirm an age-old truth: to better protect infants, we must better support the women who bear and raise them.

We could start by facilitating physical distancing for pregnant women without fear of job repercussions. We could bolster safety-net programs for families because we know that poverty, housing instability, food insecurity, and other socioeconomic factors drive COVID-19 infection risk just as powerfully as they impact other public health concerns. We could, like other developed countries, recognize that paid parental leave helps keep infants out of group child care settings in the first months of life when they are most vulnerable to respiratory syncytial virus and other infections every year, not just in 2020.

These interventions are the right thing to do for parents and children alike, but they require far more collective and political will than does a simple recommendation to separate infants and mothers. Asking a mother positive for COVID-19 to sacrifice the first days of infant bonding during

her hospital stay and then find a round-the-clock "clean infant caregiver" for a period of home quarantine is logistically challenging and a setup for postpartum depression even in the most privileged of circumstances. More often, we ask women to go home to crowded living conditions, take short unpaid maternity leaves, return to essential shift jobs that provide inadequate personal protective equipment, and place their young infants in the care of others; simultaneously, we ask them to protect those infants from infectious risk. This is an incongruous "plan of care." Physicians should advocate at higher levels for the solutions we wish we could provide.

What can we do for our patients right now? At an institutional level, we can craft policies that leave room for patient choice in the setting of incomplete data, and we can arrange physical spaces to safely care for mother-infant dyads positive for COVID-19 as well as separated mothers and infants. At the bedside, we can recognize that tangible infectious risks naturally elicit more fear than the more nebulous and longer-term harms of disrupting the postpartum experience, but when we have no clear data about the magnitude of COVID-19 risk for newborns, all factors merit consideration. And we can remember that words spoken during life's pivotal moments may have a lasting impact. Validating the importance of maternal instincts, even on a backdrop of fear or anxiety, sends an important and empowering message to women who are beginning their parenting journey amid challenging circumstances.

For me, today, the benefits of bonding outweigh concerns about infection. If I tested positive for COVID-19, I would wear a mask, wash my hands, and breastfeed my infant skin to skin. If I were ill, that might tip the scales toward isolation. But I am not sure.

This is the point: none of us can truly know what choices we would make until we are the one in the hospital bed. Wise counsel in the face of unclear risk requires not fixed recommendations but an ability to sit comfortably with uncertainty and mold it into something therapeutic for the patient before us. As a mother, uncertainty is part of the everyday job description; as a physician, acknowledging it can be the best way to help patients make informed decisions that are right for themselves and their families.

ACKNOWLEDGMENTS

I thank Jill Rinehart for inspiring inquiry into this subject; members of the University of Vermont Medical Center's family medicine, pediatrics, and internal medicine departments for thoughtful discussions; and Lewis First, John Boscia, Philip Murphy, and Nancy Murphy for helpful comments on the article.

ABBREVIATION

COVID-19: coronavirus disease 2019

REFERENCES

- Sutton D, Fuchs K, D'Alton M, Goffman D. Universal screening for SARS-CoV-2 in women admitted for delivery [published online ahead of print April 13, 2020]. *N Engl J Med*. doi:10.1056/NEJMc2009316
- Puopolo KM, Hudak M, Kimberlin DW, Cummings J. *Initial Guidance: Management of Infants Born to Mothers With COVID-19*. Itasca, IL: American Academy of Pediatrics Committee on Fetus and Newborn, Section on Neonatal and Perinatal Medicine, and Committee on Infectious Diseases; 2020
- World Health Organization. Clinical management of severe acute respiratory infection (SARI) when COVID-19 disease is suspected: interim guidance, 13 March 2020. Available at: <https://apps.who.int/iris/handle/10665/331446>. Accessed May 3, 2020
- Ma X, Zhu J, Du L. Neonatal management during coronavirus disease (COVID-19) outbreak: Chinese experiences. *NeoReviews*. 2020;21(5):e293–e297
- Savino M, Lank O. Questions linger whether a newborn baby from CT died from COVID-19. *WFSB News*. April 6, 2020. Available at: www.wfsb.com/news/questions-linger-whether-a-newborn-baby-from-ct-died-from-COVID-19/article_edaba41a-783e-11ea-b8da-139cba5c5d60.html. Accessed May 3, 2020
- Hinkel D. Chicago's top doctor says investigation has raised questions about whether infant died of COVID-19. *Chicago Tribune*. April 19, 2020. Available at: www.chicagotribune.com/coronavirus/ct-coronavirus-chicago-baby-death-20200418-vm6bsba45rfptpyrcmml4u3nh4-story.html. Accessed May 3, 2020
- Li AM, Ng PC. Severe acute respiratory syndrome (SARS) in neonates and children. *Arch Dis Child Fetal Neonatal Ed*. 2005;90(6):F461–F465
- Zeng H, Xu C, Fan J, et al. Antibodies in infants born to mothers with COVID-19 pneumonia. *JAMA*. 2020;323(18):1848–1849
- Melendi GA, Coviello S, Bhat N, Zea-Hernandez J, Ferolla FM, Polack FP. Breastfeeding is associated with the production of type I interferon in infants infected with influenza virus. *Acta Paediatr*. 2010;99(10):1517–1521
- Chen H, Guo J, Wang C, et al. Clinical characteristics and intrauterine vertical transmission potential of COVID-19 infection in nine pregnant women: a retrospective review of medical records. [published correction appears in *Lancet*. 2020;395(10229):1038]. *Lancet*. 2020;395(10226):809–815
- Jose RJ, Manuel A. COVID-19 cytokine storm: the interplay between inflammation and coagulation. *Lancet Respir Med*. 2020;8(6):E46–E47
- Moore ER, Bergman N, Anderson GC, Medley N. Early skin-to-skin contact for mothers and their healthy newborn infants. *Cochrane Database Syst Rev*. 2016;11(11):CD003519
- Saxton A, Fahy K, Rolfe M, Skinner V, Hastie C. Does skin-to-skin contact and breast feeding at birth affect the rate of primary postpartum haemorrhage: results of a cohort study. *Midwifery*. 2015;31(11):1110–1117
- Mehler K, Hucklenbruch-Rother E, Trautmann-Villalba P, Becker I, Roth B, Kribs A. Delivery room skin-to-skin contact for preterm infants-A randomized clinical trial. *Acta Paediatr*. 2020;109(3):518–526
- Chowdhury R, Sinha B, Sankar MJ, et al. Breastfeeding and maternal health outcomes: a systematic review and meta-analysis. *Acta Paediatr*. 2015;104(467):96–113
- Duijts L, Jaddoe VW, Hofman A, Moll HA. Prolonged and exclusive breastfeeding reduces the risk of infectious diseases in infancy. *Pediatrics*. 2010;126(1). Available at: www.pediatrics.org/cgi/content/full/126/1/e18
- Le Doare K, Holder B, Bassett A, Pannaraj PS. Mother's milk: a purposeful contribution to the development of the infant microbiota and immunity. *Front Immunol*. 2018;9:361
- Cho ES, Kim SJ, Kwon MS, et al. The effects of kangaroo care in the neonatal intensive care unit on the physiological functions of preterm infants, maternal-infant attachment, and maternal stress. *J Pediatr Nurs*. 2016;31(4):430–438
- Chen L, Li Q, Zheng D, et al. Clinical characteristics of pregnant women with COVID-19 in Wuhan, China [published online ahead of print April 17, 2020]. *N Engl J Med*. doi:10.1056/NEJMc2009226
- Chen C, Zhang JW, Xia HW, et al. Preterm birth in China between 2015 and 2016. *Am J Public Health*. 2019;109(11):1597–1604

Skin-to-Skin Care and COVID-19

Carolyn Boscia

Pediatrics 2020;146;

DOI: 10.1542/peds.2020-1836 originally published online May 19, 2020;

Updated Information & Services	including high resolution figures, can be found at: http://pediatrics.aappublications.org/content/146/2/e20201836
References	This article cites 13 articles, 1 of which you can access for free at: http://pediatrics.aappublications.org/content/146/2/e20201836#BIBL
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Hospital Medicine http://www.aappublications.org/cgi/collection/hospital_medicine_sub Patient Education/Patient Safety/Public Education http://www.aappublications.org/cgi/collection/patient_education:patient_safety:public_education_sub Infectious Disease http://www.aappublications.org/cgi/collection/infectious_diseases_sub Epidemiology http://www.aappublications.org/cgi/collection/epidemiology_sub
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.aappublications.org/site/misc/Permissions.xhtml
Reprints	Information about ordering reprints can be found online: http://www.aappublications.org/site/misc/reprints.xhtml

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®



PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Skin-to-Skin Care and COVID-19

Carolyn Boscia

Pediatrics 2020;146;

DOI: 10.1542/peds.2020-1836 originally published online May 19, 2020;

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/146/2/e20201836>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 345 Park Avenue, Itasca, Illinois, 60143. Copyright © 2020 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®

