

Equity in Emergency Mental Health Care

Polina Krass, MD,^{ab} Stephanie K. Douppnik, MD, MSHP^{ac}

Over the past 2 decades, young people have increasingly sought acute mental health care in emergency departments (EDs).^{1,2} In this issue of *Pediatrics*, Nash et al³ examine ED length of stay for children with mental health conditions. They draw on 10 years of data from the National Hospital Ambulatory Medical Care Survey (2005–2015) to compare length of stay for visits with and without a mental health diagnosis. In their sample, 4.9% of pediatric ED visits had an associated mental health diagnosis, and mental health visits had 3.3-fold increased odds of a length of stay of ≥ 6 hours compared with non-mental health visits. Length of stay increased over the 10-year study period to nearly a quarter of visits lasting ≥ 6 hours. This increase in length of stay over time was driven primarily by visits with a mental health diagnosis only (ie, no accompanying physical health diagnoses). The study's findings point to increasing friction in the process of accessing emergency mental health care for young people and highlight the importance of improving access to mental health services in an equitable way.

Hispanic patients were nearly 3 times more likely to experience an ED stay > 12 hours, compared with non-Hispanic white patients. Although the secondary data analysis in Nash et al³ was not positioned to explain this difference, racism, preferred language, and illness severity might play important roles. Hispanic youth are known to receive less mental health care and lower-quality mental health care than their non-Hispanic white

peers.⁴ Hispanic youth are also underrepresented in clinical trials of mental health treatments, and there are disproportionately few mental health professionals of Hispanic ethnicity,⁵ which may further reduce the efficacy of available mental health treatments for this population. Nash et al³ suggest that individual institutions examine local data to ensure that children of all races and ethnicities, particularly Hispanic children, have equitable access to mental health services.

Institutions may benefit from several interventions to improve the quality of mental health services. As a first step, they may audit how race, ethnicity, and preferred language are captured to ensure that the medical record accurately reflects a patient's self-identity and sheds light on the disparities that prevent families from receiving language-concordant, high-quality mental health care. To further improve quality of care, institutions might seek feedback from Hispanic and Black families and develop interventions that address inequities in care, such as reducing delays to accessing interpreter services and/or mental health evaluations. Some barriers may be difficult to resolve in the ED setting alone and will require a systems-level approach. For example, ED leaders might participate in regional partnerships to work toward ensuring that expedient and language-concordant services are available not just in the ED but also in inpatient psychiatric units and community mental health services.



^aPolicyLab and Department of Pediatrics, Children's Hospital of Philadelphia, Philadelphia, Pennsylvania; and ^bNational Clinician Scholars Program and ^cDepartment of Pediatrics, Perelman School of Medicine, University of Pennsylvania, Philadelphia, Pennsylvania

Drs Krass and Douppnik conceptualized and designed the study, drafted the initial manuscript, and reviewed and revised the manuscript; and all authors approved the final manuscript as submitted and agree to be accountable for all aspects of the work.

DOI: <https://doi.org/10.1542/peds.2020-049843>

Accepted for publication Feb 12, 2021

Address correspondence to Polina Krass, MD, Department of Pediatrics, Children's Hospital of Philadelphia, 3401 Civic Center Blvd, Philadelphia, PA 19104. E-mail: krassp@chop.edu

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2021 by the American Academy of Pediatrics

FINANCIAL DISCLOSURE: The authors have indicated they have no financial relationships relevant to this article to disclose.

FUNDING: Dr Douppnik's time was supported by National Institute of Mental Health award K23MH115162. Dr Krass received no external funding. Funded by the National Institutes of Health (NIH).

POTENTIAL CONFLICT OF INTEREST: The authors have indicated they have no potential conflicts of interest to disclose.

COMPANION PAPER: A companion to this article can be found online at www.pediatrics.org/cgi/doi/10.1542/peds.2020-030692.

To cite: Krass P and Douppnik SK. Equity in Emergency Mental Health Care. *Pediatrics*. 2021; 147(5):e2020049843

The coronavirus disease 2019 pandemic has introduced new challenges to pediatric mental health care. Increases in frequency and severity of mental illness⁶ are compounded by disruptions in nonessential health care access and increased risks associated with being in a crowded ED or an inpatient psychiatric unit. Black and Hispanic families have borne an undue burden of grief,⁷ financial instability,⁸ and educational disruption⁹ because of the pandemic and may suffer worse mental health outcomes as well.¹⁰

Policies to mitigate delays in accessing high-quality mental health services can help to alleviate these negative effects. The American Rescue Plan Act, a federal stimulus package approved by Congress on March 10, 2021,¹¹ offers federal funding to states to develop and sustain robust state crisis response systems. Such funding may help broaden the network of mental health crisis services beyond EDs and serve as a streamlined access point for care, regardless of ability to pay. However, even if additional crisis services become available, some youth will require stabilization in an ED. For these patients, the highest-quality ED mental health services would expand beyond triage, assessment, and safety observation. Billing and reimbursement strategies that incentivize the provision of mental health care by ED providers, including safety planning,¹² lethal-means reduction,¹³ and postdischarge mental health outreach,¹⁴ would enable more children to receive evidence-based mental health treatments more quickly.

As health care leaders plan for the future of mental health care in the wake of the coronavirus disease 2019 pandemic, the findings in Nash et al³

can help guide their decisions. Strengthening pediatric community mental health resources, including crisis response services, would facilitate access to high-quality mental health care outside of the ED. In addition, engaging members of marginalized communities in health care services planning will help ensure equitable access to mental health care for all children.

ABBREVIATION

ED: emergency department

REFERENCES

- Cutler GJ, Rodean J, Zima BT, et al. Trends in pediatric emergency department visits for mental health conditions and disposition by presence of a psychiatric unit. *Acad Pediatr*. 2019; 19(8):948–955
- Santillanes G, Axeen S, Lam CN, Menchine M. National trends in mental health-related emergency department visits by children and adults, 2009–2015. *Am J Emerg Med*. 2020; 38(12):2536–2544
- Nash KA, Zima BT, Rothenberg C, et al. Prolonged emergency department length of stay for US pediatric mental health visits (2005–2015). *Pediatrics*. 2021;147(5):e2020030692
- Alegria M, Vallas M, Pumariega AJ. Racial and ethnic disparities in pediatric mental health. *Child Adolesc Psychiatr Clin N Am*. 2010;19(4):759–774
- Santiago CD, Miranda J. Progress in improving mental health services for racial-ethnic minority groups: a ten-year perspective. *Psychiatr Serv*. 2014; 65(2):180–185
- Guessoum SB, Lachal J, Radjack R, et al. Adolescent psychiatric disorders during the COVID-19 pandemic and lockdown. *Psychiatry Res*. 2020;291: 113264
- Centers for Disease Control and Prevention. COVID-19 racial and ethnic health disparities. 2020. Available at: <https://www.cdc.gov/coronavirus/2019-ncov/community/health-equity/racial-ethnic-disparities>. Accessed February 2, 2021
- Couch KA, Fairlie RW, Xu H. Early evidence of the impacts of COVID-19 on minority unemployment. *J Public Econ*. 2020;192:104287
- Saavedra A, Rapaport A, Polikoff M, Silver D, Haderlein S, Garland M. Evidence of COVID-19's impact on K-12 education points to critical areas of intervention. The evidence blog. 2020. Available at: <https://healthpolicy.usc.edu/evidence-base/evidence-of-covid-19s-impact-on-k-12-education-points-to-critical-areas-of-intervention/>. Accessed February 2, 2021
- Purtle J. COVID-19 and mental health equity in the United States. *Soc Psychiatry Psychiatr Epidemiol*. 2020; 55(8):969–971
- American Rescue Plan Act of 2021, H.R.1319, 117th Cong. (2021). Available at: <https://www.congress.gov/bill/117th-congress/house-bill/1319/text>. Accessed March 11, 2021
- Stanley B, Brown GK, Brenner LA, et al. Comparison of the safety planning intervention with follow-up vs usual care of suicidal patients treated in the emergency department. *JAMA Psychiatry*. 2018;75(9):894–900
- Mueller KL, Naganathan S, Griffey RT. Counseling on Access to Lethal Means-Emergency Department (CALM-ED): a quality improvement program for firearm injury prevention. *West J Emerg Med*. 2020;21(5):1123–1130
- Milner AJ, Carter G, Pirkis J, Robinson J, Spittal MJ. Letters, green cards, telephone calls and postcards: systematic and meta-analytic review of brief contact interventions for reducing self-harm, suicide attempts and suicide. *Br J Psychiatry*. 2015; 206(3):184–190

Equity in Emergency Mental Health Care

Polina Krass and Stephanie K. Doupnik

Pediatrics 2021;147;

DOI: 10.1542/peds.2020-049843 originally published online April 5, 2021;

Updated Information & Services

including high resolution figures, can be found at:
<http://pediatrics.aappublications.org/content/147/5/e2020049843>

References

This article cites 11 articles, 2 of which you can access for free at:
<http://pediatrics.aappublications.org/content/147/5/e2020049843#BL>

Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):
Emergency Medicine
http://www.aappublications.org/cgi/collection/emergency_medicine_sub
Psychiatry/Psychology
http://www.aappublications.org/cgi/collection/psychiatry_psychology_sub

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
<http://www.aappublications.org/site/misc/Permissions.xhtml>

Reprints

Information about ordering reprints can be found online:
<http://www.aappublications.org/site/misc/reprints.xhtml>

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®



PEDIATRICS[®]

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Equity in Emergency Mental Health Care

Polina Krass and Stephanie K. Doupnik

Pediatrics 2021;147;

DOI: 10.1542/peds.2020-049843 originally published online April 5, 2021;

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/147/5/e2020049843>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 345 Park Avenue, Itasca, Illinois, 60143. Copyright © 2021 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN[®]

