

# Ambient Air Pollution: Respiratory Hazards to Children

Committee on Environmental Health

Levels of many outdoor air pollutants decreased substantially after the passage of the Clean Air Act of 1970; however, levels of ozone, carbon monoxide, and particulate matter are still high enough to present hazards to children. Failure to meet the federal standards for these pollutants was a major force driving the adoption of the revised Clean Air Act of 1990. In addition, recent research indicates that acidic aerosols, for which there are no health-based standards, may be associated with adverse respiratory effects.

As an ambient air pollutant, ozone is formed by the action of sunlight on nitrogen oxides and reactive hydrocarbons (both of which are emitted by motor vehicles and industrial sources). Ozone levels therefore tend to be highest on warm, sunny days, which are conducive to outdoor activities. In many areas ozone concentrations peak in the midafternoon, when children are likely to be playing outside. It is important to distinguish ground-level ozone air pollution from stratospheric ozone depletion by chlorofluorocarbons. These issues are unrelated. Carbon monoxide, a product of incomplete combustion, is emitted mainly from cars and other mobile sources. Airborne particulate matter is a variable and complex mixture of natural materials and substances released from numerous industries, motor vehicles, residential wood burning, construction and demolition, and other sources. Acidic aerosols are traceable mainly to combustion of sulfur-containing fossil fuels and to reactions of photochemical free radicals with nitrogen dioxide.

Exposure to ambient air pollution in North America has been clearly associated with acute and subacute effects in epidemiologic investigations and in controlled exposure studies in environmental chambers. For example, ozone causes airway inflammation and hyperreactivity, bronchial epithelial permeability, decrements in pulmonary function, cough, chest tightness, pain on inspiration, and upper respiratory tract irritation.<sup>1-7</sup> Nonrespiratory effects associated with ozone exposure include nausea, headache, malaise, and decreased ability to perform sustained exercise.<sup>7-9</sup> Epidemiologic studies link increased ozone concentrations with exacerbations of asthmatic symptoms.<sup>10,11</sup> Controlled chamber studies suggest that low concentrations of ozone do not cause dramatic bronchoconstriction in asthmatic vol-

unteers, although at higher concentrations asthmatics experience greater airway obstruction than healthy study subjects.<sup>12-14</sup>

Although healthy children appear to experience losses in pulmonary function comparable with those observed in adults for a given dose of ozone, children do not report symptoms to the same extent.<sup>15-18</sup> This suggests that children may not experience or recognize somatic signals to curtail exposure. Field studies suggest that ozone effects on pulmonary function in children are much greater than would be predicted from chamber studies.<sup>19</sup> Moreover, decreased peak flow in children has been reported to persist for up to a week following exposure to ozone concentrations lower than 0.2 ppm, suggesting the presence of damage to the respiratory tract.<sup>20</sup> Repeated exposures may result in persistent bronchial hyperresponsiveness.<sup>21</sup>


Controlled exposures to low ozone concentrations (at and below the current federal standard of 0.12 ppm) involving moderate levels of exercise for several hours have resulted not only in pulmonary function changes and respiratory symptoms, but also in dramatic increases in inflammatory markers in bronchoalveolar lavage fluid.<sup>1,2,5</sup> These findings are consistent with animal studies indicating that repeated exposures to ozone concentrations found in typical urban air result in centriacinar inflammation and small-airway structural changes.<sup>22-24</sup> Epidemiologic studies suggest that repeated exposures to ozone and other photochemical oxidants and particulates are associated with an accelerated decline in lung function and with symptoms of chronic respiratory disease; however, the quantitative aspects of such a relationship have not been adequately explored.<sup>25,26</sup> One recent study links oxidants (primarily ozone) and other air pollutants in Los Angeles, CA, with daily mortality.<sup>27</sup>

Epidemiologic studies undertaken in a variety of locations indicate a relationship between outdoor air pollution and adverse respiratory effects in children. The pollutants most frequently implicated in these studies have been respirable particles (notably acidic sulfates) and ozone. Examples of health outcomes found to be correlated with air pollution levels include increased prevalence of chronic cough, chest illness and bronchitis (measured by questionnaire), hospital admissions for various respiratory conditions, and decrements in lung function.<sup>28-33</sup> The prevalence of respiratory symptoms was markedly increased among children with a history of asthma or wheezing.<sup>34</sup>

The recommendations in this statement do not indicate an exclusive course of treatment or serve as a standard of medical care. Variations, taking into account individual circumstances, may be appropriate.

Received for publication Mar 18, 1993; accepted Mar 19, 1993.

PEDIATRICS (ISSN 0031 4005). Copyright © 1993 by the American Academy of Pediatrics.



**Figure.** Relationship between ambient ozone levels and respiratory tract effects in humans and experimental animals. EPA, Environmental Protection Agency.

Controlled studies involving adolescents with asthma have found that exposure to acidic aerosols affects results of pulmonary function tests.<sup>35,36</sup> A recent study of asthmatic adults found a significant association between ambient airborne acidity and daily fluctuations of asthmatic symptoms, including cough and shortness of breath.<sup>37</sup> Acidic aerosols also have been found to provoke changes in tracheobron-

chial clearance and increased airway reactivity in normal subjects.<sup>38,39</sup>

The effects of exposures to multiple pollutants are difficult to study in humans. A few controlled investigations and field studies indicate, however, that exposures to complex mixtures of air pollutants may have synergistic acute effects on pulmonary function and, possibly, on symptoms.<sup>19,40,41</sup> A recent report

suggests that even brief exposure to ozone can potentiate allergic asthmatic responses to aeroallergens.<sup>42</sup> There is, moreover, a substantial body of experimental evidence in animals indicating that ozone can lower resistance to infection, facilitate sensitization and airway responses to airborne allergens, and act synergistically with airborne acidity to damage deep lung tissues.<sup>43-47</sup>

### CONCLUSIONS

Existing epidemiologic and toxicologic data indicate that exposure to ambient air pollution is associated with respiratory toxicity. The decrements in pulmonary function observed in epidemiologic and experimental studies involving children exposed to ozone and other pollutants may last longer than the episodes of pollution that initiate these changes.

A factor that increases children's vulnerability to airborne pollution is that their airways are narrower than those of adults. Thus, irritation caused by air pollution that would produce only a slight response in an adult can result in potentially significant obstruction in the airways of a young child. Moreover, children have markedly increased needs for oxygen relative to their size. They breathe more rapidly and inhale more pollutant per pound of body weight than do adults. In addition, they often spend more time engaged in vigorous outdoor activity than adults. Experimental and epidemiologic data provide grounds for concern about chronic lung damage from repeated exposures.

Current strategies in the United States for attaining clean air and protecting public health have been only partially successful. Thus, the American Academy of Pediatrics offers the following recommendations:

#### Recommendations to Government Agencies

1. *Ambient standards.* The federal ambient air standard for ozone of 0.12 ppm (averaged over 1 hour) contains little or no margin of safety for children engaged in active outdoor activity. In view of recent research indicating the occurrence of adverse effects at ozone concentrations lower than the current standard, the Academy recommends that the standard be reconsidered for possible lowering (see Figure). Similarly, epidemiologic evidence has shown that the current federal standard for particulate matter provides less than optimal protection of public health and should be lowered.

2. *Smog alerts.* State and local government agencies have a responsibility to issue pollution or smog alerts in a clear and timely manner. These alerts should warn specifically of the hazards that air pollution presents to children. Furthermore, recent evidence indicates that respiratory toxicity occurs at ozone concentrations lower than the stage 1 smog alert concentration (0.20 ppm, 1 hour average) recommended by the US Environmental Protection Agency, suggesting the need for reconsideration of this advisory level. Among other things, the stage 1 alert level triggers advisories to schools that outdoor activities should be restricted.

3. *Source control.* State and federal governments must act more vigorously in the arena of pollution

prevention, in terms of both technologic requirements and public education. Regulatory agencies should act aggressively to implement the requirements of the Clean Air Act of 1990.

#### Recommendations to Pediatricians

1. Pediatricians should become informed about air pollution problems in the community.
2. Pediatricians caring for children at special risk, such as those with asthma and cystic fibrosis, should be aware that current levels of air pollution may cause deterioration in these children's pulmonary function and may aggravate their symptoms.
3. Pediatricians who serve as physicians for schools and for students participating in team sports need to be aware of the health implications of pollution alerts in order to provide appropriate guidance to schools and other public agencies on the health hazards of air pollution.
4. Pediatricians can make parents aware of the predictable daily variation in ozone, especially the tendency to peak in the afternoon. This awareness is essential in areas with recognized high ozone levels. When ozone levels are elevated, it may be possible to decrease children's exposure by scheduling outdoor sports earlier in the day.
5. Pediatricians can help children by expressing their concern about the child health hazards of air pollution to their representatives and to policymakers within state and federal governmental agencies.

#### COMMITTEE ON ENVIRONMENTAL HEALTH, 1992 TO 1993

J. Routt Reigart, MD, Chairperson

Ruth A. Etzel, MD

Lynn R. Goldman, MD, MPH

Jim G. Hendrick, MD

Howard C. Mofenson, MD

Peter R. Simon, MD

#### Liaison Representatives

Henry Falk, MD, Centers for Disease Control

Robert W. Miller, MD, National Cancer Institute

Walter Rogan, MD, National Institute of Environmental Health Sciences

#### Consultants

Herbert L. Needleman, MD

Richard J. Jackson, MD

Philip Landrigan, MD

Michael Lipsett, MD

#### REFERENCES

1. Folinsbee LJ, McDonnell WF, Horstman DH. Pulmonary function and symptom responses after 6.6-hour exposure to 0.12 ppm ozone with moderate exercise. *J Air Pollut Control Assoc.* 1988;38:28-35
2. Horstman D, Folinsbee L, Ives PJ, Abdul-Salaam S, McDonnell WF. Ozone concentration and pulmonary response relationships for 6.6-hour exposures with five hours of moderate exercise to 0.08, 0.10 and 0.12 ppm. *Am Rev Respir Dis.* 1990;142:1158-1163
3. Kinney PL, Ware JH, Spengler JD, Dockery DW, Speizer FE, Ferris BG Jr. Short-term pulmonary function change in association with ozone levels. *Am Rev Respir Dis.* 1989;139:56-61
4. Kehrl HR, Vincent LM, Kowalsky RJ, et al. Ozone exposure increases respiratory epithelial permeability in humans. *Am Rev Respir Dis.* 1987;135:1124-1128
5. Koren HS, Devlin RB, Graham DE, et al. Ozone-induced inflammation in the lower airways of human subjects. *Am Rev Respir Dis.* 1989;139:407-415
6. Seltzer J, Bigby BG, Stulbarg M, et al. O<sub>3</sub>-induced change in bronchial

- reactivity to methacholine and airway inflammation in humans. *J Appl Physiol*. 1986;60:1321-1326
7. Lippmann M. Health effects of ozone: a critical review. *J Air Pollut Control Assoc*. 1983;54:1345-1352
  8. McDonnell WF, Horstman DH, Hazucha MJ, et al. Pulmonary effects of ozone exposure during exercise: dose-response characteristics. *J Appl Physiol*. 1983;54:1345-1352
  9. Adams WC. Effects of ozone exposure at ambient air pollution episode levels on exercise performance. *Sports Med*. 1987;4:395-424
  10. Whittmore AS, Korn EL. Asthma and air pollution in the Los Angeles area. *Am J Public Health*. 1980;70:687-696
  11. Schoettlin CE, Landau E. Air pollution and asthmatic attacks in the Los Angeles area. *Public Health Rep*. 1961;76:545-548
  12. Linn WS, Buckley RD, Spier CE, et al. Health effects of ozone exposure in asthmatics. *Am Rev Respir Dis*. 1978;117:835-843
  13. Koenig JQ, Covert DS, Morgan MS, et al. Acute effects of 0.12 ppm ozone or 0.12 ppm nitrogen dioxide on pulmonary function in healthy and asthmatic adolescents. *Am Rev Respir Dis*. 1985;132:648-651
  14. Kreit JW, Gross KB, Moore TB, Lorenzen TJ, D'Arcy J, Eschenbacher WL. Ozone-induced changes in pulmonary function and bronchial responsiveness in asthmatics. *J Appl Physiol*. 1989;66:217-222
  15. Avol EL, Linn WS, Shamoo DA, et al. Respiratory effects of photochemical oxidant air pollution in exercising adolescents. *Am Rev Respir Dis*. 1985;132:619-622
  16. Avol EL, Linn WS, Shamoo DA, et al. Short-term respiratory effects of photochemical oxidant exposure in exercising children. *J Air Pollut Control Assoc*. 1987;37:158-162
  17. McDonnell WF, Chapman RS, Leigh MW, Strobe GL, Collier AM. Respiratory responses of vigorously exercising children to 0.12 ppm ozone exposure. *Am Rev Respir Dis*. 1985;132:875-879
  18. Spektor DM, Lippmann M, Liou PJ, et al. Effects of ambient ozone on respiratory function in active, normal children. *Am Rev Respir Dis*. 1988;137:313-320
  19. Spektor DM, Thurston GD, Mao J, He D, Hayes C, Lippmann M. Effects of single- and multiday ozone exposures on respiratory function in active normal children. *Environ Res*. 1991;55:107-122
  20. Liou PJ, Vollmuth TA, Lippmann M. Persistence of peak flow decrement in children following ozone exposures exceeding the National Ambient Air Quality Standard. *J Air Pollut Control Assoc*. 1985;35:1069-1071
  21. Zwick H, Popp W, Wagner C, et al. Effects of ozone on the respiratory health, allergic sensitization, and cellular immune system in children. *Am Rev Respir Dis*. 1991;144:1075-1079
  22. Evans MJ. Oxidant gases. *Environ Health Perspect*. 1984;55:85-95
  23. Fujinaka LE, Hyde DM, Plopper CG, Tyler WS, Dungworth DL, Lollini LO. Respiratory bronchiolitis following long-term ozone exposure in bonnet monkeys: a morphometric study. *Exp Lung Res*. 1985;8:167-190
  24. Crapo JD, Barry BE, Chang LY, Mercer RR. Alterations in lung structure caused by inhalation of oxidants. *J Toxicol Environ Health*. 1984;13:301-321
  25. Detels R, Tashkin DP, Sayre JW, et al. The UCLA population studies of chronic obstructive respiratory disease, 9: lung function changes associated with chronic exposure to photochemical oxidants; a cohort study among never-smokers. *Chest*. 1987;92:594-603
  26. Euler GL, Abbey DE, Hodgkin JE, Magie AR. Chronic obstructive pulmonary disease symptom effects of long-term cumulative exposure to ambient levels of total oxidants and nitrogen dioxide in California Seventh-Day Adventist residents. *Arch Environ Health*. 1988;43:279-285
  27. Kinney PL, Ozkaynak H. Associations of daily mortality and air pollution in Los Angeles County. *Environ Res*. 1991;54:99-120
  28. Ware JH, Ferris BG Jr, Dockery DW, Spengler JD, Stram DO, Speizer FE. Effects of ambient sulfur oxides and suspended particles on respiratory health of preadolescent children. *Am Rev Respir Dis*. 1986;133:834-842
  29. Dassen W, Brunekreef B, Hoek G, et al. Decline in children's pulmonary function during an air pollution episode. *J Air Pollut Control Assoc*. 1986;36:1223-1227
  30. Bates DV, Sizto R. Air pollution and hospital admissions in Southern Ontario: the acid summer haze effect. *Environ Res*. 1987;43:317-331
  31. Bates DV, Sizto R. The Ontario air pollution study: identification of the causative agent. *Environ Health Perspect*. 1989;79:69-72
  32. Pope CA III. Respiratory disease associated with community air pollution and a steel mill, Utah Valley. *Am J Public Health*. 1989;79:623-628
  33. Pope CA III, Dockery DW, Spengler JD, Raizenne ME. Respiratory health and PM<sub>10</sub> pollution: a daily time series analysis. *Am Rev Respir Dis*. 1991;144:668-674
  34. Dockery DW, Speizer FE, Stram DO, Ware JH, Spengler JD, Ferris BG Jr. Effects of inhalable particles on respiratory health of children. *Am Rev Respir Dis*. 1989;139:587-594
  35. Koenig JQ, Covert DS, Pierson WE. Effects of inhalation of acidic compounds on pulmonary function in allergic adolescent subjects. *Environ Health Perspect*. 1989;79:173-178
  36. Koenig JQ, Pierson WE, Horike M. The effects of inhaled sulfuric acid on pulmonary function in adolescent asthmatics. *Am Rev Respir Dis*. 1983;128:221-225
  37. Ostro BD, Lipsett MJ, Wiener MB, Selner JC. Asthmatic responses to airborne acid aerosols. *Am J Public Health*. 1991;81:694-702
  38. Spektor DM, Yen BM, Lippmann M. Effect of concentration and cumulative exposure of inhaled sulfuric acid on tracheobronchial particle clearance in healthy humans. *Environ Health Perspect*. 1989;79:167-172
  39. Utell MJ, Morrow PE, Hyde RW. Airway reactivity to sulfate and sulfuric acid aerosols in normal and asthmatic subjects. *J Air Pollut Control Assoc*. 1984;34:931-935
  40. Koenig JQ, Covert DS, Hanley QS, Van Belle G, Pierson WE. Prior exposure to ozone potentiates subsequent response to sulfur dioxide in adolescent asthmatic subjects. *Am Rev Respir Dis*. 1990;141:377-380
  41. Spektor DM, Lippmann M, Thurston GD, et al. Effects of ambient ozone on respiratory function in healthy adults exercising outdoors. *Am Rev Respir Dis*. 1988;138:821-828
  42. Molfino NA, Wright SC, Katz I, et al. Effect of low concentrations of ozone on inhaled allergen responses in asthmatic subjects. *Lancet*. 1991;338:199-203
  43. Gardner DE. Oxidant-induced enhanced sensitivity to infection in animal models and their extrapolations to man. *J Toxicol Environ Health*. 1984;13:423-439
  44. Miller FJ, Illing JW, Gardner DE. Effect of urban ozone levels on laboratory-induced respiratory infections. *Toxicol Lett*. 1978;2:163-169
  45. Yanai M, Ohru T, Aikawa T. Ozone increases susceptibility to antigen inhalation in allergic dogs. *J Appl Physiol*. 1990;68:2267-2273
  46. Osebold JW, Gershwin LJ, Zee YC. Studies on the enhancement of allergic lung sensitization by inhalation of ozone and sulfuric acid aerosol. *J Environ Pathol Toxicol*. 1980;3:221-234
  47. Warren DL, Last JA. Synergistic interaction of ozone and respirable aerosols on rat lungs, III: ozone and sulfuric acid aerosol. *Toxicol Appl Pharmacol*. 1987;88:203-216

**Ambient Air Pollution: Respiratory Hazards to Children**  
*Pediatrics* 1993;91;1210

**Updated Information & Services**

including high resolution figures, can be found at:  
<http://pediatrics.aappublications.org/content/91/6/1210>

**Permissions & Licensing**

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:  
<http://www.aappublications.org/site/misc/Permissions.xhtml>

**Reprints**

Information about ordering reprints can be found online:  
<http://www.aappublications.org/site/misc/reprints.xhtml>

**American Academy of Pediatrics**

DEDICATED TO THE HEALTH OF ALL CHILDREN®





# PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

**Ambient Air Pollution: Respiratory Hazards to Children**  
*Pediatrics* 1993;91;1210

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/91/6/1210>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 345 Park Avenue, Itasca, Illinois, 60143. Copyright © 1993 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®

