

# Adverse Events Associated With Exchange Transfusion in Healthy and Ill Newborns

J. Craig Jackson, MD

**ABSTRACT.** *Objective.* To determine the incidence of adverse events attributable to exchange transfusion during the past 15 years and compare the incidence of severe complications between healthy and ill infants.

*Design.* Medical records for the past 15 years from two teaching hospitals with neonatal intensive care units were reviewed. Those newborns who underwent exchange transfusions were classified as healthy or ill. Adverse events were analyzed to determine whether they were attributable to the procedure.

*Results.* Of the 106 patients who underwent exchange transfusion, 81 were healthy and had no medical problems other than jaundice. The remaining 25 patients were classified as ill and had medical problems ranging from mild to severe. At least 2 (2%) of the 106 patients died of complications probably attributable to exchange transfusion. None of the 81 healthy infants died, but 1 had severe necrotizing enterocolitis requiring surgery. Of the 25 ill infants, at least 3 (12%) experienced severe complications (including 2 deaths) probably attributable to exchange transfusion. Serious complications from the most common adverse events, hypocalcemia and thrombocytopenia, were limited to the group of infants already ill with other medical problems.

*Conclusions.* Because of the significantly greater rate of severe complications in ill infants, exchange transfusion should be delayed until the risk of bilirubin encephalopathy is as high as the risks of severe complications from the procedure itself (12%). These results do not support recommendations to use lower exchange levels in ill infants compared with healthy infants. *Pediatrics* 1997;99(5). URL: <http://www.pediatrics.org/cgi/content/full/99/5/e7>; *exchange transfusion, whole-blood; adverse events, jaundice, neonatal; kernicterus, infant, newborn.*

---

ABBREVIATION. NICHD, National Institute of Child Health and Human Development.

---

The bilirubin level at which exchange transfusion is indicated remains controversial.<sup>1-3</sup> This is because it is very difficult to define the risk of bilirubin encephalopathy in various categories of patients, such as those with or without hemolysis, healthy or ill, term or premature. The recommendations attempt to balance the benefits of preventing bilirubin toxicity with the risks of exchange transfusion. However,

there are few recent reports of the complication rates from exchange transfusion or attempts to stratify the risk of adverse events based on clinical condition.

Mortality rates attributable to exchange transfusion ranged from .65% to 3.2% in studies performed in the 1960s<sup>4-7</sup> and from .4% to 3.2% during the 1970s and 1980s.<sup>8-10</sup> Causes of death ascribed to exchange transfusion included cardiovascular collapse during the transfusion, and the subsequent complications of necrotizing enterocolitis, bacterial sepsis, and pulmonary hemorrhage.

The most frequently cited review of adverse events from exchange transfusion is from the 1974 to 1976 prospective National Institute of Child Health and Human Development (NICHD) phototherapy study. Keenan et al<sup>11</sup> reported that 190 infants underwent 331 exchange transfusions. Adverse clinical problems were observed in 6.7% of the exchange transfusions, and the observed rate of serious morbidity was 5.2%. Based on one death attributed to the procedure, they calculated mortality rate to be .53 per 100 patients and .3 per 100 procedures. Only 2 of the 14 serious adverse events occurred in infants defined as being in good condition at the initiation of the exchange transfusion.

Most of the published experience regarding adverse events from exchange transfusions comes from studies performed more than 15 years ago. There have been many advances in neonatal intensive care since that time that may have reduced the incidence of adverse events. For instance, the 1985 report by Hovi and Simmes<sup>8</sup> indicated that electronic monitors were not routinely used, and that only in the final year of their study—and only in some cases—did they use continuous monitoring of electrocardiogram, blood pressure, or transcutaneous oxygen tension.

Improvements in outcome from advances in neonatal care may have been offset by the increased risk of adverse events attributable to inexperience with the procedure. Due to tolerance of higher bilirubin levels in term infants without evidence of hemolysis<sup>1-3</sup> and improved obstetric management of Rh-sensitized mothers, the last 15 years have seen a much lower frequency of exchange transfusion. Furthermore, the incidence of kernicterus in preterm infants appears to be lower in recent years and exchange transfusion is frequently deferred in patients with serum bilirubin greater than the exchange levels used in the NICHD study.<sup>12</sup>

In previously published reports of the mortality and morbidity from exchange transfusion, many of

---

From the Department of Pediatrics, University of Washington, Children's Hospital and Medical Center, Seattle, Washington.

Received for publication Feb 23, 1996; accepted Jun 4, 1996.

Address correspondence to: J. Craig Jackson, MD, Children's Hospital and Medical Center, CH-35, 4800 Sand Point Way NE, Box 5371, Seattle, WA 98105-0371.

PEDIATRICS (ISSN 0031 4005). Copyright © 1997 by the American Academy of Pediatrics.

the patients were sick or premature or both. The rates of complications in healthy newborns undergoing the procedure were not usually reported separately, although most of the adverse events occurred in infants with other significant medical problems. As clinicians and parents weigh the risks of the procedure against its benefits in individual situations, it would be helpful to have estimates of the mortality and morbidity attributable to exchange transfusion stratified by clinical condition.

The present study was undertaken to determine the incidence of adverse events attributable to exchange transfusion performed during the past 15 years. All adverse events possibly caused by exchange transfusion were evaluated regardless of how late the event occurred after the procedure. The rate of complications from exchange transfusion in jaundiced but otherwise healthy infants was analyzed separately from those occurring in ill infants with other medical problems.

## METHODS

The computerized discharge abstract summaries of all patients admitted in the first month of life to Children's Hospital and Medical Center during 1981 through 1995 and the University of Washington Medical Center during 1980 through 1995 were searched for the procedure code for exchange transfusion. After eliminating records for patients who underwent only partial exchange transfusion for polycythemia, the medical records of the 106 remaining patients were reviewed in detail. Those newborns admitted solely for asymptomatic hyperbilirubinemia were classified as healthy. The remaining infants—those with any other medical conditions—were classified as ill. All patients were cared for in neonatal intensive care units by University of Washington pediatric residents and neonatal fellows under the close supervision of academic and clinical neonatologists.

The cause of jaundice reported in the record was classified in the following way: Rh disease was defined as jaundice in Rh positive newborns from Rh negative mothers with elevated titers to the Rh antigen and evidence of hemolysis. ABO disease was defined as jaundice in newborns with positive direct Coombs test against the A or B antigens from type O mothers; hemolysis was often but not always documented. Other antigen sensitization was defined as hemolytic jaundice in Coombs-positive newborns from mothers with antibodies to other blood group antigens.

All blood used for exchange transfusion was obtained from the Puget Sound Blood Center. Blood from volunteer donors was anticoagulated with citrate phosphate dextrose adenosine-1 and was <5 days old. Either whole blood ABO compatible with both the baby and mother, or group O red cells resuspended in compatible (usually AB) plasma, were used. Exchange transfusions were performed by the fellow or attending neonatologist, or by pediatric residents under their direct supervision. The double-volume exchange procedures were generally completed in about 2 hours by repeatedly removing and replacing small aliquots of blood (<5 mL/kg) according to standard published guidelines.

The records were reviewed for adverse events possibly attributable to exchange transfusion and classified into six prospectively defined categories of severity (Table 1). For simplicity of data presentation and because it was difficult to assign complications to a specific transfusion in those undergoing multiple procedures, rates of complications were calculated on the number of treated infants rather than the number of procedures performed.

Each infant was assigned to the one category best describing his or her most serious complications, to allow for the calculation of cumulative percentages of adverse events of decreasing severity. However, all adverse events were recorded to determine the overall frequency of the most common complications. The rate of severe complications in healthy infants was compared with that in ill infants with the Fisher's exact test. The 95% confidence intervals for estimates of risk were determined from standard tables.<sup>13</sup>

**TABLE 1.** Classification of Adverse Event Severity

Asymptomatic lab abnormalities
During or immediately after exchange transfusion, previously normal value for sodium, potassium, glucose, or calcium values fell outside the normal range, or platelets fell below 50 000/ $\mu$ L. No symptoms and no treatment given.
Asymptomatic complications, treated
Asymptomatic patients whose laboratory values were judged severe enough to require additional treatment (platelet transfusion, antibiotics) or who had complications of management (eg, clotting of catheter).
Serious, transient complications
Symptomatic patients whose adverse events were treated and which resolved quickly.
Serious, prolonged complications
Symptomatic patients with serious problems whose problems eventually resolved.
Permanent serious sequelae
Serious complications that resulted in permanent bodily alterations.
Death
Died due to adverse events, regardless of the interval after the transfusion.

## RESULTS

During the 15-year study period, there were approximately 15 000 neonatal intensive care unit admissions. One hundred six patients underwent exchange transfusion (Table 2). Of these, 81 were healthy and had no medical problems other than jaundice. The remaining 25 patients were classified as ill because they had additional medical problems; 10 required mechanical ventilation for hyaline membrane disease and the remainder had an assortment of conditions ranging from mild to severe (Table 3).

The 106 patients underwent 140 exchange transfusions. Ill infants underwent more multiple transfusions than healthy infants (Table 4). In 77% of the healthy infants, exchange transfusion was initiated via a catheter in the umbilical vein, whereas in 58% of the ill infants, it was initiated via a catheter in the umbilical artery.

The most common cause for jaundice was hemolysis from sensitization to Rh (n = 51), ABO (n = 21), or other blood group antigens (n = 6). Other causes of jaundice included unknown (n = 18), prematurity (n = 6), and one each of the following: hemolysis with polycythemia, hereditary spherocytosis, cholestasis from gastroschisis, and adrenal hemorrhage with cholestasis.

### Death

Two infants died of respiratory failure that did not appear to be attributable to exchange transfusion. The deaths of five other infants were possibly related to complications from exchange transfusion (Appendix A). All of the deaths were in the group of 25 ill

**TABLE 2.** Gestational Age and Body Weight (Mean  $\pm$  SD and Range)

	All Infants (n = 106)	Healthy Infants (n = 81)	Ill Infants (n = 25)
Gestational age, weeks	36.6 $\pm$ 3.6	37.5 $\pm$ 2.6	33.4 $\pm$ 4.7
Range, weeks	(24–41)	(31–41)	(24–40)
Body weight, g	2846 $\pm$ 806	2996 $\pm$ 588	2367 $\pm$ 1167
Range, g	(630–4520)	(1237–4436)	(630–4520)

**TABLE 3.** Primary Additional Medical Problems Before Exchange Transfusion in the 25 Ill Infants

12 required mechanical ventilation:
Hyaline membrane disease (n = 10)
Hydrops
Hyperkalemia and severe asphyxia
13 had no requirement for mechanical ventilation:
Hyaline membrane disease
Hydrops
Respiratory distress, desaturations, lethargy, irritability
Asphyxia and polycythemia
Lethargy, emesis, melena
Apnea, bradycardia, posturing
Polycythemia and fever
Dehydration
<i>Staphylococcus epidermidis</i> sepsis and adrenal hemorrhage
Large cephalohematoma
Antibiotics for maternal pretreatment
Stenosis of aortic and pulmonic valves
Gastroschisis

**TABLE 4.** Number of Exchange Transfusions

Number of Exchange Transfusions Per Patient	Healthy Infants		Ill Infants	
	Patients	Transfusions	Patients	Transfusions
1	65	65	16	16
2	15	30	4	8
3	1	3	3	9
4	0	0	1	4
5	0	0	1	5
Total	81	98	25	42

infants. The deaths of two infants (#3 and #4 in Appendix A) were classified as probably attributable to exchange transfusion, ie, more likely than not to have been a complication of the procedure (Table 5).

**Permanent Serious Sequelae**

Four infants suffered permanent serious sequelae possibly attributable to exchange transfusion (Appendix B). One was from the group of 81 healthy infants, and 3 were from the 20 remaining ill infants. The permanent serious complications of 2 infants (A and C in Appendix B) were classified as probably attributable to exchange transfusion (Table 5).

**Serious, Prolonged Complications**

In the group of 25 ill infants, 1 term infant with umbilical artery catheter developed renovascular hypertension requiring chronic antihypertensive medication. Although this complication might have occurred even if the exchange transfusions had not been performed through the catheter, it was attributed to exchange transfusion because blood products increase the risk of catheter thrombus and because

this infant received five exchange transfusions through the catheter.

In the group of 81 healthy newborns, 4 additional infants developed serious prolonged complications probably attributable to exchange transfusion. One developed *Staphylococcus aureus* bacteremia 2 days after exchange transfusion, and two others developed omphalitis in association with umbilical catheter placement for exchange transfusion; all required prolonged courses of antibiotics. One developed a serious purpuric eruption similar to porphyria, thought to be secondary to exchange transfusion. Adverse events in all 4 infants were considered probably attributable to exchange transfusion.

**Serious, Transient Complications**

In the group of 81 healthy newborns, 14 additional infants experienced serious transient complications probably attributable to exchange transfusion. These complications included apnea or bradycardia with cyanosis requiring resuscitation (usually with positive pressure ventilation by mask) during or immediately after exchange transfusion (n = 5), hypocalcemia associated with electrocardiographic abnormalities, marked jitteriness, or pedal spasm (n = 3), rectal bleeding leading to work-up and brief medical treatment for presumed necrotizing enterocolitis (n = 2), and one each of the following complications: surgery for removal of knotted femoral vein catheter guidewire, hypertension and hematuria associated with umbilical artery catheter, transient bacteremia, and petechial rash from thrombocytopenia.

In the group of 25 ill newborns, 4 additional infants suffered serious transient complications: rectal bleeding associated with thrombocytopenia from exchange transfusion, bleeding from umbilical stump requiring platelet transfusion, severe apnea and bradycardia with cyanosis requiring resuscitation with positive pressure ventilation, and jitteriness associated with hypocalcemia.

**Asymptomatic Treated Complications**

Of the 81 healthy infants, 21 had only asymptomatic complications, including clotting of umbilical catheter requiring replacement or discontinuation of the exchange transfusion (n = 10), intravenous calcium for hypocalcemia (n = 9), and one each of the following: both intravenous calcium for hypocalcemia and clotting of umbilical catheter, platelet transfusion attributable to severe thrombocytopenia, and discovery shortly after beginning exchange that the

**TABLE 5.** Complications Probably Due to Exchange Transfusion

	Previously Healthy (n = 81)			Previously Ill (n = 25)		
	n	%	Cumulative %	n	%	Cumulative %
Death	0	0%	0%	2	8%	8%
Permanent serious sequelae	1	1%	1%	1	4%	12%
Serious, prolonged complications	4	5%	6%	1	4%	16%
Serious, transient complications	14	17%	23%	4	16%	32%
Asymptomatic, treated	22	27%	51%	6	24%	56%
Asymptomatic lab abnormalities	10	12%	63%	1	4%	60%

wrong type of blood (packed cells) had been delivered for the exchange.

Of the 25 ill infants, 6 had only asymptomatic complications, including administration of calcium for hypocalcemia (n = 3), clotting of umbilical catheter requiring replacement (n = 2), and 1 who was treated for both hypocalcemia and thrombocytopenia.

#### Asymptomatic Laboratory Abnormalities

Of the 81 healthy infants, 10 had only asymptomatic and untreated laboratory abnormalities, including hypocalcemia (n = 6), thrombocytopenia (n = 2), hypocalcemia and thrombocytopenia (n = 1), and hyponatremia and thrombocytopenia (n = 1). Of the 25 ill infants, 1 had only hypocalcemia.

#### Implications for Laboratory Monitoring

The most common adverse events were related to hypocalcemia and thrombocytopenia (Table 6 and Table 7). Only two infants had hypoglycemia during or after exchange transfusion (mild and asymptomatic in both cases) and one had asymptomatic hyponatremia (121 mEq/dL).

#### Estimates of Risk for Severe Complications

For statistical comparisons between healthy and ill infants, death and permanent serious sequelae were combined and defined as severe complications. The 12% rate of severe complications in the ill infants (3 of 25) is significantly greater ( $P < .05$ ) than the 1.2% observed in the healthy infants (1 of 81). Based on the number of infants in this review, the 95% confidence interval for the 12% estimate of severe complications in ill infants is 3% to 31%. The 95% confidence interval for the 1.2% estimate in healthy infants is 0% to 7%.

### DISCUSSION

Despite improvements in neonatal intensive care in the past two decades, exchange transfusion remains a high-risk procedure. Two of the 106 patients in this study (2%) died of complications probably attributable to exchange transfusion, similar to previous reports.<sup>4-11</sup> In the present study, none of the 81 healthy infants undergoing exchange transfusion died. This observation, and the paucity of reports of transfusion-related deaths in healthy infants, suggests that the mortality rate for the procedure in this population is well below 1%. The rate of permanent serious sequelae from exchange transfusion is also very low, approximately 1%, and both this study and prior reports indicate that necrotizing enterocolitis is the most common severe complication. Thus, the

bilirubin exchange level for healthy infants should be set at a level at which the risk of bilirubin encephalopathy is no higher than 1%.

In ill infants, the incidence of procedure-related complications leading to death was 8% and the rate of severe complications (death or permanent serious sequelae) was 12%. Clinicians and parents should be aware of the much greater incidence of severe complications in this population when judging whether to perform an exchange transfusion. The bilirubin exchange level for ill infants should be at a point where the risk of bilirubin encephalopathy is approximately 12%. It is commonly assumed—although with limited evidence—that infants with risk factors such as prematurity, asphyxia, acidosis, and respiratory distress syndrome are at higher risk for bilirubin encephalopathy. Unless their risk is more than 10-fold greater than for healthy infants at any given level of bilirubin, the exchange levels for ill and healthy infants should be the same. Unfortunately, we probably have better data on the risks of exchange transfusion than on the risks of hyperbilirubinemia.

Unless the procedure can be made safer in this high-risk population, severe complications are inevitable. Overuse of the procedure may reduce the incidence of bilirubin encephalopathy at the cost of increased incidence of severe procedure-related complications. If exchange transfusions are delayed until the risks of severe complications from the procedure are equal to the risks of bilirubin encephalopathy, the number of patients with each will be roughly equal. A review of the medical records from our neurodevelopmental clinics indicated that only three children cared for in our nurseries during the past 15 years have been diagnosed as having bilirubin encephalopathy or kernicterus. Thus, during 15 years, four infants suffered severe complications from exchange transfusion in order to prevent a disease that developed in only three children.

Uncertainty regarding the figure of 12% incidence of severe complications in the ill infants is due first to the small size of the population, only 25 patients. The 95% confidence interval for three events out of a total of 25 is 3% to 31%. Second, there is uncertainty in ascribing complications in already ill infants to exchange transfusion. The estimate is too high if none of the severe complications outlined in Appendices A and B were attributable to exchange transfusion, and too low if all of them were.

The results of this study have implications for monitoring both healthy and ill infants who undergo exchange transfusion. The most common serious morbidities included symptomatic hypocalcemia,

**TABLE 6.** Complications Associated With Hypocalcemia

	81 Healthy Infants	25 Ill Infants
Cardiac arrest associated with hypocalcemia	0	1 (4%)
Serious, transient complications*	4 (5%)	1 (4%)
Asymptomatic hypocalcemia (treated)	15 (19%)	6 (24%)
Asymptomatic hypocalcemia (untreated)	13 (16%)	4 (16%)
Normal calcium or not measured	49 (60%)	13 (52%)

\* ECG changes, pedal spasm, irritability, marked jitteriness.

**TABLE 7.** Complications Associated With Thrombocytopenia

	81 Healthy Infants	25 Ill Infants
Death or permanent serious sequelae*	0	2 (8%)
Serious transient complications†	2 (2%)	3 (12%)
Asymptomatic, but received platelet transfusion	2 (2%)	8 (32%)
Asymptomatic thrombocytopenia (not treated)	5 (6%)	2 (8%)
Normal platelet count or not measured	72 (89%)	10 (40%)

\* Symptomatic pulmonary hemorrhage, intraventricular hemorrhage.

† Petechial rash, rectal bleeding, hematuria, bleeding umbilicus.

bleeding from thrombocytopenia, catheter-related complications, and apnea and bradycardia with cyanosis requiring resuscitation. These complications are common enough that exchange transfusion, even in healthy newborns, should be performed only in nurseries prepared to respond to these adverse events. Because 5% of healthy infants in this study developed symptomatic hypocalcemia, such infants should at a minimum have blood-ionized calcium checked at the first signs of hypocalcemia. However, it should be noted that the administration of supplemental calcium during exchange transfusion is usually ineffective and unnecessary.<sup>14</sup> Because 10% of healthy infants had platelet counts decrease to <50 000 per microliter, such infants should at a minimum be observed closely for petechiae or other signs of bleeding. Invasive procedures such as lumbar puncture or surgery should be delayed until a safe level of platelets has been achieved. Furthermore, both calcium and platelet counts should be checked before repeat transfusions. Catheters should be removed as soon as appropriate, and there should be a high index of suspicion for thrombotic complications. Although apnea, bradycardia, and cyanosis occur rarely during exchange transfusion of healthy infants, cardiorespiratory and oxygen saturation monitoring appear indicated. In ill infants, in addition to the above precautions, routine measurement of both ionized calcium and platelet count after exchange transfusion is indicated, given the occasionally severe complications associated with hypocalcemia and thrombocytopenia in this population.

Because many of the complications of exchange transfusion are probably unavoidable, the best way to reduce complications is to prevent the need for exchange transfusion. The use of effective phototherapy, including optimization of the wavelength and power of the lamps, and maximization of skin light exposure including use of fiberoptic pads, can greatly reduce the need for exchange transfusion. Other innovative approaches for treating jaundice include intravenous gammaglobulin in Rh-sensitized newborns and the administration of tin mesoporphyrin.

In summary, this report indicates that adverse events remain common after exchange transfusion. Some complications are as severe as, or worse than, the bilirubin encephalopathy the exchange transfusion was intended to prevent. These severe complications must be balanced against the benefits of lowering serum bilirubin. Because of the much higher rate of complications in ill infants, the results do not support recommendations to use lower exchange levels in ill infants compared with healthy infants.

#### APPENDIX A. DEATHS POSSIBLY ATTRIBUTABLE TO EXCHANGE TRANSFUSION

Infant 1 was a 26-week gestation, 824-g infant with hyaline membrane disease who was improving until sudden respiratory failure 36 hours after exchange transfusion. The infant died of *Pseudomonas* sepsis 2 days after exchange transfusion, possibly attributable to infectious complications of exchange transfusion.

Infant 2 was a 30-week gestation, 900-g infant with gastroschisis whose exchange was performed through a Broviac catheter in the superior vena cava. After the exchange transfusion, a clot was noted on the catheter. The infant subsequently developed *Staphylococcus aureus* sepsis, *Escherichia coli* sepsis, superior vena cava syndrome, *Candida tropicalis* fungemia, and died 1 month later.

Infant 3 was a 24-week gestation, 630-g infant with severe asphyxia, hyperkalemia, and hypernatremia. During the exchange transfusion, the patient suffered cardiac arrest associated with severe hypocalcemia, and died several days later due to intraventricular hemorrhage and intractable seizures.

Infant 4 was a 33-week gestation, 2435-g infant with respiratory failure from hydrops fetalis. During the exchange transfusion, the infant suffered respiratory deterioration necessitating discontinuation of the exchange transfusion. After the infant died a few days later of respiratory failure and heart block, an autopsy was performed. Necrosis of the cardiac conduction system possibly related to emboli from the exchange transfusion was observed.

Infant 5 was a 35-week gestation, 3380-g infant with mild hyaline membrane disease. Several weeks after two exchange transfusions, the patient developed chronic hepatitis, possibly acquired from blood transfusion, and later died of hepatic encephalopathy.

#### APPENDIX B. PERMANENT SERIOUS SEQUELAE POSSIBLY ATTRIBUTABLE TO EXCHANGE TRANSFUSION

Infant A was a 34-week gestation, 2030-g healthy infant whose umbilical venous catheter was replaced several times for clotting during the exchange transfusion. The next day the infant developed massive pneumoperitoneum, acidosis, and perforation of the splenic flexure of the bowel. The infant underwent resection of necrotic bowel and creation of colostomy, and later underwent surgery for bowel reanastomosis and closure of the colostomy. The infant also developed necrosis of the thumb; this complication and the bowel injury were classified as probable thrombotic complications of exchange transfusion.

Infant B was a 27-week gestation, 1080-g infant who developed chronic aortic obstruction from exchange transfusion through the umbilical artery catheter (and later died of causes unrelated to exchange transfusion).

Infant C was a 27-week gestation, 700-g infant who developed *Staphylococcus epidermidis* sepsis after the third exchange transfusion. The infant required repeated platelet transfusions due to severe thrombocytopenia from multiple exchange transfusions and then suffered intraventricular bleeding, hydrocephalus, and developmental delay.

Infant D was a 32-week gestation, 2630-g infant who had erythroblastosis fetalis, but did not require mechanical ventilation. The infant developed severe thrombocytopenia after exchange transfusion, had a sudden and severe respiratory deterioration attributable to pulmonary hemorrhage, and required intubation and mechanical ventilation. Subsequent tracheal injury resolved after bronchoscopy, but the infant was noted to have global developmental delay.

## REFERENCES

1. Watchko JF, Oski FA. Bilirubin 20 mg/dL = vigintiphobia. *Pediatrics*. 1983;71:660–663
2. Newman TB, Maisels MJ. Evaluation and treatment of jaundice in the term infant: a kinder, gentler approach. *Pediatrics*. 1992;89:809–818
3. American Academy of Pediatrics, Provisional Committee for Quality Improvement and Subcommittee on Hyperbilirubinemia. Practice parameter: management of hyperbilirubinemia in the healthy term newborn. *Pediatrics*. 1994;94:558–565
4. Boggs TR, Westphal MC. Mortality of exchange transfusion. *Pediatrics*. 1960;26:745–755
5. Weldon VV, Odel GB. Mortality risk of exchange transfusion. *Pediatrics*. 1968;41:797–801
6. Panagopoulos G, Valaes T, Doxiadis SA. Morbidity and mortality related to exchange transfusions. *J Pediatr*. 1969;74:247–254
7. Kitchen WH. Neonatal mortality in infants receiving an exchange transfusion. *Aust Paediatr J*. 1970;6:30–40
8. Hovi L, Siimes MA. Exchange transfusion with fresh heparinized blood is a safe procedure. *Acta Paediatr Scand*. 1985;74:360–365
9. Dikshit SK, Gupta PK. Exchange transfusion in neonatal hyperbilirubinemia. *Indian Pediatr*. 1989;26:1139–1145
10. Guaran RL, Drew JH, Watkins AM. Jaundice: clinical practice in 88,000 liveborn infants. *Aust N Z J Obstet Gynaecol*. 1992;32:186–192
11. Keenan WJ, Novak KK, Sutherland JM, Bryla DA, Fetterly KL. Morbidity and mortality associated with exchange transfusion. *Pediatrics*. 1985;75(suppl):422–426
12. Watchko JF, Claassen D. Kernicterus in premature infants: current prevalence and relationship to NICHD Phototherapy Study exchange criteria. *Pediatrics*. 1994;93:996–999
13. Beyer WH, ed. *Handbook of Tables for Probability and Statistics*. 2nd ed. Cleveland, OH: The Chemical Rubber Co; 1968:222–229
14. Maisels JM, Li T, Piechocki JT, Werthman MW. The effect of exchange transfusion on serum ionized calcium. *Pediatrics*. 1974;53:683–686

## Adverse Events Associated With Exchange Transfusion in Healthy and Ill Newborns

J. Craig Jackson

*Pediatrics* 1997;99:e7

DOI: 10.1542/peds.99.5.e7

### Updated Information & Services

including high resolution figures, can be found at:  
<http://pediatrics.aappublications.org/content/99/5/e7>

### References

This article cites 12 articles, 7 of which you can access for free at:  
<http://pediatrics.aappublications.org/content/99/5/e7#BIBL>

### Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:  
<http://www.aappublications.org/site/misc/Permissions.xhtml>

### Reprints

Information about ordering reprints can be found online:  
<http://www.aappublications.org/site/misc/reprints.xhtml>

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



# PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

## **Adverse Events Associated With Exchange Transfusion in Healthy and Ill Newborns**

J. Craig Jackson

*Pediatrics* 1997;99:e7

DOI: 10.1542/peds.99.5.e7

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/99/5/e7>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 1997 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

