



POLICY STATEMENT

Baseball and Softball

abstract

FREE

Baseball and softball are among the most popular and safest sports in which children and adolescents participate. Nevertheless, traumatic and overuse injuries occur regularly, including occasional catastrophic injury and even death. Safety of the athlete is a constant focus of attention among those responsible for modifying rules. Understanding the stresses placed on the arm, especially while pitching, led to the institution of rules controlling the quantity of pitches thrown in youth baseball and established rest periods between pitching assignments. Similarly, field maintenance and awareness of environmental conditions as well as equipment maintenance and creative prevention strategies are critically important in minimizing the risk of injury. This statement serves as a basis for encouraging safe participation in baseball and softball. This statement has been endorsed by the Canadian Paediatric Society. *Pediatrics* 2012;129:e842–e856

INTRODUCTION

The pediatrician needs to have an understanding of baseball and softball. This will allow the pediatrician to offer appropriate counseling and guidance to the many boys and girls, their parents, and members of the sporting community who participate in baseball and softball each year.

Baseball is one of the most popular sports in the United States, with an estimated 8.6 million children ages 6 to 17 participating annually in organized and recreational baseball.¹ Although baseball is a relatively safe sport in comparison with many other athletic activities, highly publicized catastrophic impact injuries from contact with a ball or a bat frequently raise safety concerns.^{2,3} These incidents, as well as the high frequency of shoulder and elbow injuries resulting from overload and overuse, provide the impetus for this review and new guidance to reduce injury risk and improve safety in baseball for 5- to 18-year-old participants. This policy statement replaces the previous statement written in 2001.⁴ This statement has been endorsed by the Canadian Paediatric Society.

Beginning in the early 1990s, epidemiological and injury surveillance research in baseball and softball intensified. Data from these scientific efforts paved the way for organized baseball to create medical advisory committees, which generated policies designed to reduce the risks of injury in baseball and softball. Advances in equipment also continue to offer new opportunities to make the game safer for youth athletes; similarly, the dissemination and use of automatic lightning detectors (which produce a clear and loud warning signal) and automated external defibrillators (AEDs) provide additional means of reducing catastrophic events on the baseball field.^{5,6} Moreover, organized youth baseball coaches, officials, and administrators must remain knowledgeable and

COUNCIL ON SPORTS MEDICINE AND FITNESS

KEY WORDS

balls, bats, commotio cordis, elbow, equipment, helmets, injury, pitch count, safety, shoulder

ABBREVIATIONS

AED—automated external defibrillator
CPSC—Consumer Protection Safety Commission
NEISS—National Electronic Injury Surveillance System
NOCSAE—National Operating Committee on Standard for Athletic Equipment
UCL—ulnar collateral ligament

This document is copyrighted and is property of the American Academy of Pediatrics and its Board of Directors. All authors have filed conflict of interest statements with the American Academy of Pediatrics. Any conflicts have been resolved through a process approved by the Board of Directors. The American Academy of Pediatrics has neither solicited nor accepted any commercial involvement in the development of the content of this publication.

All policy statements from the American Academy of Pediatrics automatically expire 5 years after publication unless reaffirmed, revised, or retired at or before that time.

www.pediatrics.org/cgi/doi/10.1542/peds.2011-3593

doi:10.1542/peds.2011-3593

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2012 by the American Academy of Pediatrics

sensitive to the developmental and skill levels of young baseball players and continue to modify the rules, when necessary, for the safety of the players. This policy statement focuses principally on baseball, but softball will also be considered where relevant literature is available.

PARTICIPATION

There are over a dozen youth baseball organizations whose combined annual participation is nearly 5 million. Little League Baseball reports baseball participation at approximately 2.3 million annually and softball participation at 400 000 per year⁷; the Babe Ruth League (and its 12-and-under Cal Ripken Baseball division) has more than 1 million youth participants annually.⁸ Over 2 million girls between the ages of 12 and 18 compete in fast pitch softball annually. The National Federation of State High School Associations 2009–2010 participation data counted 473 503 baseball players (ages 14–18) and 393 578 softball athletes.⁹

INJURY DATA

There are a variety of sources of data for injuries in youth baseball, which include hospital emergency departments, insurance claims, national catastrophic injury reporting system, and independent researchers.^{1,10–13} Each of these sources captures a limited view of total injuries. Taken together, however, a fairly consistent pattern emerges on the overall injury incidence and relative frequency (prevalence) of baseball injuries compared with other sports. Comprehensive data also come from the high school level reporting systems using certified athletic trainers to record the key injury-related information on this 14- to 18-year-old population.^{14–16}

Prevalence

The US Consumer Product Safety Commission (CPSC) maintains the National

Electronic Injury Surveillance System (NEISS) through a network of approximately 100 participating US emergency departments. From the data collected from this cross section of national emergency departments, the CPSC extrapolates national estimates of injury prevalence in various age groups. The overall prevalence of all sports-related injuries in children younger than 15 years seen in an emergency department is just under a million per year; in 2007, the CPSC estimated that 109 202 baseball- and softball-related injuries among 5- to 14-year-old children were treated in US emergency departments nationally.¹⁷ More specifically, in 2007, the CPSC/NEISS recorded exactly 3343 baseball injuries among 5- to 18-year-old children from the participating US emergency departments (T.J. Schroeder, MS, personal communication, 2009). The frequency distribution of injuries by age followed a bell-shaped curve for ages 5 to 18 years, with the highest number of injuries clustered in children from 11 to 14 years of age (more than 300 injuries per year of age).¹⁷ Because 11- to 14-year-old children represent the majority of participants, valid injury risk comparisons with other age groups cannot be directly determined from these raw numbers alone. Nearly half of the 3343 injuries (44%) involved the head (25% to the face, including eyes and nose; 14% to the head and neck; and 5% to the mouth), with the highest frequency in children 9 to 11 years of age.¹⁷ According to several years of annual NEISS reports, approximately one-fourth to one-third of all youth baseball injuries are to the upper extremities, including the fingers (10%–13%), wrists (4%–5%), and hands (4%–5%). Just under 20% of injuries were to the lower extremities, with knees (5%) and ankles (6%–7%) almost equally affected. Injuries to the trunk and pubic area accounted for approximately 6% to 10% of all injuries.^{1,11,17}

Also in 2007, the NEISS recorded exactly 1188 youth softball injuries in the same 100 participating emergency departments. Girls accounted for 1062 injuries, and boys accounted for 126.¹⁷ For girls, the frequency of injuries by age also followed a bell-shaped curve for ages 5 to 18 years, with the highest numbers of injuries clustered in children from 13 to 16 years of age (more than 130 injuries per year of age). The distribution of injuries by body region was more evenly distributed for girls playing softball compared with boys playing baseball, with 28% of girls' softball injuries involving the head and neck, 35% involving the upper extremities, 31% involving the lower extremities, and 5% involving the trunk and pubic area. The 5 most common areas for injury were the face (14%), ankle (14%), finger (13%), knee (11%), and head and neck (11%).¹⁷ Thirty-nine percent of injuries were classified as contusions, abrasions, lacerations, or hematomas/hemorrhages; 31% were classified as sprains or strains; 21% were classified as fractures, dislocations, or avulsions; 4% were classified as internal organ injuries; and 4% were classified as concussions.¹⁷

Baseball is one of the safest high school sports in the United States with a reported injury rate of 1.26 injuries per 1000 athletic exposures.¹⁴ Baseball athletes have the third lowest rates of injury lasting longer than 1 week among 18 different sports played in high school. However, although the overall injury rate is low, the degree of injury severity is relatively high. In baseball, fractures represent a larger percentage of total injuries than in other sports. Baseball also ranks second highest for the percentage of injuries resulting in a time loss from sport participation longer than 7 days.¹⁴ In high school baseball and softball, as in youth baseball, most injuries to the head and face (48%) and mouth

and teeth (16.0%) are attributed to being hit by a batted ball.¹⁴ More of these head and facial injuries also required surgery (18.0%), compared with other baseball-related injuries (6.8%).

USA Baseball, through its Medical and Safety Advisory Committee, sponsored injury-surveillance research and produced a report on injury patterns in youth baseball in 2008 which concluded that, in younger players, injuries are more often associated with lack of skill, whereas in older players, the greater skill, muscle power, and body size become injurious forces.¹⁵ More detailed study findings were as follows:

1. Across all age groups, pitchers, catchers, and fielders have a relative frequency of injury in a season comparable to batters and base runners, with one exception: as age increases, the proportion of injuries to catchers increases, whereas the proportion of injuries to fielders decreases.
2. The younger the age group, the more frequently injuries occur during practice sessions, not games. Also, in younger age groups, relatively higher frequencies of injuries came from “before/after” baseball play, such as warm-ups or postgame horseplay.
3. Pitcher injuries are also age-related, with noncontact overuse injuries increasing in each older age group. Younger pitchers are more likely to be hit by a batted or thrown ball.
4. Catchers principally are injured in catching a pitch, getting hit by a foul tip, or tagging a base runner. Younger catchers are more frequently injured trying to catch the pitched ball, whereas older catchers are more likely to be injured by foul tips. Although home-plate collisions during a tag play generate a modest but constant pattern of injury for catchers of all ages, these collisions produce a greater injury risk to the older catcher.
5. Batters are most frequently injured by a pitched ball (and by a foul ball as they get older). The younger the batter, the more likely they are to be injured by a swung bat of another player while getting into position. Approximately 10% of batting injuries among older batters are of a noncontact nature, indicating some form of overswinging resulting in an injury to the muscles, bones, or connective tissues of the rib cage.
6. Injuries to base runners occurred almost equally at all bases, with slightly more injuries at home plate than at first base. Two-thirds of these injuries occurred from sliding, and one-third occurred from base running. Base running injuries are not related to age, occurring almost equally from collisions, being hit by a thrown ball, or simply running (such as falling, stepping wrong on the base, or straining a muscle).
7. Fielder injuries occur at the base (33%), in foul territory (10%), and in the field (57%). Younger fielders are more likely to be hit by a batted ball, whereas older fielders are more likely to be injured by colliding with a sliding opponent.

Catastrophic Injuries

Research from the National Amateur Baseball Catastrophic Injury Surveillance Program places the rate of catastrophic injury very low, at approximately 1 injury per 1 million participants annually. Such injuries may be caused by trauma related to participating in the skills of the sport, such as contact with a bat, baseball, or softball (direct catastrophic injury), by a body system failure resulting from exertion while participating in a sport or activity (eg, cardiac collapse or heat stroke) or by a complication that resulted from a nonfatal injury (indirect catastrophic injury). Catastrophic injuries are further classified according to outcome as fatal (athlete

died), nonfatal (athlete left with permanent disability), and serious (athlete experienced severe injury but recovered, to be left with no permanent functional disability).¹² The National Amateur Baseball Catastrophic Injury Surveillance Program data revealed an annual fatality rate of 0.05 per 100 000 participants, 0.03 for nonfatal catastrophic injuries and 0.04 for serious catastrophic injuries.¹⁸ Between 1996 and 2006, deaths in youth baseball averaged just over 2 per year.¹⁸ These deaths occurred from impact to the head resulting in intracranial bleeding and from blunt chest impact, likely prompting ventricular fibrillation or asystole (commotio cordis).^{11,12}

Another study based on statistics compiled by the CPSC¹¹ indicated that there were 88 baseball-related deaths (approximately 4 per year) to children 5 to 18 years old between 1973 and 1995.^{11,19} The injury causing death in 38 cases (43%) was direct-ball impact with the chest (commotio cordis); in 21 cases (24%), the injury causing death was direct-ball contact with the head; in 13 cases (15%), the injury causing death was from impact from bat contact; in 9 cases (10%), the injury causing death was from direct contact with a ball impacting the neck, ears, or throat; and in 7 (8%), the injury causing death was unknown.^{11,19}

Among high school baseball athletes (with approximately 475 000 participating annually), there have been 10 “direct” deaths, 14 “indirect” deaths, 17 nonfatal catastrophic injuries, and 22 serious catastrophic injuries between 1982 and 2008.¹² Direct injury rates are between 2 and 5 times greater in high school than in youth baseball (5–14 years of age) and 4 times greater in college than high school baseball (but still less than 1/100 000 participants annually in any of the 3 categories).

Among high school softball athletes (371 000 annual participants), there has been 1 direct death, no indirect deaths, 2 nonfatal catastrophic injuries, and 1 serious catastrophic injury in the 15 seasons from 1993 and 2008; these rates are approximately one-tenth of those for high school baseball.¹²

Comotio Cordis

Young baseball and softball players who receive direct ball impact to the anterior chest wall over the cardiac silhouette may develop cardiac arrest. Comotio cordis is the second highest cause of death in athletes younger than 14 years and is considered to be only a pediatric problem because of its unique occurrence in children, usually younger than 16 years.^{20–22} Children 5 to 14 years of age may be uniquely vulnerable to this blunt chest impact, because their chest walls are more elastic and more easily compressed.^{20–24}

Although protective gear can be a key preventive measure, it is not always effective. Research has shown that even with protective gear, the fatality rate for comotio cordis is alarmingly high at 90%.^{19–24,25,26} Proper coaching and execution can augment protective equipment. Batters who learn to avoid the ball and turn away from an inside pitch can greatly reduce their risk. This is particularly difficult, however, while bunting and, thus, requires special attention and instruction. The risk to pitchers can also be reduced by teaching proper fielding position after ball release as well as ball avoidance when necessary.¹⁹ The capacity to recognize that a batted ball is coming at you quickly and the ability to react to that event before the ball arrives are crucial traits for protecting these athletes. Speed of processing and reaction time are commonly measured during the computer-based neuropsychological tests given to high

school athletes as a baseline in concussion management programs. Presumably, these are skills that can be improved marginally with proper training and sufficient repetitions. It is estimated that 400 milliseconds is required for a pitcher to complete a protective movement, such as lifting a hand to the chest or face; this corresponds to a batted ball speed of 42 m/second.²⁷ Because balls hit from some metal bats may exceed 42 m/second, performance standards for composite metal bats were developed in 2003 to limit their capacity to transfer force (velocity) to the batted ball.

It is noteworthy that cardiac death caused by comotio cordis is potentially preventable. If cardiac arrest occurs, an immediate and appropriate response may, in many cases, save a young player's life. AEDs are becoming more prevalent in the sports world at athletic complexes and certainly in patrol cars and emergency medical vehicles. The use of these devices, along with cardiopulmonary resuscitation, has become a standard first step in treating cardiac arrest.^{6,28} If no AED is available near the affected child, the emergency medical care system should be activated by using 911 to get an AED to the athlete's side as soon as possible (ideally within 3 minutes). Baseball coaches should be routinely reminded to have a local cellular phone and emergency medical numbers at every youth baseball and softball game and practice in the event of a medical emergency.

Concussion

Concussion in sports continues to garner ever-increasing public attention and concern. Data on high school concussion rates in baseball and softball were generated through the Reporting Information Online system, developed from the National High School Sports-Related Injury Surveillance Study, over

the 5-year period from the 2005–2006 to 2009–2010 school years. The rate of concussions for baseball was 0.2 per 1000 athletic exposures, and for softball the rate of concussions was 0.5 per 1000 athletic exposures. Of the 9 sports surveyed (including football, wrestling, soccer, and basketball), baseball had the lowest rate, and softball was seventh. The top 2 activities associated with concussion in high school boys playing baseball were batting (36.7%) and running the bases (21.5%), whereas for girls playing softball, the top 2 activities were catching (33.3%) and fielding (25.0%).²⁹ The most recent concussion recommendations, including the AAP Clinical Report published in 2010, stress removing an athlete suspected of sustaining a concussion from play immediately with no return to play on the day of injury. Resting the brain is the key to the most rapid resolution of the concussion; such rest includes no physical activity, with minimal cognitive and social activity. The concussed athlete should seek prompt medical attention from an appropriate health professional. Once the athlete is fully free of symptoms and cleared by a physician knowledgeable in the treatment and management of concussion, the athlete should enter a graduated return-to-play protocol to ensure a safe return to sporting activity.³⁰

Summary of Acute Injuries

Although the rates of injury for baseball and softball are low in comparison with other sports, the combination of relatively high severity of injuries that do occur and the very large number of participants in these sports produces a substantial number of individuals with significant injuries each year. Focused media attention on these cases often produces a groundswell for action, even when the scientific evidence for change is lacking.

OVERLOAD CAUSING OVERUSE INJURY OF THE SHOULDER AND ELBOW IN BASEBALL

Overuse arm injuries in youth baseball primarily affect pitchers. The repetitive stress of throwing can lead to muscle fatigue and then to muscle, tendon, and ligament damage. To control the number of these injuries, it is critical that pediatricians, coaches, athletes, and parents understand the anatomy, biomechanics, and kinetics of throwing a baseball pitch.^{31–33}

In throwing, especially pitching, the arm force and speed generated during the full windup motion is astonishingly high (7200 degrees, or 20 revolutions of a circle, per second for professional pitchers). At ball release, the rotator cuff muscles and long head of the biceps contract to decelerate arm motion and prevent the arm from following the ball (“throwing one’s arm out”).³⁴

The shoulder girdle (including the clavicle [collarbone], scapula [shoulder blade], glenohumeral joint, humerus, and all of the various muscular attachments to the thorax and scapula) permits the arm to enjoy remarkable freedom of movement. Greater joint range of motion sacrifices the inherent joint stability that is usually provided by bony articulation or ligamentous limitation. Unlike other joints, such as the knee and ankle, the glenohumeral joint of the shoulder acquires nearly all of its stability during activity from the muscles of the shoulder with little or no assistance from bones or ligaments. Therefore, when the muscles become fatigued (and cannot do their job properly), the shoulder becomes an unstable joint. Pitchers who continue to throw when the arm is fatigued risk serious and possibly permanent injury to their muscles, ligaments, capsules, labra, and bones.³⁵

The scapula plays a critical role in the functioning kinetic chain of the upper

extremity. The scapula is attached to the distal end of the clavicle by the broad acromioclavicular ligament and the 2 coracoclavicular ligaments that arise from the hooklike coracoid. The clavicle attaches proximally to the sternum at the sternoclavicular joint; the sternoclavicular joint is the only joint connection of the entire upper extremity to the main skeleton of the body. The scapula rests on the thorax (chest wall) and is connected through numerous muscles (scapulothoracic joint) that permit the shoulder blade to rise (shrugging shoulders), to glide around the body (protraction), and to glide back toward the spinal column (retraction). Strength and endurance of these extrinsic shoulder girdle muscles, especially the serratus anterior, are essential to ensure safe pitching. These muscles must move the scapula to its appropriate position and stabilize its base before the acceleration phase of the throwing motion can proceed. With excellent scapular muscular control, the athlete can then use the long lever of the upper extremity to throw with power, speed, and accuracy.³⁶

It also has long been recognized, however, that the source of power (velocity) in throwing is part of a larger kinetic chain, which begins in the legs, travels through the pelvis into the torso (“the core”), and finally comes out through the scapula into the upper extremity. Strong legs and especially a strong core help generate a majority of the force behind a pitched ball.³⁷

Relative weakness of the serratus anterior muscle is often responsible for injuries to the rotator cuff muscles, biceps, and elbow (farther down the kinetic chain). Scapular winging (being able to slide one’s hand under the medial or inferior edge of the scapula when relaxed), scapular depression (inferior pole sits lower than the other side or below T-7), and protraction (sits farther away from the vertebral

spines than the opposite side) are noted on inspection. Observing an athlete (from behind) performing the arm component of a “jumping jack” approximately a dozen times will demonstrate poor movement patterns, or scapulothoracic dyskinesia. A second test entails forward flexing the arm to the fully overhead position and slowly lowering the arm 180 degrees to its natural position. Scapulothoracic dyskinesia consists of unsmooth or uncontrolled scapular motion while bringing the arms up or down, including marked scapular winging during descent. Although the athlete may perform the first few repetitions well, scapulothoracic dyskinesia may become evident toward the end of the set, when fatigue sets in.^{36,38}

The scapula also forms part of the glenohumeral joint (true shoulder joint). The scapula’s glenoid is a shallow socket surrounded by a cuplike rim called the labrum, which provides additional stability. The 2 main ligaments of the glenohumeral joint perform most of their stabilizing function when the shoulder is not being used dynamically. During sporting activity and especially during throwing, the 4 rotator cuff muscles and the long head of the biceps are responsible for keeping the humeral head in the glenoid socket. The larger muscles (deltoids, trapezius, pectorals, latissimus dorsi, short head of the biceps, and triceps) provide the powerful dynamic forces to move the arm and generate ball velocity.

The term “Little League shoulder” refers to the widening of the proximal humeral physis resulting from chronic recurrent traction stress across the growth plate. This lesion is felt to be primarily the result of throwing a large number of pitches or maximum-effort throws or insufficient rest between pitching assignments, but can also be related to improper mechanics or premature attempts with certain pitch

types. When unrecognized, or when recognized in an advanced stage of overuse, Little League shoulder can lead to chronic pain with throwing, early shoulder instability, or degenerative arthritis.³⁹

The term “Little League elbow” refers to medial elbow pain in skeletally immature athletes resulting from throwing issues similar to those mentioned in the description of “Little League shoulder.” Pitchers are most likely to be affected by this condition, but it can also occur in other players at positions requiring frequent and forceful throwing. Traction forces occur on the medial elbow and compression forces laterally. The medial traction forces can cause separation or avulsion of the humeral medial epicondyle apophysis and overuse injury to the common flexor tendon. The compression forces laterally can cause collapse and deformity of the distal humerus (capitulum), also known as osteochondritis dissecans in adolescents and Panner disease in children younger than 10 years. Early recognition of symptoms is important to avoid chronic elbow pain, instability, and arthritis.³⁵ The inability to fully straighten the elbow is usually an indication of an overuse elbow injury. Complete tears of the ulnar collateral ligament (UCL) are repaired by “Tommy John” surgery, which is followed by nearly a year of rehabilitation before return to pitching is allowed. UCL tears have risen to nearly epidemic proportions in the past 5 years among youth and high school pitchers, increasing almost 20-fold from the previous decade.^{35,40} When the pitch velocity exceeds 80 miles per hour, the forces experienced at the UCL are near the point of failure (32 Newton meters).^{35,41}

Prevention of Shoulder and Elbow Overuse Injuries in Baseball

In 2006, the Medical and Safety Advisory Committee of USA Baseball set

pitching limits for a season and a calendar year, based on age.⁴² In 2007 and 2008, on the basis of extensive research, Little League Baseball released its pitch count regulation guide in an attempt to reduce the risk of overuse injuries to elbows and shoulders.^{43,44} This guide for parents, coaches, and league officials sets a maximum number of pitches to be thrown in a day on the basis of age as well as the number of rest days between pitching assignments on the basis of number of pitches thrown and age. It replaces the previous regulations, which set a maximum number of innings pitched in a calendar week. Beginning in 2010, Little League tournament pitching regulations became the same as for regular season play, as shown in Tables 1 and 2.^{44,45} It is the responsibility of the coach to know when an athlete has reached his or her pitch count limit. When pitchers exceed recommended pitch count maximums for age for an entire season or a full year, the injury rates for shoulders and elbows increase dramatically.⁴¹

Young pitchers should avoid pitching on multiple teams with overlapping seasons; the guidelines for rest requirements must be enforced across all teams. Youth pitchers should not pitch competitively in more than 8 months in any 12-month period; 3 consecutive months of complete rest from pitching each year is recommended. A pitcher should also not be a catcher for his or her team. Catchers throw even more frequently than pitchers, and although the return throws are only moderately stressful, catchers also make many hard throws to the bases during a game. Thus, playing both positions greatly increases the repetitive stress to the arm.^{35,46} A preseason conditioning program that includes strengthening the core, the rotator cuff, and the shoulder-stabilizing

muscles (scapular stabilizers) also may help reduce throwing injuries. A prospective epidemiological research study among asymptomatic adolescent and preadolescent tennis athletes found those with decreased scapula stabilizing muscular strength (noted by scapula winging, protraction, and depression) demonstrated a higher rate of shoulder injury during the subsequent season.⁴⁷

Teaching proper pitching mechanics may also prevent serious overuse injuries.^{35,42,43,48} Allowing sufficient time during the early part of the season to gradually increase the amount and intensity of throwing may provide young arms an opportunity to adapt to the stresses of throwing. During the off-season and preseason, after allowing for a period of several months of complete rest from throwing, daily throwing of a strictly limited number of pitches may enhance strength development (as regular strength training does for other muscle regions) with little risk of overuse injury or fatigue.^{35,49} A core-stabilizing exercise program will improve strength and should also be adopted for pitchers and hitters.^{35,37,47}

Four keys to successful pitching within the 5- to 14-year age group are (1) the development of a fastball; (2) accuracy of locating pitches (control); (3) development of an off-speed pitch (change-up); and (4) changing pitch speeds and plate locations to keep batters guessing. Conventional wisdom has long considered the curve ball and slider to be stressful to the young elbow, and recommendations to delay their introduction until later years (when the skeletal maturity is more significantly advanced) were common.⁴³ Recent studies challenge those theories and indicate that, when properly thrown, the curve ball may not overly stress the elbow.^{35,50,51} Nevertheless, on the basis of those

TABLE 1 Little League Baseball 2010 Pitching Guidelines: Maximum Pitches per Game

10 y and younger	75 pitches per d
11–12 y	85 pitcher per d
13–16 y	95 pitches per d
17–18 y	105 pitches per d

Source: Little League (<http://www.littleleague.org/media/newsarchive/2009/Sep-Dec/LLTournamentRegularSeasonPitchingRulesMadeSame.htm>).

studies that show increased injury among those who throw curve balls and sliders at early ages, researchers currently continue to recommend delaying introduction of the curve ball until after age 14 or when pubertal development has advanced to the stage when the athlete has started to shave (sliders should be delayed until age 16).⁴¹ The American Academy of Pediatrics endorses this recommendation. Finally, on the basis of the increasing number of elbow ligament surgeries in younger and younger pitchers, it is clear that players, parents, and coaches require more respect for the limits of the developing child's arm to withstand the forces incurred while pitching and should understand that the consequence of overload causing overuse injuries which can permanently damage anatomic structures.³⁵

OVERUSE INJURY FROM SOFTBALL PITCHING

Data on softball injuries were included in the previous sections; in general, these

demonstrate injury patterns similar to baseball with several minor variations. One unique feature of softball, however, is the nature of pitching; pitchers throw underhand from a flat mound. Although the softball windmill pitching motion may be less stressful to the pitcher's shoulder and elbow than throwing a hardball overhand, there are still significant forces placed on various body structures to cause overuse injury. The same concerns exist for arm safety: do not throw when fatigued. As of 2011, inning limitations and rest days for softball windmill pitchers in the various age divisions of Little League apply to tournament play only. Many softball teams rely on fewer pitchers per team than baseball. Before such inning limitations, some pitchers pitched in multiple games during weekend tournaments, running up pitch counts between 1500 and 2000 pitches in a 3-day period.⁵²

The driving force of the windmill softball pitch is in the lower body⁵³; 1 study reported more than 50% of the total kinetic energy of upper extremity during overhead movements is supplied by the trunk and legs.⁵⁴ The softball pitcher engages the gluteal muscles to achieve stabilization of the pelvis, which in turn helps the scapula achieve adequate control.

The lower extremity supports the mechanics of the upper body during the

softball windmill pitch, especially during the single-leg support component; however, the site of failure (injury) may not be located where the problem originates, but at a more remote site (the weakest link). This kinetic chain theory produces “culprits” and “victims” in describing injuries. For example, although 70% of 131 injuries to collegiate softball pitchers were from overuse, only 13% were reported in the lower extremity.⁵⁵ The majority occurred in the shoulder or elbow. Thus, although the “culprit” may be weak gluteal muscles and lack of pelvic stabilization, the athlete may present with shoulder pain, because the “victim” shoulder is inherently “the weakest link.”

The pelvis and torso work to accelerate the segments of the upper extremity in a sequential manner. Scapular retraction is stimulated by ipsilateral hip extension and trunk extension. A stable scapula is vital for optimal rotator cuff function that helps to keep the humerus in the glenoid fossa (thus avoiding impingement).

Accordingly, examination and treatment of shoulder pain in softball windmill pitchers should include an assessment of and rehabilitation for pelvic stability and gluteal strength. Scapular stabilization must also be evaluated and treated as well as the affected shoulder or elbow. Off-season and preseason conditioning programs for softball windmill pitchers should include gluteal strengthening and pelvic stabilization exercises.

ROLE OF EQUIPMENT IN INJURY RISK AND PREVENTION

Playing Equipment

The game of baseball requires playing equipment (ball, bat, gloves, bases), as well as equipment to dress and protect the athlete. Although there are overarching concerns about equipment

TABLE 2 Little League Baseball 2010 Pitching Guidelines: Rest Requirements for Pitchers

Pitchers 14 y and younger	
66 or more pitches in a day	Four (4) calendar days
51–65 pitches in a day	Three (3) calendar days
36–50 pitches in a day	Two (2) calendar days of rest must be observed
21–35 pitches in a day	One (1) calendar day of rest must be observed
1–20 pitches in a day	NO (0) calendar day of rest must be observed
Pitchers 15–18 y	
76 or more pitches in a day	Four (4) calendar days
61–75 pitches in a day	Three (3) calendar days
46–60 pitches in a day	Two (2) calendar days of rest must be observed
31–45 pitches in a day	One (1) calendar day of rest must be observed
1–30 pitches in a day	NO (0) calendar day of rest must be observed

Source: Little League (<http://www.littleleague.org/media/newsarchive/2009/Sep-Dec/LLTournamentRegularSeasonPitchingRulesMadeSame.htm>).

performance and safety that apply to every athlete who plays baseball, some concerns typically apply to only a small segment of players depending on age, developmental level, and skill.^{56,57}

The ball is the cause of most baseball injuries, through being hit by a pitched ball, being struck while attempting to field a batted ball, trying to catch a thrown ball, or being hit by a thrown or batted ball while running the bases. Accordingly, modifications in the hardness and compressibility of baseballs (and softballs) were developed by equipment manufacturers several decades ago for use by children of different ages with the intent of reducing the force of impact by the ball while maintaining adequate performance characteristics; they were known as reduction-in-force balls. These softer balls, however, are alleged to possess a more “lively” bounce than traditional baseballs.^{56–58} The National Operating Committee on Standard for Athletic Equipment (NOCSAE) has developed standards for these softer baseballs (levels 1, 2, and 3),^{59,60} and an expert review panel and other researchers have indicated that softer balls meeting the NOCSAE standard are less likely to cause injury, specifically serious head injury or commotio cordis by impact.^{22–24,60,61} Children with the lowest skill level (in general, those younger than 10 years) should use the lowest-impact NOCSAE-approved balls (level 1). Level 2 low-impact balls are designed for children 10 to 12 years of age with moderate skill levels. Children younger than 10 years of age with moderate skills may use either level 1 or level 2 balls. Level 3 balls are designated for youths older than 12 years and those 10 to 12 years of age with advanced skills.⁶¹

Baseball has the highest number of sports-related eye injuries in children, with the highest incidence in 5- to 14-year-olds (T.J. Schroeder, MS, personal

communication, 2009).^{11,15} Approximately one-third of baseball-related eye injuries result from being struck by a pitched ball; other common causes of eye injuries are attempting to field a batted ball or catch a thrown ball.

The composition of the bat has provoked much discussion in recent years, especially for baseball players older than 15 years. Composite (metal) bats have largely replaced wooden bats.⁶² Initially, the introduction of metal bats led to an increased velocity of the batted ball, putting fielders at greater risk of injury, especially pitchers and third basemen.⁶² Considerable research has been conducted comparing both types of bats, but no consensus exists regarding safety of composite metal bats compared with wooden bats in youth baseball.^{62,63} All bats, wooden or metal, are momentarily deformed upon contact with the baseball; the recoil or restitution characteristics of the bat to its usual shape result in imparting additional velocity to the batted ball. Industry performance standards for composite metal bats were developed in 2003 through the leadership of the National Collegiate Athletic Association and National Federation of State High School Associations. These standards limit the dimensions of the bat and its capacity to transfer force (velocity) to the batted ball (ball exit speed ratio). Bat composition for National Collegiate Athletic Association baseball was required to meet the Batted Ball Coefficient of Restitution guidelines in 2011, and high school baseball will follow suit in 2012.⁶⁴ For Little League Baseball, the maximum bat length is 33 inches and maximum barrel diameter may not exceed 2 1/4 inches. Since 2009, all Little League bats must be labeled with a bat performance factor of 1.15 or lower.^{65,66} Bat performance factor is a measure of a nonwood bat's performance relative to wood bats. For 2011,

Little League International has imposed a moratorium on the use of composite bats for all baseball divisions citing research that found that “composite bats, while they meet the standard when new, can exceed that standard after a break-in process.”⁶⁷ Although the media continue to report anecdotal stories of significant head and facial injuries associated with metal bats, epidemiological injury surveillance studies do not show an increase in the rate of such injuries since metal bats were introduced.⁶²

Equipment-related injuries also include those caused by bases and cleats. Foot and ankle injuries occur regularly in baseball and softball, especially during sliding. Early studies showed that at least 35% of all injuries in softball occurred while sliding feet first into a base, and, in one study, 82% of those injuries resulted in ankle fractures or dislocations.^{68,69} Subsequently, a 2-year study comparing traditional stationary bases and bases capable of disengaging noted a greater than 95% reduction in foot and ankle injuries with breakaway bases.^{70–74} In 2008, Little League baseball and softball mandated the use of breakaway bases.⁷⁴ Youth softball leagues have also incorporated the use of a separate “runner's base” at first base to avoid collisions; the orange “runner's base” sits adjacent to the white regular base, but in foul territory, providing the runner and fielder separate, equal-size bases. Metal spikes are dangerous to basemen covering bases; rubber spikes are preferred for youth baseball.

Protective Equipment

Protective equipment should always be properly fitted, well maintained, and clean; equipment hygiene is important to prevent infections, such as methicillin-resistant *Staphylococcus aureus*. Although not all equipment adaptations have been tested scientifically for

efficacy, experts believe those mentioned in this article will help reduce injuries in softball and baseball players.²⁴

Padded sliding pants worn underneath the baseball pants provide good protection against contusions and abrasions to the hips and thigh when athletes slide feet first into a base. For athletes who slide head and hands first, wearing gloves can reduce hand abrasions; similarly, use of batting gloves improves the grip on the bat and reduces the risk of blisters. Head-first sliding creates the risk of head injuries from collisions with the fielder or from being hit by a thrown ball; upper-extremity injuries can occur from the contact and deceleration forces that occur when the hands of the diving body encounter the stationary base. According to current Little League Baseball rules, a runner may not slide head-first except when retreating to a previously held base.⁷⁵ Chest protectors for batters were introduced in the 1990s to protect the heart from ball impact, which can cause commotio cordis.¹⁹ However, this protective barrier has not been shown to be reliable in either the human experience or in animal laboratory studies.^{22–25,28,63} More study is required to develop and target equipment that will better prevent commotio cordis.²⁴ Recent design modifications indicate that progress is being made, but these advances have not been validated in evidence-based peer-reviewed scientific journal articles or abstracts presented at professional meetings.⁷⁶ Even the chest protectors for catchers, which are effective for blunting the usual forces encountered from pitches, foul tips, throws, and collisions do not reliably prevent commotio cordis.^{24,77}

Hard plastic shell batting helmets have long been an integral part of protecting baseball and softball players from head injuries while batting and

running the bases. The NOCSAE developed standards for helmets and their testing decades ago.⁵⁹ Their design and characteristics have been modified over the years to better protect the eyes, ears, face, mouth, and nose.⁷⁸ In 2009, a new batting helmet was introduced with the capacity to withstand the impact of a 100-mph fastball; it has a layer of expanded polypropylene, the hard, foamlike material used in bicycle helmets.^{79,80}

Face and eye protection can be achieved by securing additional protection to the batting helmet, such as a polycarbonate plastic face guard, metal cage, or a polycarbonate full-face shield. Although full plastic face shields have been demonstrated to reduce injury risk significantly, their acceptance by players and league officials has been mixed, primarily because of compromised visibility through the shield over time.⁵⁶ More recent epidemiological research among high school baseball athletes has led to a call for infielders and pitchers to wear face shields.¹⁴

Eye protection may be worn to reduce the risk for eye injury.^{81–84} This protection may be particularly important for young athletes who have undergone eye surgery or experienced a previous serious eye injury.⁸⁵ Polycarbonate face guards and shields must meet the F910 standard of the American Society of Testing and Materials.^{84,86} Face guards cover the lower part of the face from the tip of the nose to below the chin, directly protecting the teeth and facial bones; the space between the top of the guard and the brim of the helmet is less than the diameter of a baseball, thereby indirectly protecting the eyes without impairing vision. Functionally one-eyed athletes (best corrected vision in the worse eye of less than 20/50) must use one of these protectors when batting and also must protect

their good eye when fielding the ball by using polycarbonate sports goggles that meet American Society of Testing and Materials standard F803.⁸⁶ Some parents may also prefer the use of a face guard or eye goggles when their children are in the field. All athletes who wear glasses or sunglasses while playing baseball or softball should make sure that the lenses and frames meet appropriate safety standards, thus minimizing the risk of breaking or shattering on contact with a baseball or softball.⁸⁶

An essential piece of equipment for all boys playing baseball or softball is use of a hard plastic athletic cup to protect the testicles. Although catchers, pitchers, and infielders are at the greatest risk, testicular injuries can occur to any player while in the field, batting, or running the bases.

The development of triangular-shaped foam pads (“knee savers”) for catchers is another advance in safety equipment that has enjoyed widespread use since its introduction.⁸⁷ These pads fit into the popliteal fossa behind the knee and are designed to minimize the strain on the knee joint while the catcher is in the crouched position. Some batters opt for shin guards to protect their medial distal tibia; another innovation gaining notice is the elbow protective pad for batters.

ENVIRONMENTAL RISKS

Environmental risks include weather-related concerns and field conditions and boundaries (perimeters).

Baseball and softball players are at risk for injury from weather-related factors, including lightning, sun, and heat. Recreational facilities managers, coaches, and umpires need to be well-versed in lightning safety guidelines^{5,88} and prepare an emergency action prevention plan in advance. In particular, lightning presents a unique challenge because of the difficulty in

predicting and appreciating the presence of electrical storms. More and more facilities are installing automatic lightning detectors and alarm systems to ensure that everyone responds appropriately to the presence of lightning. In the absence of an automated warning system, estimating the distance of the storm from the fields can be accomplished by counting the number of seconds between seeing lightning and hearing thunder. Divide the number of seconds by 5 to estimate the distance in miles. A “flash-to-bang” of 30 seconds or less is an indication to move athletes and spectators to a safe area. A preassigned individual should be responsible for the decision to evacuate. In general, a period of 30 minutes should elapse from the last sound of thunder and from the last visible lightning flash before play should be resumed.

Hot and/or humid weather conditions pose significant risk for heat-related illness in children and adolescents. Ambient temperature, relative humidity, wind speed, and solar radiant heat all affect risk for heat illness.^{89,90} The ability to dissipate body heat into the surrounding environment decreases as the ambient temperature rises closer to body temperature. Similarly, as the humidity rises and the air becomes more saturated with water, the ability to cool oneself through evaporation decreases. Wind and sun exposure can damage the skin and eyes. Being acclimatized to the environment, ensuring adequate hydration, wearing a baseball cap, and having a covered dugout are important features of minimizing the risk of sun- and heat-related illness. Recommendations regarding sunscreen, sunglasses, and proper uniforms for practice and games are appropriate for players of all ages. Similarly, in cold weather, athletes need to be properly dressed. When necessary, games should be postponed or

game times altered when weather conditions are extreme.

Fields need to be inspected regularly for hazards, including problems with sprinkler heads, gopher holes in the outfield, or rocky infields. A warning track (usually 15 feet wide) should surround the entire grass or artificial turf playing surface and should meet all recommended specifications; the change in surface texture serves to alert fielders that they are nearing the perimeter barriers, such as when outfielders are going back on long hits to the fence. All fences should be in good repair, especially the backstop and the fence along both foul lines. If a chain-link fence is curled up and the grass becomes high, an athlete may receive a laceration when reaching down to pick up the ball (not noticing the hazard produced by the defective fencing).

Elimination of the on-deck circle, placing dugouts behind a fence, and the wearing of batting helmets by the first-base and third-base coaches have further reduced the risk of injury on the playing field.

DEVELOPMENTAL CONSIDERATIONS

The pediatrician needs to have an understanding of developmental considerations as they apply to baseball and softball participation at various ages. This will allow the pediatrician to offer appropriate counseling and guidance to the many boys and girls, their parents, and members of the sporting community who participate in baseball and softball each year.

Compared with older players, children between 5 and 10 years often have less coordination, slower reaction times, a reduced ability to pitch accurately, and a greater fear of being struck by the ball.^{91,92} Some developmentally appropriate rule modifications, therefore, are advisable for this age group,

including the use of batting tee, a pitching machine, or an adult pitcher. The avoidance of head-first sliding for children younger than 10 years is especially important, because there have been anecdotal reports of rare but serious cervical spine injuries occurring when a young player slides head-first, hitting an opponent with the top of the helmet. This injury is similar to that caused by “spearing” (using the head as the lead object) in football. Feet-first sliding should be taught consistently (as players become developmentally ready) so that all players become comfortable with this key technique of the game. The use of softer balls, as described in a previous section of this statement, is also recommended in this age group to reduce serious impact injury. Youth baseball has long recognized that the game must be modified to meet the developmental level and skills of its players. The distance between bases begins at 45 feet for Tee Ball, moves to 60 feet for Minor League and Little League (Majors) for ages 10 to 12 years, goes to 70 to 75 feet for ages 13 to 15 years, and eventually advances to 90 feet for boys 16 to 18 years of age; similarly, the pitcher’s mound increases from 46 feet, to 50 feet, to 60.5 feet from home plate, respectively. The distances of the outfield fences also grow as the field enlarges. The length and weight of bats is another variable that changes as the youth grow in size, strength, and ability.

Some children are able to track ball movement at early ages and can accurately move to and catch a fly ball or hit live pitching as early as 5 years old; most children, however, do not achieve these capabilities until ages 8 or 9.⁹¹ Therefore, it is appropriate to begin introduction to baseball batting for 5- to 7-year-olds using a ball that is not moving but placed on a tee. For

children who are capable of hitting a thrown pitched ball, however, being forced to hit the tee ball may not enhance their skill development.

Coaches who are not knowledgeable in child development may become frustrated trying to teach basic skills to young boys and girls who are simply not ready to acquire these skills, no matter how many balls are pitched to them or how many fly balls they are asked to catch during practice. For these reasons, Little League baseball tracks its athletes into the minors and the majors for boys 10 through 12, separating those children who are developmentally advanced from those who are developing at the typical rate. It must be remembered, however, that “late bloomers” may ultimately turn out to be the finest athletes of all, despite their slower beginning. Proper instruction and persistent repetition in the fundamentals of baseball are essential for youth to develop the talent and skills required to perform proficiently at a high level during late adolescence and adulthood.

RECOMMENDATIONS

The American Academy of Pediatrics recommends the following:

1. Because baseball and softball for children 5 through 18 years of age are relatively safe sports, participation should be encouraged by pediatricians.
2. Preventive measures should be used to protect young baseball pitchers from throwing injuries. Adequate core strength and scapular muscle strength provide a critical foundation; proper instruction in throwing mechanics, conditioning, and training is another essential component. Delay introduction of the curve ball until after age 14 and the slider until age 16. Three key elements in preventing overuse injuries in pitchers include: (a) age guidelines regarding the number of pitches thrown daily; (b) rest requirements between pitching assignments; and (c) season and yearly total pitch limits.^{38,42,44} Young pitchers should avoid pitching on multiple teams with overlapping seasons; the guidelines for rest requirements must be enforced across all teams. Youth pitchers should not pitch competitively in more than 8 months in any 12-month period; 3 months of rest from pitching are recommended each year. A pitcher should also not be a catcher for his or her team.
3. Parents, coaches, and players should be educated about the early warning signs of elbow and shoulder overuse injuries as well as the importance of parascapular muscle strength. Athletes should cease pitching immediately when signs of arm fatigue or pain occur; they should be encouraged to seek timely and appropriate treatment of significant or persistent pain.³⁵
4. Serious and potentially catastrophic baseball injuries can be minimized by the proper use of available safety equipment. This safety equipment includes: (a) for hitters, approved batting helmets with face protection; (b) for catchers, helmets, masks with throat guards, chest protectors, and shin guards; (c) for all male players, hard plastic athletic cup; and (d) for all baseball and softball players, rubber-spiked soles. In light of the relatively high percentage and severity of head, face, and mouth injuries in baseball, strong consideration should be given to head and facial protection for pitchers and infielders, especially in younger age groups and for less-skilled players. Protective equipment should always be properly fitted, well maintained, and clean.
5. Current data show that chest barriers or protection are not sufficiently effective in preventing commotio cordis; thus, the routine use of these heart protectors is not recommended for baseball players at this time. Catchers must continue to wear approved chest protectors.
6. Coaches and officials must be prepared to activate the emergency response system (call 911) and to obtain rapid access to an AED, ideally in less than 3 minutes, to assist young baseball or softball players who are experiencing cardiac arrest or another medical emergency. Coaches should have a local cellular phone and emergency medical phone numbers available for use at all times.
7. Coaches, parents, umpires, and league officials need to be knowledgeable regarding the cause, prevention, recognition, and response to concussion. New policies exist with excellent free educational and prevention materials, especially from the Centers for Disease Control and Prevention. Furthermore, free on-line trainings are available through the Centers for Disease Control and Prevention (<http://www.cdc.gov/concussion/>) and the National Federation of State High School Associations (www.nfhs.org).
8. Youth baseball and softball players should wear polycarbonate eye protection guards or shields or metal cages on their batting helmets to reduce the risk of eye injury. For functionally one-eyed athletes (best corrected vision in the worse eye of less than 20/50) and those who have undergone eye surgery or experienced previous severe eye injuries, eye protection (one of the 3 types of

face shields or polycarbonate sports goggles) should be required at all times while the athlete is batting or fielding.

9. Children with the lowest skill level (in general, those younger than 10 years of age) should use the lowest-impact NOCSAE-approved balls (level 1). Level 2 low-impact balls are recommended for children 10 to 12 years of age with moderate skill levels. Children younger than 10 years with moderate skills may use either level 1 or level 2 balls. Level 3 balls are designated for youths older than 12 years and those 10 to 12 years of age with advanced skills.
10. Continued research into the relative safety of composite metal versus wooden bats is appropriate and necessary.
11. Awareness of heat and lightning safety is essential for all coaches and officials of Little League baseball and softball. The Web site <http://www.lightningsafety.noaa.gov/outdoors.htm>⁸⁸ and weather forecasts are critical resources for prevention of lightning injuries.

When necessary, games or practices should be postponed or game times altered when weather conditions are extreme.

12. Protective fencing of dugouts and benches and the elimination of the on-deck circle are recommended during games and practices in organized and informal participation. Proper field maintenance is required to minimize injury risks.
13. New developmentally appropriate rule modifications should continue to be implemented when indicated.
14. Surveillance of baseball and softball injuries should be continued. Continual strong support of research is essential to develop other new, improved, and efficacious safety equipment and rule modifications.

LEAD AUTHORS

Stephen G. Rice, MD, PhD, MPH
Joseph A. Congeni, MD

COUNCIL ON SPORTS MEDICINE AND FITNESS EXECUTIVE COMMITTEE, 2010-2011

Teri M. McCambridge, MD, Chairperson
Joel S. Brenner, MD, MPH, Chairperson-Elect

Holly J. Benjamin, MD
Charles T. Cappetta, MD
Rebecca A. Demorest, MD
Mark E. Halstead, MD
Chris G. Koutures, MD
Cynthia R. LaBella, MD
Michele Labotz, MD
Keith J. Loud, MDCM, MSc
Stephanie S. Martin, MD
Amanda K. Weiss Kelly, MD

PAST COUNCIL EXECUTIVE COMMITTEE MEMBERS

Joseph A. Congeni, MD
Andrew J. M. Gregory, MD
Stephen G. Rice, MD, PhD, MPH

LIAISONS

Lisa K. Kluchurosky, MEd, ATC – *National Athletic Trainers Association*
John F. Philpot, MD – *Canadian Paediatric Society*
Kevin D. Walter, MD – *National Federation of State High School Associations*

PAST COUNCIL EXECUTIVE LIAISON

Claire M. A. LeBlanc, MD – *Canadian Paediatric Society*

CONSULTANTS

Michael F. Bergeron, PhD
Robert Cordes, MD
James Raynor, MS, ATC

STAFF

Anjie Emanuel, MPH

REFERENCES

1. Lawson BR, Comstock RD, Smith GA. Baseball-related injuries to children treated in hospital emergency departments in the United States, 1994-2006. *Pediatrics*. 2009;123(6). Available at: www.pediatrics.org/cgi/content/full/123/6/e1028
2. Henson S. Pitcher head injuries to trigger cries for protection. *Yahoo! Sports*. May 28, 2010. Available at: http://sports.yahoo.com/mlb/news;_ylt=Ashw9AFXDSrC4iQ6Md2d-WA85nYcB?slug=sh-headinjuries052710. Accessed December 19, 2010
3. Killion A. Viewpoint: California proposes ban on metal bats after kid suffers head injury. *SI.com*. May 25, 2010. Available at: http://sportsillustrated.cnn.com/2010/writers/ann_killion/05/25/sandberg/index.html?xid=shareFB&hpt=Sbin Accessed December 19, 2010
4. Committee on Sports Medicine and Fitness; American Academy of Pediatrics. Risk of injury from baseball and softball in children. *Pediatrics*. 2001;107(4):782-784
5. Baseball USA, Medical and Safety Advisory Committee. Lightning at the Ballpark. February 28, 2002. Available at: http://web.usabaseball.com/news/article.jsp?ymd=20090813&content_id=6408814&vkey=news_usab&gid. Accessed December 19, 2010
6. Baseball USA, Medical and Safety Advisory Committee. AEDs at the Ballpark. December 2008. Available at: http://web.usabaseball.com/news/article.jsp?ymd=20090813&content_id=6407404&vkey=news_usab&gid. Accessed February 28, 2010
7. Little League Baseball Inc. Participation in Little League Reaches 3-Year High. Available at: www.littleleague.org/media/newsarchive/03_2006/06participation.htm. Accessed December 19, 2010
8. Wikipedia. List of Organized Baseball League – Youth Leagues. Available at: http://en.wikipedia.org/wiki/List_of_organized_baseball_leagues#Youth_leagues. Accessed December 19, 2010
9. National Federation of State High School Athletic Associations. 2009–2010 High School Athletics Participation Data. Available at: www.nfhs.org/participation. Accessed December 19, 2010

10. US Consumer Product Safety Commission. National Electronic Injury Surveillance System. CPSC Document 3002. Available at: www.cpsc.gov/cpscpub/pubs/3002.html. Accessed December 19, 2010
11. Kyle SB. *Youth Baseball Protective Equipment Project: Final Report*. Washington, DC: US Consumer Product Safety Commission; 1996
12. Mueller FO, Cantu RC. National Center for Catastrophic Injury Research. 26th Annual Report. Available at: www.unc.edu/depts/nccsi/. Accessed December 19, 2010
13. Clarke KK. Patterns of Injury in Youth Baseball. November 30, 2008. Available at: http://web.usabaseball.com/news/article.jsp?ymd=20090810&content_id=6354562&vkey=news_usab&gid. Accessed December 19, 2010
14. Collins CL, Comstock RD. Epidemiological features of high school baseball injuries in the United States, 2005-2007. *Pediatrics*. 2008;121(6):1181-1187
15. Powell JW, Barber-Foss KD. Sex-related injury patterns among selected high school sports. *Am J Sports Med*. 2000;28(3):385-391
16. Krajnik S, Fogarty KJ, Yard EE, Comstock RD. Shoulder injuries in US high school baseball and softball athletes, 2005-2008. *Pediatrics*. 2010;125(3):497-501
17. US Consumer Product Safety Commission. NEISS Data Highlights – 2007. Available at: www.cpsc.gov/library/neiss.html. Accessed December 19, 2010
18. Baseball USA, Medical and Safety Advisory Committee. National Amateur Baseball Catastrophic Injury Surveillance Program (NAB-CISP). August 13, 2009. Available at: http://web.usabaseball.com/news/article.jsp?ymd=20090813&content_id=6409874&vkey=news_usab&gid. Accessed December 19, 2010
19. Baseball USA, Medical and Safety Advisory Committee. What is Commotio Cordis in Baseball? November 30, 2008. Available at: http://web.usabaseball.com/news/article.jsp?ymd=20090813&content_id=6410334&vkey=news_usab&gid. Accessed December 19, 2010
20. Link MS, Wang PJ, Maron BJ, Estes NA. What is commotio cordis? *Cardiol Rev*. 1999;7(5):265-269
21. Link MS, Maron BJ, VanderBrink BA, et al. Impact directly over the cardiac silhouette is necessary to produce ventricular fibrillation in an experimental model of commotio cordis. *J Am Coll Cardiol*. 2001;37(2):649-654
22. Janda DH, Bir CA, Viano DC, Cassatta SJ. Blunt chest impacts: assessing the relative risk of fatal cardiac injury from various baseballs. *J Trauma*. 1998;44(2):298-303
23. Link MS, Maron BJ, Wang PJ, Pandian NG, VanderBrink BA, Estes NA III. Reduced risk of sudden death from chest wall blows (commotio cordis) with safety baseballs. *Pediatrics*. 2002;109(5):873-877
24. Link MS, Bir C, Dau N, Madias C, Estes NA, III, Maron BJ. Protecting our children from the consequences of chest blows on the playing field: a time for science over marketing. *Pediatrics*. 2008;122(2):437-439
25. Janda DH, Viano DC, Andrzejak DV, Hensinger RN. An analysis of preventive methods for baseball-induced chest impact injuries. *Clin J Sport Med*. 1992;2(3):172-179
26. Weinstock J, Maron BJ, Song C, Mane PP, Estes NA, III, Link MS. Failure of commercially available chest wall protectors to prevent sudden cardiac death induced by chest wall blows in an experimental model of commotio cordis. *Pediatrics*. 2006;117(4). Available at: www.pediatrics.org/cgi/content/full/117/4/e656
27. Nicholls RL, Elliott BC, Miller K. Impact injuries in baseball : prevalence, aetiology and the role of equipment performance. *Sports Med*. 2004;34(1):17-25
28. Haskell SE, Kenney MA, Patel S, et al. Awareness of guidelines for use of automated external defibrillators in children within emergency medical services. *Resuscitation*. 2008;76(3):354-359
29. Comstock RD. High School RIO: The National High School Sports Injury Surveillance Study. Paper presented at: American Academy of Pediatrics National Conference and Exhibition; San Francisco, CA; October 4, 2010
30. Halstead ME, Walter KD; Council on Sports Medicine and Fitness. American Academy of Pediatrics. Clinical report—sport-related concussion in children and adolescents. *Pediatrics*. 2010;126(3):597-615
31. Fleisig GS. *The Biomechanics of Baseball Pitching* [dissertation]. Birmingham, AL: University of Alabama at Birmingham; 1994
32. Fleisig GS, Andrews JR, Dillman CJ, Escamilla RF. Kinetics of baseball pitching with implications about injury mechanisms. *Am J Sports Med*. 1995;23(2):233-239
33. Fleisig GS, Barrentine SW, Zheng N, Escamilla RF, Andrews JR. Kinematic and kinetic comparison of baseball pitching among various levels of development. *J Biomech*. 1999;32(12):1371-1375
34. Nissen CW, Westwell M, Ounpuu S, et al. Adolescent baseball pitching technique: a detailed three-dimensional biomechanical analysis. *Med Sci Sports Exerc*. 2007;39(8):1347-1357
35. Fleisig GS, Weber A, Hassell N, Andrews JR. Prevention of elbow injuries in youth baseball pitchers. *Curr Sports Med Rep*. 2009;8(5):250-254
36. Sciascia A, Kibler WB. The pediatric overhead athlete: what is the real problem? *Clin J Sport Med*. 2006;16(6):471-477
37. Seroyer ST, Nho SJ, Bach BR, Bush-Joseph CA, Nicholson GP, Romeo AA. Shoulder pain in the overhead throwing athlete. *Sports Health*. 2009;1(2):108-120
38. Putnam CA. Sequential motions of body segments in striking and throwing skills: descriptions and explanations. *J Biomech*. 1993;26(suppl 1):125-135
39. Sabick MB, Kim YK, Torry MR, Keirns MA, Hawkins RJ. Biomechanics of the shoulder in youth baseball pitchers: implications for the development of proximal humeral epiphysiolysis and humeral retrotorsion. *Am J Sports Med*. 2005;33(11):1716-1722
40. Petty DH, Andrews JR, Fleisig GS, Cain EL. Ulnar collateral ligament reconstruction in high school baseball players: clinical results and injury risk factors. *Am J Sports Med*. 2004;32(5):1158-1164
41. Andrews JR. Young throwers. Presented at: American Medical Society for Sports Medicine Annual Meeting; Albuquerque, NM; April 2007
42. Baseball USA, Medical and Safety Advisory Committee. Youth Baseball Pitching Injuries. November 30, 2008. Available at: http://web.usabaseball.com/news/article.jsp?ymd=20090813&content_id=6409508&vkey=news_usab&gid. Accessed December 19, 2010
43. Lyman S, Fleisig GS, Andrews JR, Osinski ED. Effect of pitch type, pitch count, and pitching mechanics on risk of elbow and shoulder pain in youth baseball pitchers. *Am J Sports Med*. 2002;30(4):463-468
44. Little League Baseball Inc. Protecting young pitching arms. The Little League Pitch Count Regulation Guide for Parents, Coaches and League Officials. Updated for 2008. Available at: www.littleleague.org/Assets/old_assets/media/Pitch_Count_Publication_2008.pdf. Accessed December 19, 2010
45. Little League Tournament and Regular Season Pitching Rules Made the Same by Adoption of New Rule. November 24, 2009. Available at: www.littleleague.org/media/newsarchive/2009/Sep-Dec/LLTournamentRegularSeasonPitchingRulesMadeSame.htm. Accessed December 19, 2010
46. Fleisig GS, Andrews JR, Cutter GR, et al. Risk of serious injury for young baseball pitchers: a 10-year prospective study. *Am J Sports Med*. 2011;39(2):253-257
47. Kibler WB, McQueen C, Uhl T. Fitness evaluations and fitness findings in competitive junior tennis players. *Clin Sports Med*. 1988;7(2):403-416

48. Keeley DW, Hackett T, Keirns M, Sabick MB, Torry MR. A biomechanical analysis of youth pitching mechanics. *J Pediatr Orthop*. 2008;28(4):452–459
49. Axe M, Hurd W, Snyder-Mackler L. Data-based interval throwing programs for baseball players. *Sports Health*. 2009;1(2):145–153
50. Dun S, Loftice J, Fleisig GS, Kingsley D, Andrews JR. A biomechanical comparison of youth baseball pitches: is the curveball potentially harmful? *Am J Sports Med*. 2008;36(4):686–692
51. Nissen CW, Westwell M, Ounpuu S, Patel M, Solomito M, Tate J. A biomechanical comparison of the fastball and curveball in adolescent baseball pitchers. *Am J Sports Med*. 2009;37(8):1492–1498
52. Oliver GD, Dwelly PM, Kwon Y-H. Kinematic motion of the windmill softball pitch in pre-pubescent and pubescent girls. *J Strength Cond Res*. 2010;24(9):2400–2407
53. Oliver GD. Powering the windmill: lower body mechanics of softball pitching. *Lower Extremity Review*. 2011;3(6):18–22
54. Kibler WB. Biomechanical analysis of the shoulder during tennis activities. *Clin Sports Med*. 1995;14(1):79–85
55. Hill JL, Humphries B, Weidner T, Newton RU. Female collegiate windmill pitchers: influences to injury incidence. *J Strength Cond Res*. 2004;18(3):426–431
56. Mueller FO, Marshall SW; Baseball USA, Medical and Safety Advisory Committee. Safety equipment used in little league baseball. June 2000; Updated August 11, 2009. Available at: http://web.usabaseball.com/news/article.jsp?ymd=20090811&content_id=6371824&vkey=news_usab&gid. Accessed December 19, 2010
57. Crisco JJT, Greenwald R; National Institute for Sports Science and Safety; USA Baseball, Medical and Safety Advisory Committee. The why and how of bat and ball regulation. June 2000. Available at: http://web.usabaseball.com/news/article.jsp?ymd=20090810&content_id=6354586&vkey=news_usab&gid. Accessed December 19, 2010
58. USA Baseball Medical/Safety Advisory Committee. Use of the reduced impact ball in youth baseball, ages 5–12. Available at: http://web.usabaseball.com/news/article.jsp?ymd=20090813&content_id=6410310&vkey=news_usab&gid November 2008. Accessed February 28, 2010
59. National Operating Committee on Standards for Athletic Equipment, Baseball Helmet Task Force. *Standard Method of Impact Test Performance Requirements for Baseball/Softball Batters: Helmets, Baseballs, and Softballs*. Kansas City, MO: National Operating Committee on Standards for Athletic Equipment; 1991
60. National Operating Committee on Standards for Athletic Equipment. *Standard Performance Specification for Newly Manufactured Youth Baseballs*. NOCSAE Document No. 027. Kansas City, MO: National Operating Committee on Standards for Athletic Equipment; April 2006; modified December 2006
61. Marshall SW, Mueller FO, Kirby DP, Yang J. Evaluation of safety balls and faceguards for prevention of injuries in youth baseball. *JAMA*. 2003;289(5):568–574
62. Wilson D. Effort to get metal to act like wood. *The New York Times*. October 18, 2008. Available at: www.nytimes.com/2008/10/18/sports/baseball/18bats.html. Accessed December 19, 2010
63. Baseball USA, Medical and Safety Advisory Committee. Wood and metal bats. March 27, 2007. Available at: http://mlb.mlb.com/usa_baseball/article.jsp?story=medsafety10. Accessed December 19, 2010
64. The NCAA News and NCAA.com. Bat-testing regulations modified. October 8, 2008. Available at: <http://fs.ncaa.org/Docs/NCAANewsArchive/2008/association-wide/bat-testing%2Bregulations%2Bmodified%2B%2B10-08-08%2B-%2Bncaa%2Bnews.html>. Accessed December 19, 2010
65. Little League Online. Statement on non-wood bats. Available at: www.littleleague.org/media/newsarchive/Unknown_Dates/bats.htm. Accessed December 19, 2010
66. Little League Online. Little League International issues update regarding composite bats: moratorium imposed immediately. December 30, 2010. Available at: www.littleleague.org/media/newsarchive/2010/Sep-Dec/CompositeBatMoratorium.htm. Accessed January 4, 2011
67. Youth-Athlete's Sports Blog. Bat rule changes for 2009 season. Available at: www.youthathlete.org/blog/post/2009/02/26/Bat-Rule-Changes-for-2009-Season.aspx. Accessed December 19, 2010
68. Wheeler BR. Slow-pitch softball injuries. *Am J Sports Med*. 1984;12(3):237–240
69. Nadeau MT, Brown T, Boatman J, Houston WT. The prevention of softball injuries: the experience at Yokota. *Mil Med*. 1990;155(1):3–5
70. Janda DH, Bir C, Kedroske B. A comparison of standard vs. breakaway bases: an analysis of a preventative intervention for softball and baseball foot and ankle injuries. *Foot Ankle Int*. 2001;22(10):810–816
71. Janda DH, Maguire R, Mackesy D, Hawkins RJ, Fowler P, Boyd J. Sliding injuries in college and professional baseball—a prospective study comparing stationary and break-away bases. *Clin J Sport Med*. 1993;3(2):78–81
72. Janda DH, Wojtys EM, Hankin FM, Benedict ME. Softball sliding injuries. A prospective study comparing standard and modified bases. *JAMA*. 1988;259(12):1848–1850
73. Little League Online. Disengage-able base rule. Available at: www.littleleague.org/learn/rules/Disengage-Able_Base_Rule.htm. Accessed December 19, 2010
74. Consumer Product Safety Commission. Baseball safety. CPSC Publication No. 329. Available at: www.cpsc.gov/cpscpub/pubs/329.pdf. Accessed December 19, 2010
75. Little League Playing Rules and Regulations. Rule 7.08 (a) (4). Available at: www.wad4llb.org/wp-content/uploads/2009/01/II_safety_rules.pdf. Accessed December 19, 2010
76. Pro Vest, Inc. Pro Vest baseball batting safety vest test results. Available at: www.pro-vest.com/testresults.cfm. Accessed January 19, 2011
77. Berkson D, Queller H, Holmes N, Yun DS, Sandella B, Sargent T. Commotio cordis in a 17-year-old baseball catcher. *Pediatr Cardiol*. 2010;31(5):689–692
78. National Operating Committee on Standards for Athletic Equipment. Standard test method and equipment used in evaluating the performance characteristics of protective headgear/equipment. NOCSAE Doc. 001-08m08b. Modified December 2008. Available at: www.nocsae.org/standards/pdfs/Standards%20%2709/ND001-08m08b-Drop%20Impact%20Test%20Method.pdf. Accessed December 19, 2010
79. Waldstein D. Building a better batting helmet. *The New York Times*. August 12, 2009: B11–B12
80. Rawlings Inc. S100 batting helmet. Available at: www.rawlings.com. Accessed December 19, 2010
81. Grin TR, Nelson LB, Jeffers JB. Eye injuries in childhood. *Pediatrics*. 1987;80(1):13–17
82. Caveness LS. Ocular and facial injuries in baseball. *Int Ophthalmol Clin*. 1988;28(3):238–241
83. Nelson LB, Wilson TW, Jeffers JB. Eye injuries in childhood: demography, etiology, and prevention. *Pediatrics*. 1989;84(3):438–441
84. American Society for Testing Materials. Standard F910-04. Standard specifications for face guards for youth baseball. Available at: www.astm.org/Standards/F910.htm. Accessed December 19, 2010
85. American Society for Testing Materials. Standard F803-03. Standard specifications for eye protectors for selected sports.

- Available at: www.astm.org/Standards/F803.htm. Accessed December 19, 2010
86. Rice SG; American Academy of Pediatrics Council on Sports Medicine and Fitness. Medical conditions affecting sports participation. *Pediatrics*. 2008;121(4):841–848
87. United States International Trade Commission Rulings and Harmonized Tariff Schedule. NY G82707. Available at: www.faqs.org/rulings/rulings2000NYG82707.html. Accessed December 19, 2010
88. Coaches and sports officials guide to lightning safety. Available at: www.lightningsafety.noaa.gov/resources/CoachGuide.pdf. Accessed December 19, 2010
89. Centers for Disease Control and Prevention. Heat illness among high school athletes—United States 2005–2009. *MMWR Morb Mortal Wkly Rep*. 2010;59(32):1009–1013
90. Bergeron MF, Rice SG, Devore CD; Council on Sports Medicine and Fitness and Council on School Health. Climatic heat stress and exercising children and adolescents. *Pediatrics*. 2011;128(3). Available at: www.pediatrics.org/cgi/content/full/128/3/e741
91. Malina RM. 1988 C.H. McCloy research lecture: children in the exercise sciences. *Res Q Exerc Sport*. 1989;60(4):305–317
92. Malina RM, Bouchard C. *Growth, Maturation and Physical Activity*. Champaign, IL: Human Kinetics; 1991

Baseball and Softball
COUNCIL ON SPORTS MEDICINE AND FITNESS
Pediatrics originally published online February 27, 2012;

Updated Information & Services

including high resolution figures, can be found at:
<http://pediatrics.aappublications.org/content/early/2012/02/22/peds.2011-3593>

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
<http://www.aappublications.org/site/misc/Permissions.xhtml>

Reprints

Information about ordering reprints can be found online:
<http://www.aappublications.org/site/misc/reprints.xhtml>

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®



PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Baseball and Softball
COUNCIL ON SPORTS MEDICINE AND FITNESS
Pediatrics originally published online February 27, 2012;

The online version of this article, along with updated information and services, is
located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/early/2012/02/22/peds.2011-3593>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 345 Park Avenue, Itasca, Illinois, 60143. Copyright © 2012 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®

