

PEDIATRICS

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

COVID-19 in Children: Looking Forward, Not Back

Benjamin Lee, MD, and William V. Raszka Jr, MD

DOI: 10.1542/peds.2020-029736

Journal: *Pediatrics*

Article Type: Solicited Commentary

Citation: Lee B, Raszka Jr WV. COVID-19 in children: looking forward, not back. *Pediatrics*. 2020; doi: 10.1542/peds.2020-029736

This is a prepublication version of an article that has undergone peer review and been accepted for publication but is not the final version of record. This paper may be cited using the DOI and date of access. This paper may contain information that has errors in facts, figures, and statements, and will be corrected in the final published version. The journal is providing an early version of this article to expedite access to this information. The American Academy of Pediatrics, the editors, and authors are not responsible for inaccurate information and data described in this version.

COVID-19 in Children: Looking Forward, Not Back

Benjamin Lee, MD and William V. Raszka Jr., MD

Affiliation: University of Vermont, Larner College of Medicine, Burlington, Vermont

Address correspondence to:

William V. Raszka Jr.
Department of Pediatrics,
Larner College of Medicine, University of Vermont
89 Beaumont Ave
Given Courtyard N210
Burlington VT 05405

Funding Source: No funding was secured for this work.

Potential Conflict of Interest: The authors have no conflicts of interest to disclose.

Abbreviations: CI, confidence interval; COVID-19, coronavirus disease 2019; MIS-C, multisystem inflammatory syndrome in children; RT-PCR, reverse transcription polymerase chain reaction; SAR, secondary attack rate; SARS-CoV-2, severe acute respiratory syndrome coronavirus 2

In the fall of 2020, some of the fiercest debates waged in both academic and public arenas concern the relative ability of children to acquire and transmit SARS-CoV-2, the causative agent of COVID-19—rightfully so, as there are enormous implications for children, parents, and schools. To date, much of the data have come from studies of COVID-19 within household clusters early during the pandemic. In this issue of *Pediatrics*, Laws and colleagues add to this body of literature with a secondary analysis of a previously reported household contact study.¹ They describe infection rates, transmission dynamics, and symptom profiles in a cohort of US children with household SARS-CoV-2 exposure between March and May 2020 in Milwaukee and Salt Lake City.²

The authors found that secondary infection rates in household contacts was similar in children (19/68 contacts; 28%) and adults (36/120 contacts; 30%). Having an infected parent was associated with a marked increase in risk for secondary infection in a child. Possible child-to-adult or child-to-child transmission was observed in 2/10 and 1/6 households, respectively, with potential for such events. Infected children generally had mild symptoms and were less likely than adults to report lower respiratory tract symptoms or loss of taste or smell. The findings provide additional confirmation that the overwhelming majority of children with SARS-CoV-2 infection develop mild symptoms,^{3,4} but question whether children are less susceptible to infection or less likely to transmit SARS-CoV-2. Limitations of the study include convenience sampling and relatively small sample size, as only 33 households analyzed included children.

Household contact tracing studies similar to that conducted by Laws and colleagues arguably provide the best evidence regarding pediatric susceptibility to SARS-CoV-2, where the intensity of exposure between household contacts is higher and more consistent than in non-household settings. While not universal,⁵ the preponderance of data from numerous countries

continues to support the notion that children are less susceptible to infection than adults.⁶⁻¹³ Recent meta-analyses suggests that overall susceptibility in children is approximately half of that of adults, with the greatest effect seen in younger children.^{14,15} In this study, younger children had lower rates of infection: the odds ratio for infection in 5-12 year olds compared with 13-18 year olds was 0.36 (95% CI, 0.13-1.05).² Notably, virtually all previous studies have relied on RT-PCR to detect infection and often only test symptomatic contacts. That four pediatric infections in this study could only be detected by antibody seroconversion suggests that RT-PCR-based case detection (symptoms-based or otherwise) in children may underestimate true infection prevalence.

At the population level as well, children continue to be underrepresented among SARS-CoV-2 infections. Numerous large-scale studies have indicated that children, particularly children <10 years, have much lower rates of infection than do adults.¹⁶⁻¹⁹ In the US however, between August 27 and September 10 pediatric SARS-CoV-2 infections increased from 9.5% to 10% of all infections, reflecting a months-long summer trend where regions experienced severe outbreaks and physical distancing recommendations were relaxed or ignored as testing capacity increased.²⁰ While concerning, children under 18 represent 22.3% of the US population,²¹ and this increase has been driven primarily by infections in older children.

The role of children in transmission of SARS-CoV-2 is similarly problematic. This study found that 20% and 17% of households had potential child-to-adult transmission and child-to-child transmission, respectively, but numbers are too small to draw definitive conclusions. Despite some reports of likely onward household transmission from infected children,^{22,23} most studies suggest that children appear less likely to transmit compared with adults or present as index cases in household clusters.^{7,13,19,24-26} A study from South Korea generated considerable

alarm when it reported that children 10-19 years old appeared as likely as adults to transmit infection to household contacts.²⁷ However, a subsequent, more detailed, analysis of 107 pediatric index cases and their household contacts found definitive evidence of only one instance of onward transmission from a teenager, giving a household SAR of 0.5%.²⁸

A significant limitation of household cluster data is that most were generated during school closures, when opportunities for non-household exposure to SARS-CoV-2 among children were scarce. Newer data--reporting experiences in summer camps, childcare settings, and schools (both pre-closure and post-reopening) are helping to fill this gap. Importantly, in areas with low prevalence rates and appropriate mitigation policies, children occasionally become infected but there have been no significant outbreaks. For example, schools or child care programs systems in Ireland, Australia, France, Singapore, Germany, and Rhode Island reported no to little facility-based transmission of SARS-CoV-2 by children despite the presence of infected children.²⁹⁻³⁴ However, in areas with widespread community transmission or less strict mitigation procedures, large outbreaks have occurred.³⁵⁻³⁸ Consistently, these outbreaks reveal that infected adults are typically responsible for introducing virus into these settings, questionable testing strategies enable outbreak initiation, and inconsistent use of masks or cloth facial coverings makes containment of the virus challenging.

So where does this leave us now? First, children clearly are capable of acquiring and transmitting SARS-CoV-2. Second, the preponderance of current data still indicates that children have reduced susceptibility and infectivity compared to adults, though this requires further monitoring as increased testing capacity and relaxation of community mitigation may continue to diminish the magnitude of these differences, which were so stark early during lockdown periods.

Third, the importance of mitigation measures, especially the use of masks, including among children, is now incontrovertible.³⁹

Moving forward, there remains a critical need for more high-quality pediatric SARS-CoV-2 research. Studies in children are often limited by small sample sizes and binning into convenient age cohorts that belie important differences in biology and behavior. The starkest example is to group all children <18 as a single age cohort, when there is ample evidence that younger children and older teens represent completely different patient populations in the context of SARS-CoV-2. For example, in the same region of France, higher rates of potential school-based transmissions were apparent in high schools compared to primary schools.^{38,40} A major step forward in pediatric SARS-CoV-2 research would be universal establishment of reasonable age strata to enable more appropriately powered and comparable studies. Elucidating the mechanisms responsible for differences in symptomatology, susceptibility, and infectivity between adults and children will remain important. However, at this point we also need to shift our focus towards the interventions most important for minimizing transmission of SARS-CoV-2 to and from children, understanding the pathogenesis of MIS-C, and advocating for appropriate pediatric clinical trials for SARS-CoV-2 vaccine candidates.

As SARS-CoV-2 continues its inexorable march through susceptible populations, we must remember that there is no setting on Earth guaranteed to be safe from SARS-CoV-2. With that sobering recognition, we must work to fulfill the medical, academic, social, and emotional needs of children, despite knowing that providing such care cannot ever be completely free of infectious disease risk. As has been shown in the US and around the world, with appropriate mitigation strategies, we can successfully minimize (although not eliminate) the risk of COVID-19. We fear that one day, we will look back on this terrible pandemic and recognize the extent to

which we have failed our children, by being more afraid of their infection and transmission risks than of the prospect of letting them down precisely when they needed us most.

References

1. Lewis NM, Chu VT, Ye D, Connors EE, Gharpure R, Laws RL, et al. Household Transmission of SARS-CoV-2 in the United States. *Clin Infect Dis*. 2020. doi:10.1093/cid/ciaa1166.
2. Laws RL, Chancey RJ, Rabold EM, et al. Symptoms and Transmission of SARS-CoV-2 among Children — Utah and Wisconsin, March-May 2020. IN PRESS this issue *Pediatrics*. 2020.
3. Team CC-R. Coronavirus Disease 2019 in Children - United States, February 12-April 2, 2020. *MMWR Morb Mortal Wkly Rep*. 2020;69(14):422-426. doi:10.15585/mmwr.mm6914e4.
4. Dong Y, Mo X, Hu Y, et al. Epidemiology of COVID-19 Among Children in China. *Pediatrics*. 2020;145(6). doi:10.1542/peds.2020-0702.
5. Bi Q, Wu Y, Mei S, et al. Epidemiology and transmission of COVID-19 in 391 cases and 1286 of their close contacts in Shenzhen, China: a retrospective cohort study. *Lancet Infect Dis*. 2020;20(8):911-919. doi:10.1016/S1473-3099(20)30287-5.
6. Cheng HY, Jian SW, Liu DP, Ng TC, Huang WT, Lin HH. Contact Tracing Assessment of COVID-19 Transmission Dynamics in Taiwan and Risk at Different Exposure Periods Before and After Symptom Onset. *JAMA Intern Med*. 2020;180(9):1156-1163. doi:10.1001/jamainternmed.2020.2020.
7. Jing Q-L, Liu M-J, Yuan J, et al. Household Secondary Attack Rate of COVID-19 and Associated Determinants. *medRxiv* 2020. doi:10.1101/2020.04.11.20056010.
8. Li W, Zhang B, Lu J, et al. The characteristics of household transmission of COVID-19. *Clin Infect Dis*. 2020. doi:10.1093/cid/ciaa450.
9. Rosenberg ES, Dufort EM, Blog DS, et al. COVID-19 Testing, Epidemic Features, Hospital Outcomes, and Household Prevalence, New York State—March 2020. *Clin Infect Dis*. 2020. doi:10.1093/cid/ciaa549.
10. Somekh E, Gleyzer A, Heller E, et al. The Role of Children in the Dynamics of Intra Family Coronavirus 2019 Spread in Densely Populated Area. *Pediatr Infect Dis J*. 2020;39(8):e202-e204. doi:10.1097/inf.0000000000002783.
11. Wang Y, Tian H, Zhang L, et al. Reduction of secondary transmission of SARS-CoV-2 in households by face mask use, disinfection and social distancing: a cohort study in Beijing, China. *BMJ Glob Health*. 2020;5(5). doi:10.1136/bmjgh-2020-002794.

12. Wang Z, Ma W, Zheng X, Wu G, Zhang R. Household transmission of SARS-CoV-2. *J Infect.* 2020;81(1):179-182. doi:10.1016/j.jinf.2020.03.040.
13. Dattner I, Goldberg Y, Katriel G, et al. The role of children in the spread of COVID-19: Using household data from Bnei Brak, Israel, to estimate the relative susceptibility and infectivity of children. *medRxiv.* 2020. doi:10.1101/2020.06.03.20121145.
14. Viner RM, Mytton OT, Bonell C, et al. Susceptibility to SARS-CoV-2 Infection Among Children and Adolescents Compared With Adults: A Systematic Review and Meta-analysis. *JAMA Pediatr.* 2020. doi:10.1001/jamapediatrics.2020.4573
15. Goldstein E, Lipsitch M, Cevik M. On the effect of age on the transmission of SARS-CoV-2 in households, schools and the community. *medRxiv.* 2020. doi:10.1101/2020.07.19.20157362.
16. Gudbjartsson DF, Helgason A, Jonsson H, et al. Spread of SARS-CoV-2 in the Icelandic Population. *N Engl J Med.* 2020;382(24):2302-2315. doi:10.1056/NEJMoa2006100.
17. Pollan M, Perez-Gomez B, Pastor-Barriuso R, et al. Prevalence of SARS-CoV-2 in Spain (ENE-COVID): a nationwide, population-based seroepidemiological study. *Lancet.* 2020;396(10250):535-544. doi:10.1016/S0140-6736(20)31483-5.
18. Report on the Epidemiological Features of Coronavirus Disease 2019 (COVID-19) Outbreak in the Republic of Korea from January 19 to March 2, 2020. 2020 Mar 16. Report No.: 1011-8934.
19. Children and COVID-19. The Dutch National Institute for Public Health and the Environment (RIVM). Updated September 18, 2020. Available at: https://www.rivm.nl/en/novel-coronavirus-covid-19/children-and-covid-19?hash=77642299-5104-4ee2-8e4f-1647bdfb414a&utm_medium=social&utm_source=facebook. Accessed September 18, 2020.
20. Children and COVID-19: State Data Report: A joint report from the American Academy of Pediatrics and the Children's Hospital Association, version 9/10/20. Available at: <https://services.aap.org/en/pages/2019-novel-coronavirus-covid-19-infections/children-and-covid-19-state-level-data-report/>. Accessed 09/18/20.
21. QuickFacts, United States United States Census Bureau. July 1, 2019. Available at: <https://www.census.gov/quickfacts/fact/table/US>. Accessed online 09/21/20.
22. Fateh-Moghadam P, Battisti L, Molinaro S, et al. Contact tracing during Phase I of the COVID-19 pandemic in the Province of Trento, Italy: key findings and recommendations. *medRxiv* 2020. doi:10.1101/2020.07.16.20127357.
23. Mannheim J, Gretsch S, Layden JE, Fricchione MJ. Characteristics of Hospitalized Pediatric COVID-19 Cases - Chicago, Illinois, March - April 2020. *J Pediatric Infect Dis Soc.* 2020. doi:10.1093/jpids/piaa070.
24. Lee B, Raszka WV, Jr. COVID-19 Transmission and Children: The Child Is Not to Blame. *Pediatrics.* 2020;146(2). doi:10.1542/peds.2020-004879.

25. Maltezou HC, Vorou R, Papadima K, et al. Transmission dynamics of SARS-CoV-2 within families with children in Greece: A study of 23 clusters. *J Med Virol*. 2020. doi:10.1002/jmv.26394.
26. Zhu Y, Bloxham CJ, Hulme KD, et al. Children are unlikely to have been the primary source of household SARS-CoV-2 infections. *medRxiv*. 2020. doi:10.1101/2020.03.26.20044826 %J.
27. Park YJ, Choe YJ, Park O, et al. Contact Tracing during Coronavirus Disease Outbreak, South Korea, 2020. *Emerg Infect Dis*. 2020;26(10). doi:10.3201/eid2610.201315.
28. Kim J, Choe YJ, Lee J, et al. Role of children in household transmission of COVID-19. *Arch Dis Child*. 2020. doi:10.1136/archdischild-2020-319910.
29. Heavey L, Casey G, Kelly C, Kelly D, McDarby G. No evidence of secondary transmission of COVID-19 from children attending school in Ireland, 2020. *Euro Surveill*. 2020;25(21). doi:10.2807/1560-7917.ES.2020.25.21.2000903.
30. Macartney K, Quinn HE, Pillsbury AJ, et al. Transmission of SARS-CoV-2 in Australian educational settings: a prospective cohort study. *Lancet Child Adolesc Health*. 2020. doi:10.1016/S2352-4642(20)30251-0.
31. Danis K, Epaulard O, Benet T, et al. Cluster of Coronavirus Disease 2019 (COVID-19) in the French Alps, February 2020. *Clin Infect Dis*. 2020;71(15):825-832. doi:10.1093/cid/ciaa424.
32. Yung CF, Kam KQ, Nadua KD, et al. Novel coronavirus 2019 transmission risk in educational settings. *Clin Infect Dis*. 2020. doi:10.1093/cid/ciaa794.
33. Armann JP, Unrath M, Kirsten C, Lueck C, Dalpke A, Berner R. Anti-SARS-CoV-2 IgG antibodies in adolescent students and their teachers in Saxony, Germany (SchoolCoviDD19): very low seroprevalence and transmission rates. *medRxiv*. 2020. doi:10.1101/2020.07.16.20155143 %J.
34. Link-Gelles R, DellaGrotta AL, Molina C, et al. Limited Secondary Transmission of SARS-CoV-2 in Child Care Programs - Rhode Island, June 1-July 31, 2020. *MMWR Morb Mortal Wkly Rep*. 2020;69(34):1170-1172. doi:10.15585/mmwr.mm6934e2.
35. Szablewski CM, Chang KT, Brown MM, et al. SARS-CoV-2 Transmission and Infection Among Attendees of an Overnight Camp - Georgia, June 2020. *MMWR Morb Mortal Wkly Rep*. 2020;69(31):1023-1025. doi:10.15585/mmwr.mm6931e1.
36. Stein-Zamir C, Abramson N, Shoob H, et al. A large COVID-19 outbreak in a high school 10 days after schools' reopening, Israel, May 2020. *Euro Surveill*. 2020;25(29). doi:10.2807/1560-7917.ES.2020.25.29.2001352.
37. Lopez AS, Hill M, Antezano J, et al. Transmission Dynamics of COVID-19 Outbreaks Associated with Child Care Facilities - Salt Lake City, Utah, April-July 2020. *MMWR Morb Mortal Wkly Rep*. 2020;69(37):1319-1323. doi:10.15585/mmwr.mm6937e3.
38. Fontanet A, Tondeur L, Madec Y, et al. Cluster of COVID-19 in northern France: A retrospective closed cohort study. *medRxiv*. 2020. doi:10.1101/2020.04.18.20071134.

39. Chu DK, Akl EA, Duda S, et al. Physical distancing, face masks, and eye protection to prevent person-to-person transmission of SARS-CoV-2 and COVID-19: a systematic review and meta-analysis. *Lancet*. 2020;395(10242):1973-1987. doi:10.1016/S0140-6736(20)31142-9.
40. Fontanet A, Grant R, Tondeur L, et al. SARS-CoV-2 infection in primary schools in northern France: A retrospective cohort study in an area of high transmission. *medRxiv*. 2020. doi:10.1101/2020.06.25.20140178.

COVID-19 in Children: Looking Forward, Not Back

Benjamin Lee and William V. Raszka Jr.

Pediatrics originally published online October 8, 2020;

Updated Information & Services

including high resolution figures, can be found at:
<http://pediatrics.aappublications.org/content/early/2020/10/06/peds.2020-029736.citation>

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
<http://www.aappublications.org/site/misc/Permissions.xhtml>

Reprints

Information about ordering reprints can be found online:
<http://www.aappublications.org/site/misc/reprints.xhtml>

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®



PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

COVID-19 in Children: Looking Forward, Not Back

Benjamin Lee and William V. Raszka Jr.

Pediatrics originally published online October 8, 2020;

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/early/2020/10/06/peds.2020-029736.citation>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 345 Park Avenue, Itasca, Illinois, 60143. Copyright © 2020 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®

