

PEDIATRICS

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

COVID-19 Transmission in US Child Care Programs

Walter S. Gilliam, PhD, Aryn A. Malik, PhD, Mehr Shafiq, BS, Madeline Klotz, BA, Chin Reyes, PhD, John Eric Humphries, PhD, Thomas Murray, MD, PhD, Jad A. Elharake, BS, David Wilkinson, JD, Saad B. Omer, MBBS, MPH, PhD, FIDSA

DOI: 10.1542/peds.2020-031971

Journal: *Pediatrics*

Article Type: Regular

Citation: Gilliam WS, Malik AA, Shafiq M, et al. COVID-19 transmission in US child care programs. *Pediatrics*. 2020; doi: 10.1542/peds.2020-031971

This is a prepublication version of an article that has undergone peer review and been accepted for publication but is not the final version of record. This paper may be cited using the DOI and date of access. This paper may contain information that has errors in facts, figures, and statements, and will be corrected in the final published version. The journal is providing an early version of this article to expedite access to this information. The American Academy of Pediatrics, the editors, and authors are not responsible for inaccurate information and data described in this version.

COVID-19 Transmission in US Child Care Programs

Walter S. Gilliam^a, PhD, Aryn A. Malik^{b,c}, PhD, Mehr Shafiq^{b,d}, BS, Madeline Klotz^a, BA, Chin Reyes^a, PhD, John Eric Humphries^e, PhD, Thomas Murray^{f,g}, MD, PhD, Jad A. Elharake^{b,h}, BS, David Wilkinson^{i,a}, JD, Saad B. Omer^{b,c,h,j}, MBBS, MPH, PhD, FIDSA

Affiliations: ^aYale Child Study Center, Yale School of Medicine, New Haven, CT; ^bYale Institute for Global Health, New Haven, CT; ^cYale School of Medicine, New Haven, CT; ^dMailman School of Public Health, Columbia University, New York, NY; ^eDepartment of Economics, Yale University, New Haven, CT; ^fDepartment of Pediatrics, Yale School of Medicine, New Haven, CT; ^gYale-New Haven Children's Hospital, New Haven, CT; ^hYale School of Public Health, New Haven, CT; ⁱTobin Center for Economic Policy, Yale University, New Haven, CT; ^jYale School of Nursing, New Haven, CT

Address correspondence to: Walter S. Gilliam, Yale Child Study Center, Yale School of Medicine, 310 Prospect Street, New Haven CT 06511, walter.gilliam@yale.edu, 203-432-4575.

Conflict of Interest Disclosures: All authors report no conflicts of interest.

Funding/support: All phases of this study were supported by the Andrew & Julie Klingenstein Family Fund, Esther A. & Joseph Klingenstein Fund, Heising-Simons Foundation, W.K. Kellogg Foundation, Foundation for Child Development, Early Educator Investment Collaborative, and Scholastic Inc.

Role of Funder/Sponsor: The funders/sponsors did not participate in the work.

Article Summary: This case-controlled study of child care providers is the first to report COVID-19 transmission risk in U.S. child care programs.

What's Known on This Subject: Though often limited by small sample sizes and conducted when child transmission opportunities were limited, studies have suggested that young children are less likely to transmit COVID-19. There are no published studies of COVID-19 transmission in child care.

What This Study Adds: Within the context of considerable infection mitigation efforts in U.S. child care programs, exposure to child care during the early months of the U.S. pandemic was not associated with elevated risk for COVID-19 transmission to providers.

Contributors' Statement Page

Prof Gilliam conceptualized the study, designed the study, conducted the literature search, was involved in aspects of data collection and analysis, and drafted the initial manuscript.

Dr Malik designed the study, conducted the literature search, analyzed data, contributed to data interpretation, and contributed to writing and critical revision of the manuscript.

Ms Mehr analyzed data, contributed to data interpretation, and contributed to critical revision of the manuscript.

Ms Klotz led data acquisition and development of the online survey tool, analyzed data, contributed to data interpretation, and contributed to critical revision of the manuscript.

Dr Reyes designed the study, analyzed the data, contributed to data interpretation, and contributed to critical revision of the manuscript.

Profs Humphries and Murray designed the study and contributed to data interpretation, and contributed to critical revision of the manuscript.

Mr Elharake conducted the literature search, contributed to data interpretation, and contributed to writing and critical revision of the manuscript.

Mr Wilkinson contributed to data interpretation and critical revision of the manuscript.

Prof Omer designed the study, contributed to the analytic approach, contributed to data interpretation, and contributed to critical revision of the manuscript.

All authors approved the final manuscript as submitted and agree to be accountable for all aspects of the work.

Abstract

Objective

Central to the debate over school and child care reopening is whether children are efficient COVID-19 transmitters and likely to increase community spread when programs reopen. We compared COVID-19 outcomes in child care providers who continued to provide direct in-person child care during the first three months of the U.S. COVID-19 pandemic versus those who did not.

Methods

Data were obtained from U.S. child care providers (N=57,335) reporting whether they had ever tested positive or been hospitalized for COVID-19 (N=427 cases), along with their degree of exposure to child care. Background transmission rates were controlled statistically, and other demographic, programmatic, and community variables were explored as potential confounders. Logistic regression analysis was used in both unmatched and propensity score matched case-control analyses.

Results

No association was found between exposure to child care and COVID-19 in both unmatched (odds ratio [OR], 1.06; 95% confidence interval [CI], 0.82 to 1.38) and matched (OR, 0.94; 95% CI, 0.73 to 1.21) analyses. In matched analysis, being a home-based provider (as opposed to center-based) was associated with COVID-19 (OR, 1.59; 95% CI, 1.14 to 2.23), but showed no interaction with exposure.

Conclusion

Within the context of considerable infection mitigation efforts in U.S. child care programs, exposure to child care during the early months of the U.S. pandemic was not associated with elevated risk for COVID-19 transmission to providers. These findings must be interpreted only within the context of background transmission rates and the considerable infection mitigation efforts implemented in child care programs.

The COVID-19 pandemic has had an unprecedented impact on child care and schools. Within one week of the World Health Organization's declaration of a pandemic, 107 countries had implemented national school closures.¹ Within three weeks, the number had grown to 194 countries, impacting 91% of the world's school-age children.² When schools closed in the U.S., so did many child care programs, demonstrated by the loss of more than 35% of jobs in the child care industry between February and April 2020.³ These school and child care closures have been controversial regarding their benefits versus costs. Precautionary closure of child care programs was reasonable, given considerable evidence that these programs may be significant vectors for viral spread.⁴ However, several studies have indicated that school and child care closures may have had little impact on slowing the spread of COVID-19.^{1,5,6} Furthermore, child care and school closures may result in several negative consequences, such as child care providers' loss of jobs and wages, parents' inability to return to work, and children's diminished educational, social, and nutritional opportunities.⁷

Pediatricians are key informers to parents about safety issues regarding child care and school attendance during the pandemic. Central to the debate over child care and school reopening is the uncertainty regarding whether children are efficient transmitters of COVID-19. Although there is a relatively infrequent risk for multisystem inflammatory syndrome in children, children appear to be far less likely to be infected and more likely to be asymptomatic or paucisymptomatic.⁸ On the other hand, because viral loads may be similar in symptomatic and asymptomatic COVID-19 carriers including children,^{9,10} there may be a high risk for transmission to other children and adults through asymptomatic children in settings serving large numbers of children, such as child care and schools. Unfortunately, most studies examining pediatric transmission risk are limited

by small sample sizes of children and were conducted during widespread child care and school closures, when child transmission opportunities were greatly limited.¹¹

This study compares COVID-19 rates in child care providers who continued providing child care during the first three months of the COVID-19 pandemic to those who did not, controlling for key individual, programmatic, and community characteristics which may have motivated closure decisions. We conducted a large-scale multi-state canvass of the U.S. child care workforce to identify a sufficient number of cases, given the cumulative incidence of about 0.8% in the U.S. adult population during the first three months of the U.S. outbreak. This is the first known study to estimate directly COVID-19 transmission within child care programs, while addressing current research limitations by focusing on transmission opportunity through children actively participating in out-of-home programs and therefore not shielded from COVID-19.¹²

METHODS

Sample and Procedures

To provide a wide canvassing of child care providers, Qualtrics¹³ survey links were emailed to individuals beginning May 22, 2020, with three rounds of reminders until the survey closed on June 8, 2020. Participant recall period for program practices was 8 weeks (from April 1 to the median survey administration date of May 27). Survey links were emailed to potential respondents through a variety of contact lists consisting mostly of individuals associated with the child care industry. These lists are maintained by two large national child care organizations (Child Care Aware of America and National Association for the Education of Young Children), as well as through various state child care workforce registries coordinated by the National Workforce Registry Alliance. Of the 41 state registries coordinated by National Workforce

Registry Alliance, 28 states agreed to participate in this study, 11 were unable to secure permissions quickly enough to participate, and two declined.

Data were included from participants self-identifying as child care providers, consenting to the study, satisfying inclusion criteria (pre-pandemic working as child care providers in direct contact with children), and providing all data necessary to determine both COVID-19 outcome and child care exposure. The research protocol was approved by the Yale University Institutional Review Board, and all participants indicated their informed consent electronically at the beginning of the survey. Participants were offered entry into a raffle to select 20 winners of a \$500 gift card. See Table 1.

Variables

Outcome. The key outcome was whether the respondent reported having COVID-19. A value of 1 (case) was assigned if the respondent indicated that they had been either: (a) tested positive for COVID-19 at least once or (b) hospitalized for COVID-19. A value of 0 (non-case) was assigned if neither of these conditions was true.

Exposure. Respondents were asked whether their program closed near the beginning of the pandemic (either as required by government or voluntarily), whether the program reopened, or whether the program closed at any time due to a suspected or confirmed case of COVID-19. Exposure was coded as either 1 (exposed to child care during the pandemic) or 0 (not exposed to child care during the pandemic). A value of 1 was assigned where the respondent indicated that during the pandemic their program remained open, closed but reopened, or closed due to a suspected or confirmed case of COVID-19. A value of 0 was assigned where the respondent indicated that their program was closed during the pandemic or where the respondent indicated

that during the pandemic they either did not report to work or only worked remotely. Child care providers working in open child care programs responded to a series of questions regarding typical group sizes and various infectious disease mitigating strategies their program may have employed during April 2020. See Table 2.

Confounders. All other variables in Table 1 were included as potential confounders. County-level variables were computed for each respondent's county based on their child care program's reported zip code or (if program zip code was not provided) IP address from where the online survey was completed. COVID-19 cumulative death rates were calculated as of May 27, 2020 (the survey median administration date) as a more reliable proxy for community infection rates relative to test-dependent case rates.¹⁵ County-level household median income was obtained from the most recent American Community Survey.¹⁶ County-level variables were trichotomized into proportionally equal thirds.

Given research demonstrating the importance of non-pharmaceutical interventions at reducing SARS-CoV-2 transmission,¹⁴ personal COVID-19 health precautions were measured across three factors. Personal COVID-19 health precautions were measured utilizing a 10-item dichotomized (yes/no) checklist of whether the provider followed during their non-work life common health recommendations. Principal component analysis yielded an interpretable 3-factor solution, accounting for 54.21% of the total variance. Factor 1 = Avoiding Social Interactions, Factor 2 = Avoiding High-Risk Situations and Travel, Factor 3 = Masks, Handwashing, and Distancing (scored 0-3, 0-4, and 0-3, respectively). See Table 3 for items and factor loadings.

RESULTS

Participants

A total of 94,390 survey links were accessed by individuals self-identifying as child care providers across all 50 U.S. states, the District of Columbia and Puerto Rico. Of these individuals, 82,741 satisfied inclusion criteria and consented to the study, and 57,335 (69.3%) chose to participate by providing data necessary to determine both COVID-19 outcome and child care exposure. See Table 1 for sample descriptive data. Smallest numbers of respondents were obtained from South Dakota ($n = 29$), Delaware ($n = 67$), and Utah ($n = 69$); and largest numbers were obtained from California ($n = 9,189$), Florida ($n = 5,232$), and Ohio ($n = 5,076$). See Table 4 for a comparison of the analytic sample demographics versus national estimates for child care providers.

Of received surveys, 30.1% of respondents accessed the surveys via a unique link emailed by the researchers, and 69.9% accessed the surveys via an anonymous link emailed by one of the various state child care workforce registries. An indicator for the unique versus anonymous link was not significant when entered as a covariate in the full logistic regression model (point estimate, 0.93; 95% confidence interval [CI], 0.69 to 1.25; $P = 0.62$). Also, tests of selective participation showed only negligible differences across gender, race, and ethnicity ($|\phi_{(c)}|$ ranging from 0.01 to 0.04), with the exception of age where non-participants ($M = 42.07$; $SD = 12.89$) were younger than participants ($M = 45.97$; $SD = 12.31$; $P < 0.001$; $d = 0.31$).

Child care conditions during the COVID-19 pandemic

Approximately half of respondents (51.4%) reported that their child care program closed near the beginning of the COVID-19 pandemic and remained closed until the time of the survey. Of the

48.6% employed in child care programs that either did not close or reopened during the pandemic, 8.6% reported that their program later closed due to a confirmed or suspected case of COVID-19.

Respondents reported that child care programs that remained open during the early months of the pandemic were operating with smaller than typical group sizes and with considerable degrees of infection mitigation efforts in place. Child group sizes across all open programs were on average smaller than typical for U.S. child care (center-based programs, $M = 7.57$, $SD = 4.94$; home-based programs, $M = 5.62$, $SD = 3.20$). Most children (81.1%) served in these open programs were under six-years old. See Table 5 for details. Respondents reported very high rates (> 90%) of frequent staff and child handwashing and daily disinfecting of indoor surfaces and fixtures. Most respondents reported surface and fixture disinfecting at least three times daily. Symptom screening, cohorting (not mixing children and items between child groups) and social distancing measures also were frequently employed. Far fewer reported daily mask wearing of staff (35.2%) or children two years and older (11.8%). See Table 2 for details.

Outcome

To assess the robustness of the findings, we analyzed results using binary logistic regression analysis in two different approaches: (a) an unmatched analysis predicting case versus non-case, controlling for covariates and (b) a case-controlled approach using propensity score matching.

In the unmatched analysis, we predicted case ($N = 427$; 0.7%) versus non-case ($N = 56,908$; 99.3%), controlling for all variables listed in Table 1 and indicators for the state where the child care program is located. Results indicated no association between COVID-19 outcome and exposure to child care (odds ratio [OR], 1.06; 95% CI, 0.82 to 1.38; $P = 0.66$). (See Table 6.)

COVID-19 was associated with high levels of county-level cumulative per capita COVID-19 deaths (OR, 1.60; 95% CI, 1.19 to 2.15; $P = 0.00$). Among covariates, race and ethnicity were associated with COVID-19, specifically being American Indian/Alaskan Native (OR, 2.37; 95% CI, 1.46 to 3.86; $P = 0.00$), Latinx (OR, 2.13; 95% CI, 1.63 to 2.77; $P < 0.001$), or African American (OR, 1.97; 95% CI, 1.53 to 2.53; $P < 0.001$). Personal COVID-19 health precautions also were associated with COVID-19, specifically avoiding high-risk situations and travel was a protective factor (OR, 0.72; 95% CI, 0.62 to 0.84; $P < 0.001$). Avoiding social interactions, such as family gatherings and restaurants was a risk factor (OR, 1.27; 95% CI, 1.13 to 1.44; $P = 0.001$), perhaps because opportunities to attend these were dependent on local transmission rates and business closures. To assess whether covariates significantly moderated the relationship between exposure and outcome, interaction terms for each with exposure were entered into the model, and none reached significance (P values ranging from 0.09 to 0.96). A sensitivity analysis was employed to determine whether results differ when directors were removed from the sample. Odds ratios for exposure were similar between the full sample (OR, 1.06; 95% CI, 0.82 to 1.38; $P = 0.66$) and with directors removed (OR, 1.08; 95% CI, 0.79 to 1.49; $P = 0.62$).

In the matched case-control analysis, we further tested the association between exposure to child care and COVID-19 outcome using propensity scores for exposure. We matched controls to cases using 1:50 random matching without replacement with caliper width set to $0.2 \times \text{SD}$ of the logit function.¹⁷ Values used in the propensity score include age, race, ethnicity, gender, director status, county-level COVID-19 death rate, county-level median household income, personal COVID-19 health precaution measures (3 continuous variables), and indicator variables for the state where the child care program is located. Results similarly were not indicative of an association between COVID-19 and exposure to child care (OR, 0.94; 95% CI, 0.73 to 1.21; $P =$

0.64). As opposed to the non-matched analysis, being a home-based child care provider was associated with COVID-19 outcome (OR, 1.59; 95% CI, 1.14 to 2.23; $P < 0.01$), but showed no interaction with exposure. See Table 7.

DISCUSSION

This is the first large-scale study of COVID-19 transmission in child care programs. The study utilized a large sample of U.S. child care providers, and results were robust to different analytic approaches and to various tests of exposure interaction effects. Overall, we found no evidence of child care being a significant contributor to COVID-19 transmission to adults. This finding is consistent with previous studies showing a lack of association between school closures and transmission rates.^{6,18-20}

COVID-19 outcome rates were higher among those who identified as American Indian/Alaskan Native, African American/Black, or Latinx, as well as those working in counties with high COVID-19 death rates. Conversely, avoiding situations with high risk of infection (e.g., people who are sick, traveling to high infection areas, social events) was a protective factor. None of these covariates, however, interacted with exposure to child care, suggesting a lack of association between child care exposure and COVID-19 outcome regardless of these other factors.

These findings must be interpreted within the context of infection mitigation practices within U.S. child care programs during the early months of the COVID-19 pandemic. Child care providers reported small group sizes, averaging about eight children per group in centers and six children in home-based programs. Also, respondents reported very high rates ($> 90\%$) of both frequent handwashing and daily disinfecting of indoor surfaces and fixtures, and most respondents reported surface and fixture disinfecting at least three times daily. Symptom screening, cohorting and social distancing measures also were frequently employed. Although it

is unclear whether any of these mitigational efforts significantly reduced transmission, there is no way to know whether these findings would hold in the absence of these infection mitigation efforts. Likewise, these results should not be applied to K-12 schools or universities, where students and contexts vary significantly from child care.

Even after adjusting for other variables, community-level transmission remained a significant predictor of child care providers testing positive or being hospitalized for COVID-19, highlighting the importance of reopening child care programs only when background transmission rates are low and decreasing (e.g., the World Health Organization threshold of < 5% positivity for at least 14 days).²¹ Although this study provides evidence that child care may pose negligible threat to community transmission, communities may pose a considerable threat to child care when background transmission rates are high.²² Conversely, when background transmission rates are low and managed, transmission within child care programs implementing reasonable infection mitigation efforts may be rare.²³

Although our study focused only on adults as the COVID-19 transmission end point—potentially from children to adults and adults to adults—it is also important to acknowledge the possible transmissions of COVID-19 from adults to children and children to children, which were not measured in our study. Children represent 9.3% of all COVID-19 cases in the U.S. (as of August 20, 2020),²⁴ and it is evident that adults are more likely to transmit the virus to children than children are to adults.¹² Due to this likelihood, child care workers should wear face coverings to protect children in child care programs and prevent potential transmission. Furthermore, protective measures against COVID-19 in child care centers are needed as the spread of COVID-19 from child care workers to children may lead to children's family members, including those

most vulnerable to the virus (e.g. the elderly and individuals with underlying medical conditions), contracting COVID-19 from their children or grandchildren.^{14,25,26}

As limitations, the sample is not fully representative. Although the odds ratios were very close to one, it is important to acknowledge the relatively large confidence intervals. Variations in state contexts and policies were controlled statistically, but important policy-level variables may not have been adequately controlled. These data were obtained during Spring, and Fall 2020 transmission dynamics may be different. This study depends upon known infection when it is possible that some providers had asymptomatic infection. Therefore, current case counts are probably underestimates, and results could change, if asymptomatic cases were known/counted. Finally, reliance on self-reported test results may bias findings toward null; however, the large sample size should allow detection of even small effects.

In conclusion, exposure to child care during the early months of the U.S. COVID-19 pandemic was not associated with elevated odds for COVID-19 among child care workers. These findings, however, must be interpreted within the context of both community transmission rates and child care infection mitigation efforts.

Acknowledgments

This study was funded by The Andrew & Julie Klingenstein Family Fund, Esther A. & Joseph Klingenstein Fund, Heising-Simons Foundation, W.K. Kellogg Foundation, Foundation for Child Development, Early Educator Investment Collaborative, Scholastic Inc, Yale Institute for Global Health, and Tobin Center for Economic Policy at Yale University. Invaluable assistance with obtaining child care provider contact information was provided by the National Workforce Registry Alliance (and its network of state child care workforce registries), Child Care Aware of America, and National Association for the Education of Young Children. Drs. Amalia Londono Tobon and Adrián Cerezo Caballero provided Spanish translations and back translations of the survey measures and recruitment information. Louise Scrivani provided project administrative assistance. Alicia Alonso, Catherine Chang, Renee Dauerman, Stella FitzGerald, Harleen Kaur, Emma Knight, and Helen Mooney provided assistance in qualitative data categorization of respondent comments.

References

1. Viner RM, Russell SJ, Croker H, et al. School closure and management practices during coronavirus outbreaks including COVID-19: a rapid systematic review. *Lancet Child Adolesc Health*. 2020;4:397–404. doi:10.1016/S2352-4642(20)30095X.
2. Hildenwall, H., Luthander, J., Rhedin, S., et al. (2020). Paediatric COVID-19 admissions in a region with open schools during the two first months of the pandemic. *Acta Paediatr*. 2020;00:1–3. doi:10.1111.apa.15432.
3. U.S. Bureau of Labor Statistics. Accessed August 23, 2020. <https://beta.bls.gov/dataViewer/view/timeseries/CES6562440010>
4. Hebbelstrup Jenson B, Jokelainen P, Nielson ACY, et al. Children attending day care centers are a year-round reservoir of gastrointestinal viruses. *Scientific Reports*. 2019;9:3286. doi:10.1038/s41598-019-40077-9.
5. Zhang J, Litvinova M, Liang Y, et al. Changes in contact patterns shape the dynamics of the COVID-19 outbreak in China. *Science*. 2020. doi:10.1126/science.abb8001.
6. Ferguson N, Laydon D, Nedjati-Gilani G, et al. Impact of non-pharmaceutical interventions (NPIs) to reduce COVID-19 mortality and healthcare demand. London: Imperial College London; 2020.
7. Esposito S, Principi N. School closure during the coronavirus disease 2019 (COVID-19) pandemic: an effective intervention at the global level? *JAMA Pediatr*. 2020. doi:10.1001/jamapediatrics.2020.1892.
8. Wu Z, McGoogan JM. Characteristics of and important lessons from the coronavirus disease 2019 (COVID-19) outbreak in China: summary of a report of 72,314 cases from the Chinese Center for Disease Control and Prevention. *JAMA*. 2020;323(13):1239–1242. doi:10.1001/jama.2020.2648.
9. Zou L, Ruan F, Huang M, et al. SARS-CoV-2 viral load in upper respiratory specimens of infected patients, *N Engl J Med*. 2020;382:1177–1179. doi:10.1056/NEJMc2001737.
10. Yonker LM, Neilan AM, Bartsch Y, et al. Pediatric SARS-CoV-2: clinical presentation, infectivity, and immune responses. *Pediatrics*. 2020. doi:10.1016/j.peds.2020.08.037.

11. Park YJ, Choe YJ, Park O, et al. Contact tracing during coronavirus disease outbreak, South Korea, 2020. *Emerg Infect Dis.* 2020;26(10). doi:10.3201/eid2610.201315.
12. Lee B, Raszka WV. COVID-19 transmission and children: the child is not to blame. *Pediatrics.* 2020;146(2):e2020004879. doi:10.1542/peds.2020-004879.
13. Qualtrics. Provo, UT. 2019. <https://www.qualtrics.com>
14. Ngonghala CN, Iboi E, Eikenberry S, et al. Mathematical assessment of the impact of non-pharmaceutical interventions on curtailing the 2019 novel coronavirus. *Mathematical Biosciences.* 2020;325:108364. doi:10.1016/j.mbs.108364.
15. Dong E, Du H, Gardner L. An interactive web-based dashboard to track COVID-19 in real time. *Lancet Inf Dis.* 2020;20(5):533–534. doi:10.1016/S1473-3099(20)30120-1.
16. U.S. Census Bureau. 2014-2018 American Community Survey 5-year public use microdata samples; 2019.
17. Austin PC. Optimal caliper widths for propensity-score matching when estimating differences in means and differences in proportions in observational studies. *Pharm Stat.* 2011;10(2):150–161. doi:10.1002/pst.433.
18. Davies NG, Klepac P, Liu Y, et al. Age-dependent effects in the transmission and control of COVID-19 epidemics. *Nature Med Letters.* 2020. doi:10.1038/s41591-020-0962-9.
19. Zachariah P, Johnson CL, Halabi KC, et al. Epidemiology, clinical features, and disease severity in patients with coronavirus disease 2019 (COVID-19) in a children’s hospital in New York City, New York. *JAMA Pediatr.* 2020;e202430. doi:10.1001/jamapediatrics.2020.2430.
20. Rajmil L. Role of children in the transmission of the COVID-19 pandemic: a rapid scoping review. *BMJ Paediatrics Open.* 2020;4:e000722. doi:10.1136/bmjpo-2020-0000722.
21. World Health Organization. Public health criteria to adjust public health and social measures in the context of COVID-19: annex to considerations in adjusting public health and social measures in the context of COVID-19; 2020. <https://www.who.int/publications/i/item/public-health-criteria-to-adjust-public-health-and-social-measures-in-the-context-of-covid-19>
22. Szablewski CM, Chang KT, Brown MM, et al. SARS-CoV-2 transmission and infection among attendees of an overnight camp – Georgia, June 2020. *MMWR Morb Mortal Wkly Rep.* 2020;69:1023–1025. doi:10.15585/mmwr.mm6931e1.
23. Link-Gelles R, DellaGrotta AL, Molina C, et al. Limited secondary transmission of SARS-CoV-2 in child care programs—Rhode Island, June 1-July 31, 2020. *MMWR Morb Mortal Wkly Rep.* ePub:21 August 2020. doi:10.15585/mmwr.mm6934e2.
24. American Academy of Pediatrics, Children’s Hospital Association. Children and COVID-19: state data report; 2020. <https://services.aap.org/en/pages/2019-novel-coronavirus-covid-19-infections/children-and-covid-19-state-level-data-report>
25. Wang X, Ferro EG, Zhou G, et al. Association between universal masking in a healthcare system and SARS-CoV-2 positivity among health care workers. *JAMA.* 2020;324(7):703–704. doi:10.1001/jama.2020.12897.
26. [Jing QL, Li MJ, Zhang ZB, et al. Household attack rate of COVID-19 and associated determinants in Guangzhou, China: a retrospective cohort study. *Lancet Infect Dis.* 2020. doi:10.1016/S1473-3099\(20\)30471-0.](https://doi.org/10.1016/S1473-3099(20)30471-0)

Table 1. Descriptive Statistics for All Variables for Cases, Non-Cases in the Logistic Regression Analysis, and Matched Controls in the Case-Control Analysis

| | Cases (N=427) | Non-Cases (N=56,908) | Matched Controls (N=21,350) |
|--|------------------|-------------------------|-----------------------------------|
| Exposure [%] | | | |
| No | 52.2 | 49.1 | 51.9 |
| Yes | 47.8 | 50.9 | 48.1 |
| Race/Ethnicity [%]¹ | | | |
| American Indian/Alaskan Native | 4.6 | 1.9 | 2.0 |
| Asian | 2.1 | 3.5 | 3.7 |
| African American/Black | 23.4 | 14.7 | 14.8 |
| Native Hawaiian/Other Pacific Islander | 0.9 | 0.5 | 0.5 |
| White | 54.3 | 72.1 | 71.7 |
| Prefer to not answer (Race) | 18.7 | 9.6 | 9.6 |
| Hispanic/Latinx/Spanish Origin | 34.2 | 18.6 | 18.2 |
| Prefer to not answer (Ethnicity) | 2.8 | 2.1 | 2.1 |
| Gender [%] | | | |
| Female | 96.0 | 97.0 | 97.0 |
| Male | 3.3 | 2.3 | 2.3 |
| Non-binary | 0.0 | 0.2 | 0.2 |
| Prefer not answer | 0.7 | 0.5 | 0.5 |
| Age (in years) [M (SD)] | 45.71 (12.49) | 45.97 (12.30) | 46.19 (12.31) |
| Child care Type [%] | | | |
| Center-based ² | 65.8 | 71.4 | 72.2 |
| Family-based/Home-based | 34.2 | 28.6 | 27.8 |
| Director/Owner (of center-based) [%] | | | |
| No | 78.7 | 73.4 | 74.2 |
| Yes | 21.3 | 26.6 | 25.8 |
| Personal COVID-19 Health Precautions [M (SD)] | | | |
| Masks/Handwashing/Distancing (0-3) | 2.86 (0.40) | 2.77 (0.52) | 2.80 (0.49) |
| Avoiding Social Interactions (0-3) | 2.34 (0.89) | 2.11 (1.04) | 2.13 (1.03) |
| Avoiding High-Risk Situations & Travel (0-4) | 2.64 (0.74) | 2.71 (0.67) | 2.71 (0.66) |
| County COVID-19 Cumulative Death Rate [%]³ | | | |
| Low (0-0.0561 deaths/1,000) | 24.6 | 33.6 | 31.2 |
| Moderate (0.0564-0.2180 deaths/1,000) | 24.1 | 34.2 | 33.0 |
| High (0.2184-13.5248 deaths/1,000) | 51.3 | 32.3 | 35.8 |
| County Household Median Income [%]³ | | | |
| Low (\$13,242-\$54,976) | 27.6 | 33.6 | 32.4 |
| Moderate (\$54,979-\$65,010) | 36.3 | 32.3 | 33.8 |
| High (\$65,027-\$136,268) | 36.1 | 34.2 | 33.8 |

¹ Race/Ethnicity options were “check all that apply;” resulting proportions may not add to 100%.

² Center-based child care included (a) for-profit centers (32.5%), (b) non-profit centers (27.4%), (c) school-based (16.7%), (d) Head Start (13.9%), (e) drop-in centers (2.0%), and (f) other (7.4%).

³ County-level data were trichotomized into proportionally equal thirds.

Table 2. Infection Control Efforts Reported in Open Child care Programs during April 2020 (N = 20,550)

| | Percent Reporting Each Effort Every Day | | |
|--|---|-------------|--------------|
| | Once Daily | Twice Daily | Thrice Daily |
| SCREENING | | | |
| Child Symptom Screening | 78.4 | 40.7 | |
| Child Temperature Checks | 76.1 | 33.3 | |
| Staff Symptom Screening | 73.9 | 35.4 | |
| Staff Temperature Checks | 69.3 | 28.3 | |
| DISINFECTING | | | |
| All Indoor Surfaces | 90.2 | 73.9 | 56.5 |
| All Doorknobs, Bathroom Fixtures, Handles | 90.3 | 71.1 | 54.3 |
| All Toys, Books, Classroom Materials | 81.7 | 55.2 | 39.6 |
| No Child-Accessible Items that Are Hard to Disinfect | 75.0 | | |
| HAND HYGEINE | | | |
| All Children Washed Hands Frequently | 92.7 | | |
| All Staff Washed Hands Frequently | 90.2 | | |
| MASKS & PERSONAL PROTECTIVE EQUIPMENT | | | |
| All Children (≥ 2 years old) Wore Facial Masks | 11.8 | | |
| All Adults Wore Facial Masks | 35.2 | | |
| All Adults Wore Aprons/Changed Clothes before Work | 61.7 | | |
| All Adults Wore Gloves when Handling Children | 33.2 | | |
| COHORTING | | | |
| No Mixing of Children between Child Groups | 53.6 | | |
| No Sharing Items between Child Groups | 67.3 | | |
| SOCIAL DISTANCING | | | |
| Seating/Cots ≥ 6 Feet Apart | 68.0 | | |
| Staggered Arrival and Departure Times | 51.2 | | |
| All Children Curbside Drop-off/Pickup | 78.8 | | |
| All Child Belongings Separated | 85.8 | | |
| No Eating "Family Style" or Sharing Food | 49.4 | | |

Note. Percentages indicate the proportion of providers reporting that a specific infection control effort was employed either every day or (for some efforts) twice or thrice per day.

Table 3. Personal COVID-19 Health Precautions: Principal Component Analysis with Factor Loadings

| | Factor 1 | Factor 2 | Factor 3 |
|---|-------------|-------------|-------------|
| ITEMS | | | |
| Asked family and friends not to visit | .79 | - | - |
| Avoided extended family and friends even if not symptomatic | .77 | - | - |
| Avoided eating outside of home | .55 | - | - |
| Avoided close contact with people who were sick | - | .79 | - |
| Avoided traveling to high COVID-19 infection places | - | .71 | - |
| Avoided social events would normally attend | - | .52 | - |
| Canceled business trips, social trips, vacations | - | .47 | - |
| Tried to maintain at least 6 feet from others when outside home | - | - | .75 |
| Facial covering/mask almost always when outside home | - | - | .60 |
| Frequent hand washing/sanitizing when outside home | - | - | .57 |
| FACTOR STATISTICS | | | |
| Eigen Value | 3.36 | 1.06 | 0.99 |
| % Variance Accounted (from rotated sums of squared loadings) | 22.0 | 18.8 | 13.4 |
| Cronbach Alpha | 0.75 | 0.86 | 0.82 |

Note. Factor loadings from Kaiser-normalized varimax rotation, suppressing < .45.

Table 4. Comparison of Total Analytic Sample versus National Estimates of Child Care Provider Demographics and County-Level Descriptions

| | Total Analytic Sample | National Estimates |
|---|-----------------------|--------------------|
| Age [M] | 45.97 | 38.70 |
| Male [%] | 2.3 | 6.5 |
| Race/Ethnicity | | |
| American Indian/Alaskan Native [%] | 2.0 | 0.8 |
| Asian [%] | 3.5 | 3.9 |
| African American/Black [%] | 14.7 | 15.1 |
| White [%] | 72.0 | 69.9 |
| Hispanic/Latinx/Spanish Origin [%] | 18.7 | 22.0 |
| County COVID-19 Deaths per 1,000 [M] | 0.34 | 0.30 |
| County Household Median Income [M\$] | 64,362 | 63,179 |

Note. Age, Gender, and Race/Ethnicity comparisons are between the total analytic sample (N = 57,335) versus national child care provider estimates based on the 2018 American Community Survey (ACS). ACS data are weighted national averages for individuals with an occupation of "childcare workers" (occupation code 4600) between the ages of 18 and 65. For the ACS, Asian = "Chinese" + "Japanese" + "Other Asian or Pacific Islander;" Hispanic/Latinx/Spanish Origin = "Mexican" + "Puerto Rican" + "Cuban" + "Other (Hispanic)." County COVID-19 Deaths per 1,000 are the mean deaths per 1,000 individuals for the counties of survey respondents versus the U.S. national average. County Household Median Income for survey respondents is the mean of the median annual household income of the counties of the survey respondents constructed using the 2018 five-year ACS data. County Household Median Income for the national child care provider estimate is the mean median income for the county in 2018 reported by U.S. Census.

Table 5. Percentage of Children by Age Served by Respondents Open during the Pandemic

| | All Providers | Center-Based Providers | Home-Based Providers |
|--------------------------------|------------------|---------------------------|-------------------------|
| Infants (under 1 year) | 8.3 | 8.2 | 8.3 |
| Toddlers (1-2 years) | 29.7 | 30.6 | 29.4 |
| Preschoolers (3-5 years) | 43.1 | 50.5 | 40.0 |
| School-Age (6 years and older) | 18.9 | 10.8 | 22.3 |

Table 6. Adjusted Logistic Regression Unmatched Analysis Predicting Case (N=427) Versus Non-case (N=56,908)

| | OR | 95% Wald CI | | P Value |
|---|------|------------------|------|---------|
| Exposure to Child care | | | | |
| No | | <i>Reference</i> | | |
| Yes | 1.06 | 0.82 | 1.38 | 0.66 |
| Race/Ethnicity | | | | |
| White | | <i>Reference</i> | | |
| American Indian/Alaskan Native | 2.37 | 1.46 | 3.86 | 0.001 |
| Asian | 0.83 | 0.42 | 1.64 | 0.60 |
| African American/Black | 1.97 | 1.53 | 2.53 | <0.001 |
| Native Hawaiian/Other Pacific Islander | 1.86 | 0.65 | 5.29 | 0.25 |
| Prefer not to answer (Race) | 1.59 | 1.17 | 2.16 | 0.003 |
| Hispanic/Latinx/Spanish Origin | 2.13 | 1.63 | 2.77 | <0.001 |
| Prefer not to answer (Ethnicity) | 1.31 | 0.69 | 2.50 | 0.41 |
| Gender¹ | | | | |
| Female | | <i>Reference</i> | | |
| Male | 1.60 | 0.93 | 2.75 | 0.09 |
| Prefer not to answer | 0.97 | 0.29 | 3.23 | 0.98 |
| Age | 1.00 | 0.99 | 1.01 | 0.78 |
| Child care Type | | | | |
| Center-based | | <i>Reference</i> | | |
| Family-based/Home-based | 1.25 | 0.87 | 1.80 | 0.22 |
| Director/Owner (of center-based) | | | | |
| No | | <i>Reference</i> | | |
| Yes | 0.95 | 0.72 | 1.25 | 0.73 |
| Personal COVID-19 Health Precautions | | | | |
| Masks/Handwashing/Distancing | 1.16 | 0.90 | 1.50 | 0.24 |
| Avoiding Social Interactions | 1.27 | 1.13 | 1.44 | 0.001 |
| Avoiding High-Risk Situations & Travel | 0.72 | 0.62 | 0.84 | <0.001 |
| County COVID-19 Cumulative Deaths | | | | |
| Low | | <i>Reference</i> | | |
| Moderate | 1.13 | 0.83 | 1.54 | 0.45 |
| High | 1.60 | 1.19 | 2.15 | 0.002 |
| County Median Household Income | | | | |
| Low | | <i>Reference</i> | | |
| Moderate | 1.26 | 0.96 | 1.65 | 0.10 |
| High | 1.03 | 0.76 | 1.41 | 0.84 |
| Interaction Effects | | | | |
| Exposure * Child Care Type | 0.85 | 0.54 | 1.34 | 0.49 |

Note. State where child care program is located (all 50 states, plus District of Columbia and Puerto Rico) was entered in the logistic regression analysis, but not reflected in the table.

¹ No cases self-identified as non-binary gender, dropping this value from analysis.

Table 7. Adjusted Case-Controlled Conditional Logistic Regression Analysis Predicting Case (N=427) Versus Matched Controls (N=21,350)

| | OR | 95% CI | | P Value |
|--|-----------|---------------|------|----------------|
| Exposure to Child care | 0.94 | 0.73 | 1.21 | 0.64 |
| Child care Type (Family/Home-based) | 1.59 | 1.14 | 2.23 | <0.01 |
| Interaction (Child care Type * Exposure) | 0.82 | 0.53 | 1.26 | 0.37 |

Note. Cases propensity score matched to controls using 1:50 random matching without replacement using caliper width 0.2*SD of the logit function. Values used in the propensity score include age, race, ethnicity, gender, director status, county-level COVID-19 death rate, county-level median household income, personal COVID-19 health precaution measures (3 continuous variables), and state where child care program is located (indicator variables).

COVID-19 Transmission in US Child Care Programs

Walter S. Gilliam, Aryn A. Malik, Mehr Shafiq, Madeline Klotz, Chin Reyes, John Eric Humphries, Thomas Murray, Jad A. Elharake, David Wilkinson and Saad B. Omer
Pediatrics originally published online October 14, 2020; originally published online October 14, 2020;

Updated Information & Services

including high resolution figures, can be found at:
<http://pediatrics.aappublications.org/content/early/2020/10/16/peds.2020-031971.citation>

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
<http://www.aappublications.org/site/misc/Permissions.xhtml>

Reprints

Information about ordering reprints can be found online:
<http://www.aappublications.org/site/misc/reprints.xhtml>

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®



PEDIATRICS[®]

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

COVID-19 Transmission in US Child Care Programs

Walter S. Gilliam, Aryn A. Malik, Mehr Shafiq, Madeline Klotz, Chin Reyes, John Eric Humphries, Thomas Murray, Jad A. Elharake, David Wilkinson and Saad B. Omer
Pediatrics originally published online October 14, 2020; originally published online October 14, 2020;

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/early/2020/10/16/peds.2020-031971.citation>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 345 Park Avenue, Itasca, Illinois, 60143. Copyright © 2020 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN[®]

